

Banded karyotype of the Konya wild sheep (*Ovis orientalis anatolica* Valenciennes, 1856) from Turkey

Atilla Arslan¹, Jan Zima²

1 Department of Biology, Faculty of Science, Selcuk University, TR-42031 Konya, Turkey **2** Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, CZ-603 65 Brno, Czech Republic

Corresponding author: Jan Zima (jzima@brno.cas.cz)

Academic editor: V. Kuznetsova | Received 18 February 2010 | Accepted 14 April 2010 | Published 30 June 2011

Citation: Arslan A, Zima J (2011) Banded karyotype of the Konya wild sheep (*Ovis orientalis anatolica* Valenciennes, 1856) from Turkey. *Comparative Cytogenetics* 5(2): 81–89. doi: 10.3897/CompCytogen.v5i2.1151

Abstract

The karyotype, C-banding, and nucleolar organizer regions (NORs) of eight specimens of Konya wild sheep from Turkey were examined. The complement included six large metacentric autosomes, 46 acrocentric autosomes of decreasing size, a medium-sized acrocentric X chromosome, and a small bi-armed Y chromosome (the diploid chromosome number $2n=54$, the number of autosomal arms $NFa=58$, the number of chromosome arms $NF=61$). G-banding allowed reliable identification of all the chromosome pairs and the pairing of homologous elements. All the autosomes possessed distinct centromeric or pericentromeric C-positive bands. The X chromosome had a pericentromeric C-positive band, and the Y chromosome was entirely C-heterochromatic. The NORs were located in the terminal regions of the long arms of three metacentric and two acrocentric autosomes. The karyotype of the Konya wild sheep and its banding patterns are quite similar to chromosome complement reported in domestic sheep and European mouflon.

Keywords

chromosomes, *Ovis*, sheep domestication, Turkey

Introduction

The systematics of the genus *Ovis* (Linnaeus, 1758) is rather complicated and there are different opinions in respect of the number of species and the actual species borders (Nadler et al. 1973a, b, Shackleton et al. 1997, Grubb 2005). One of the main problems is the origin and taxonomic position of the domestic sheep. For the domestic sheep a large number of possible ancestral species and subspecies exist, and several Eurasian wild sheep

taxa have been proposed as ancestors of domestic sheep or are believed to have contributed to specific breeds. Moufflons in the Mediterranean islands [*O. musimon* (Pallas, 1811); *O. ophion* Blyth, 1841] introduced to various parts of Europe are generally evaluated as feral populations of ancient domestic stocks (e.g. Hiendleder et al. 1998a, b). Phylogenetic analyses on mtDNA restriction fragment length polymorphism of Eurasian breeds of domestic sheep demonstrated two maternal origins among modern sheep breeds and indicated wild ancestors different from the previously considered urial (*O. vignei* Blyth, 1841) and argali sheep [*O. ammon* (Linnaeus, 1758)] (Hiendleder et al. 1998b). In a subsequent study, Hiendleder et al. (2002) confirmed two well-separated mtDNA lineages (European and Asian) suggesting domestication from two distinct wild populations.

The domestic sheep and its wild ancestors are sometimes considered conspecific, in which case the name *O. orientalis* has generally been used, although some authors use the name *O. aries* for both wild species and its domestic descendants (Grubb 2005). Likely candidates for wild ancestors of the European lineage of domestic sheep are populations found in Turkey and western Iran. At the species or subspecies level, these populations are currently referred to *O. orientalis*, *O. orientalis* Gmelin, 1774, *O. gmelinii* Blyth, 1841, *O. anatolica* Valenciennes, 1856, *O. ophion* or *O. musimon* (Kaya et al. 2000, 2004, Kryštufek and Vohralík 2009). The Konya wild sheep is a westernmost taxon of extant wild sheep endemic to Central Anatolia in Turkey, and it is usually referred to *Ovis orientalis anatolica* (Kryštufek and Vohralík, 2009) or *O. gmelinii anatolica* (Kaya et al., 2000). The only population is currently living in the area east of Konya (Bozdağ Protection of Wildlife). The Konya wild sheep was widespread 50 years ago, but the numbers have decreased rapidly because of harsh weather conditions, predator pressure, and hunting. The most severe population decline appeared in 1960s and 1970s when the population size of 30–40 individuals was reported (Kaya 1991, Kaya et al. 2000, Sezen et al. 2004). The abundance has later increased owing to protection, and the current numbers are estimated at several hundreds of heads (600 individuals reported in 2006; Yalçın et al., 2010). The apparently last surviving population inhabits a large protected area surrounded with a fence providing open steppe habitats in altitude ranging between 1000–1735 m above sea level. The Konya wild sheep is not phenotypically akin to the European or Mediterranean moufflons, because it displays the type of horn configuration typical of Asiatic wild sheep (Yalçın et al. 2010).

Chromosome research has contributed significantly to understanding of the evolutionary divergence among sheep taxa, and different chromosome numbers were found in individual geographic populations. The European mouflon and wild sheep from western Palaearctic (*O. orientalis*) have the same diploid number of 54 chromosomes as the domestic sheep. In the Iranian populations of urial sheep the diploid number of 58 chromosomes was recorded, and hybrid zones with polymorphic karyotype were found in the areas of contact with *O. orientalis* (Nadler et al. 1973a, Valdez et al. 1978). The argali sheep possess 56 chromosomes in their complements, $2n=52$ was found in *O. nivicola* Eschscholtz, 1929, and $2n=54$ in both the American species (Vorontsov et al. 1972a, b, Nadler et al. 1973a, b, Korobitsyna et al. 1974, Bunch et al. 1976, 1998,

2000, Lyapunova et al. 1997). The chromosomal findings support the view that wild sheep of the urial or argali types did not contribute to domestication of the sheep. However, despite the differences in chromosome number, different species of the genus can hybridize in captivity (Vorontsov et al. 1972b, Nadler et al. 1973b).

The karyotype of the Konya wild sheep was studied by Kaya et al. (2004), and the diploid number of 54 chromosomes was ascertained. Only a description of the basic karyotype characteristics was given, without any illustrations. Subsequently, Kırıkçı et al. (2003) reported the G-banding pattern of chromosomes of this population. In the present paper, we provide detailed description of the karyotype of the Konya wild sheep, a possible ancestor of domestic sheep, using various chromosome banding techniques.

Material and methods

We examined eight specimens of Konya wild sheep including four females and four males. All the studied animals originated from Bozdağ in the Konya province, Turkey (Fig. 1). Chromosomal preparations were made using peripheral blood culture for 72 hr at 37°C in medium supplemented with 15% fetal calf serum and 0.024 % phytohaemagglutinin (Freshney 1990). The chromosome slides were conventionally stained with Giemsa. G-banding by trypsin treatment stained with Giemsa (GTG) was performed (Seabright 1971). Constitutive heterochromatin and nucleolus organizer regions (NORs) were detected with C-banding (Sumner 1972) and Ag-NOR staining (Howell and Black 1980), respectively.




Figure 1. Collecting locality in the province of Konya, Anatolia, Turkey (●).




Figure 2. Conventionally stained karyotype of the Konya wild sheep with $2n=54$. There are three pairs of large metacentric autosomes and 23 pairs of acrocentric autosomes. The X chromosome is acrocentric and the Y chromosome metacentric.




Figure 3. G-banded karyotype of the Konya wild sheep. All the chromosomes in the complement can be identified and differentiated according to their G-banding pattern.

Results and discussion

The karyotype contained 54 chromosomes (Fig. 2) including three large metacentric autosomal pairs (no. 1–3) and 23 acrocentric autosomal pairs of decreasing size (nos. 4–26; NFa=58). Tiny short arms were observed in most of the acrocentric autosomes. The X chromosome was a large acrocentric with a distinct short arm, whereas the Y chromosome was metacentric and the smallest element of the complement (NF=61). All the autosomes and both the sex chromosomes were reliably identified on the basis of their unique G-banding patterns (Fig. 3). The C-banded karyotype of Konya wild sheep is illustrated in Fig. 4. All the autosomes possessed distinct and large centromeric C-positive bands which usually extended in the neighbouring pericentromeric areas. The X chromosome had a pericentromeric C-positive band, and the Y chromosome was entirely heterochromatic. Silver-nitrate staining revealed that NORs were localized in the distal position on the long arms of three large metacentric (no. 1–3) and two acrocentric (nos. 4 and 8) autosomes (Fig. 5). All the NORs were observed in both the homologous chromosomes, except those of the pair no. 8 in which a heterozygous condition with the signal present in only one homologue was recorded in four specimens.

The obtained karyotypic data confirm possible close relationships between the Konya wild sheep and domestic sheep. The diploid number of chromosomes and their morphology expressed in the number of chromosomal arms and the proportion of the metacentric and acrocentric autosomes are the same in both complements. The detailed structure of banded chromosomes of the Konya wild sheep appears quite similar in comparison with the standard chromosomal complement of the domestic sheep (cf. Ansari et al. 1999; ISCNDB 2000). The G-banding pattern observed in individual autosomes and the sex chromosomes is apparently identical between the karyotype of the Konya wild sheep and domestic sheep. The amount of pericentromeric C-heterochromatin is noticeably large in both the complements, and the Ag-NOR sites are localized in the same homologous chromosomes. Heteromorphism in the number of active NORs found in the acrocentric autosomal pair no. 8 may have resulted from the small sample size used in this study.

A considerable degree of similarity can also be demonstrated by comparison of our results with previously reported differentially stained chromosomes of various wild sheep taxa (Nadler et al. 1973b, Valdez et al. 1978, Bunch et al. 1998, 2000, Kırıkçı et al. 2003). Banding pattern in chromosomes and the distribution of NORs are also quite similar to that reported for European mouflon (Slavíčková et al. 1985). We may conclude that the chromosomal data support the possible position of the Konya wild sheep as one of the ancestors of domestic sheep. The surviving population should be subjected to further extensive genetical research, and its strict protection should be maintained.



Figure 4. C-banded karyotype of the Konya wild sheep. Distinct pericentromeric dark bands are apparent in all the autosomes and the X chromosome. The Y chromosome is stained entirely positively.




Figure 5. Silver-stained karyotype and the distribution of NORs in the Konya wild sheep. The NORs are localized in telomeric areas of the long arms of the three metacentric autosomal pairs (nos. 1, 2, 3) and two acrocentric autosomal pairs (nos. 4, 8). The heterozygous condition of the NORs in the acrocentric pair no. 8 was found consistently in various cells and different examined individuals.

Acknowledgments

We are obliged to prof. J. Rubeš for his kind advice. This research was supported by the Turkey Nature Protection Head Office of National Parks, and the Ministry of Environment and Forests of Turkey. J.Z. was supported by the Czech Ministry of Education (LC 06073). We are obliged to anonymous reviewers and particularly to the editor for useful comments on the earlier versions of the manuscript.

References

- Ansari HA, Bosma AA, Broad TE, Bunch TD, Long SE, Mahrer DW, Pearce PE, Popescu CP (1999) Standard G-, Q-, and R-banded ideograms of the domestic sheep (*Ovis aries*): homology with cattle (*Bos taurus*). Report of Committee for the standardization of the sheep karyotype. *Cytogenetics and Cell Genetics* 87: 134–142. doi:10.1159/000015380
- Bunch TD, Foote WC, Spillett JJ (1976) Translocations of acrocentric chromosomes and their implications in the evolution of sheep (*Ovis*). *Cytogenetics and Cell Genetics* 17: 122–136. doi:10.1159/000130704
- Bunch TD, Vorontsov NN, Lyapunova EA, Hoffman RS (1998) Chromosome number of Severtsov's sheep (*Ovis ammon severtsovi*): G-banded karyotype comparisons within *Ovis*. *Journal of Heredity* 89: 266–269. doi:10.1093/jhered/89.3.266
- Bunch TD, Wang S, Valdez R, Hoffman RS, Zhang Y, Liu A, Lin S (2000) Cytogenetics, morphology and evolution of four subspecies of the giant sheep argali (*Ovis ammon*) of Asia. *Mammalia* 64: 199–207. doi:10.1515/mamm.2000.64.2.199
- Freshney RI (1990) Culture of animal cells. A manual of basic technique. 2nd ed. Wiley Liss, New York.
- Grubb P (2005) Order Artiodactyla. Wilson DE, Reeder DM (Eds). In: *Mammal species of the world. A taxonomic and geographical reference. Volume 1. 3rd Ed.* The Johns Hopkins University Press, Baltimore, 637–722.
- Hiendleder S, Kaup B, Wassmuth R, Janke A (2002) Molecular analysis of wild and domestic sheep questions current nomenclature and provides evidence for domestication from two different subspecies. *Proceedings of the Royal Society London B* 269: 893–904. doi:10.1098/rspb.2002.1975
- Hiendleder S, Lewalski H, Wassmuth R, Janke A (1998a) The complete mitochondrial DNA sequence of the domestic sheep (*Ovis aries*) and comparison with the other major ovine haplotypes. *Journal of Molecular Evolution* 47: 441–448. doi:10.1007/PL00006401
- Hiendleder S, Mainz K, Plante Y, Lewalski H (1998b) Analysis of mitochondrial DNA indicates that domestic sheep are derived from two different ancestral maternal sources. No evidence for contribution from urial and argali sheep. *Journal of Heredity* 89: 113–120. doi:10.1093/jhered/89.2.113
- Howell WM, Black DA (1980) Controlled silver staining of nucleolar organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 36: 1014–1015. doi:10.1007/BF01953855

- ISCNDB (2000) International system for chromosome nomenclature of domestic bovinds. *Cytogenetics and Cell Genetics* 92: 283–299.
- Kaya MA (1991) The morphology, increase in weight, development of teeth and horns in Anatolian wild sheep, *Ovis orientalis anatolica* Valenciennes, 1856, living in Bozdağ Konya. *Turkish Journal of Zoology* 15: 135–149. [In Turkish, with English summary]
- Kaya MA, Arihan O, Bilgin CC (2000) General characteristics and taxonomy of the Turkish mouflon *Ovis gmelini anatolica* Valenciennes, 1856. In: Náhlik A, Uloth W (Eds) International Mouflon Symposium, October 27–29, 2000. Sopron, Hungary 92–95.
- Kaya MA, Bunch TD, Konuk M (2004) On Konya wild sheep, *Ovis orientalis anatolica*, in the Bozdag protected area. *Mammalia* 68: 229–232. doi:10.1515/mamm.2004.022
- Kırıkçı K, Zamani A, Durakbaşı G (2003) A study on the chromosomes of Konya wild sheep (*Ovis orientalis* spp.): Case report. *Turkish Journal of Veterinary and Animal Sciences* 27: 281–283. [In Turkish, with English summary]
- Korobitsyna KV, Nadler CF, Vorontsov NN, Hoffmann RS (1974) Chromosomes of the Siberian snow sheep, *Ovis nivicola*, and implications concerning the origin of amphiberian wild sheep (subgenus *Pachyceros*). *Quaternary Research* 4: 235–245. doi:10.1016/0033-5894(74)90013-1
- Kryštufek B, Vohralík V (2009) Mammals of Turkey and Cyprus. Rodentia II. Cricetinae, Muridae, Spalacidae, Calomyscidae, Capromyidae, Hystricidae, Castoridae. Knjižnica Annales Majora, Koper.
- Lyapunova EA, Vorontsov NN, Hoffmann R (1997) Chromosomal sets and taxonomical position of *Ovis ammon severtzovi* Nasonov. *Zoologicheskii Zhurnal* (Zoological journal). 76: 1083–1093. [in Russian, with English summary]
- Nadler CF, Korobitsyna KV, Hoffmann RS, Vorontsov NN (1973a) Cytogenetic differentiation, geographic distribution and domestication of Palaearctic sheep (*Ovis*). *Zeitschrift für Säugetierkunde* 38: 109–125.
- Nadler CF, Hoffmann RS, Woolf A (1973b) G-band patterns as chromosomal markers, and the interpretation of chromosomal evolution in wild sheep (*Ovis*). *Experientia* 29: 117–119. doi:10.1007/BF01913288
- Seabright M (1971) A rapid banding technique for human chromosomes. *Lancet* 2: 971–972. doi:10.1016/S0140-6736(71)90287-X
- Sezen Z, Akaçakaya HR, Bilgin CC (2004). Turkish mouflon (*Ovis gmelini anatolica*) in Central Anatolia: Population viability under scenarios of harvesting for trophy. In: Akcakaya HR, Burgman MA, Kindvall O, Wood CC, Sjögren-Gulve P, Hatfield JS, McCarthy MA (Eds). *Species Conservation and Management: Case Studies*. Section VI, Chapter 41, Mammals. Oxford University Press, UK, 459–467.
- Shackleton DM, the IUCN/SSC Caprinae Specialist Group (1997) Wild sheep and goats and their relatives. Status survey and conservation action plan for Caprinae. IUCN, Gland and Cambridge.
- Slavičková M, Havránková J, Zima J, Černý M (1985) The karyotype and chromosome banding in mouflon in the Czech Republic. *Veterinary Medicine (Czech)* 30: 681–686.
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Experimental Cell Research* 75: 304–306. doi:10.1016/0014-4827(72)90558-7

- Valdez R, Nadler CF, Bunch TD (1978) Evolution of wild sheep in Iran. *Evolution* 32: 56–72. doi:10.2307/2407410
- Vorontsov NN, Korobitsina KV, Nadler Ch, Hoffmann RS, Sapozhnikov GN, Gorelov YuK (1972a) Cytogenetic differentiation and species borders in true Palaearctic sheeps (*Ovis* s. str.). *Zoologicheskii Zhurnal* 51: 1109–1123. [In Russian, with English summary]
- Vorontsov NN, Korobitsyna KV, Nadler CF, Hoffmann R, Sapozhnikov GN, Gorelov JK (1972b) Chromosomi dikich baranov i proiskhozhdenie domashnikh ovets. *Priroda* (Moscow) 3: 74–82. [In Russian]
- Yalçın H, Kaya MA, Arslan A (2010) Comparative geometrical morphometries on the mandibles of Anatolian wild sheep (*Ovis gmelini anatolica*) and Akkaraman sheep (*Ovis aries*). *Kafkas Universitesi Veteriner Fakültesi Dergisi* 16: 55–61.

