

Chromosomal and molecular evidence for presence of *Polyommatus (Agrodiaetus) poseidon* (Lepidoptera, Lycaenidae) in Caucasus region

Vladimir A. Lukhtanov^{1,2}, Valentin V. Tikhonov³

1 Department of Karyosystematics, Zoological Institute of Russian Academy of Sciences, Universitetskaya nab. 1, 199034 St. Petersburg, Russia **2** Department of Entomology, Faculty of Biology, St. Petersburg State University, Universitetskaya nab. 7/9, 199034 St. Petersburg, Russia **3** North Caucasus Federal University, Pushkin str., 1, Stavropol 355009 Russia

Corresponding author: Vladimir A. Lukhtanov (lukhtanov@mail.ru)

Academic editor: S. Nokkala | Received 1 April 2015 | Accepted 10 April 2015 | Published 3 June 2015

<http://zoobank.org/6073BC1E-59EB-40B8-8AB5-EEE8449C9E70>

Citation: Lukhtanov VA, Tikhonov VV (2015) Chromosomal and molecular evidence for presence of *Polyommatus (Agrodiaetus) poseidon* (Lepidoptera, Lycaenidae) in Caucasus region. *Comparative Cytogenetics* 9(2): 249–255. doi: 10.3897/CompCytogen.v9i2.5020

Abstract

We show how combination of chromosomal and molecular markers can be applied for proper species identification in *Agrodiaetus* Hübner, 1822 blue butterflies. Using this approach we provide first evidence for presence of *P. (A.) poseidon* (Herrich-Schäffer, [1851]) in Georgia.

Keywords

Agrodiaetus, chromosome number, *COI*, karyotype, Lycaenidae, *Polyommatus*

Introduction

The blue butterfly subgenus *Agrodiaetus* Hübner, 1822 belongs to the genus *Polyommatus* Latreille, 1804 (Talavera et al. 2013). In the last years, this group become a model system for study of speciation and chromosome evolution (Lukhtanov et al. 2015, Verшинina et al. 2015). Despite this, its taxonomy is still poorly elaborated and identification of individual species is difficult due to their morphological similarity. Species within the subgenus are mostly uniform and exhibit few differences in

characters traditionally used in classification, such as wing pattern and/or aspects of the male and female genitalia (Lukhtanov et al. 2006, Vila et al. 2010). The genus was estimated to have originated very recently (Kandul et al. 2004) and, thus, many *Agrodiaetus* species may have not had sufficient time to acquire extensive genetic differences. In particular, *COI* barcode gap is low or even absent between numerous closely related species of *Polyommatus* (*Agrodiaetus*) (Wiemers and Fiedler 2007). In opposite to majority of other butterflies and moths (Lukhtanov 2014), many *Agrodiaetus* species have evolved distinctive karyotypes. They show one of the highest interspecific karyotypic diversities known in the animal kingdom with haploid chromosome numbers ranging from $n = 10$ to $n = 134$ (Lukhtanov et al. 2005). Therefore, karyotypic features provide important identification characters for many described species that are virtually indistinguishable by their morphology. However, it should be noted that in few cases the chromosome number may be identical in different species (see Results and Discussion).

Here we show how combination of chromosomal and molecular markers can be applied for proper species identification in *Agrodiaetus*. Using this approach we provide first evidence for presence of *P. (A.) poseidon* (Herrich-Schäffer, [1851]) in Georgia.

Material and methods

The samples used for molecular and chromosomal analysis were collected in Georgia (Akhaltsikhe, 41.60N, 43.06E, 1000 m alt., 18 July 2014, V. Lukhtanov et V. Tikhonov leg., samples 2014VL56, 2014VL57, 2014VL58, 2014VL62, 2014VL63, 2014VL64, 2014VL65, 2014VL68, 2014VL69, 2014VL70). The methods of DNA sequencing, chromosomal analysis and phylogenetic inference were described previously (Lukhtanov and Dantchenko 2002a, Lukhtanov et al. 2008, 2014, Vershinina and Lukhtanov 2010, Przybyłowicz et al. 2014). Additional samples of *Polyommatus* belonging to *P. (A.) poseidon* species complex (Kandul et al. 2007) were used for comparison.

Results and discussion

The species *P. (A.) poseidon* (= *Lycaena poseidon* var. *mesopotamica* Staudinger, 1892, synonymized with *P. poseidon* by Schurian et. 1992) is known to be an endemic of the Middle East sporadically distributed from Kütahya in West Turkey to Artvin in North-East Turkey (Hesselbarth et al. 1995). Phenotypically similar, but chromosomally distinct species *P. (A.) putnami* (Lukhtanov & Dantchenko, 2002) was described from East Turkey (provinces Erzurum and Ağrı) (Lukhtanov and Dantchenko 2002b). The last taxon is allopatric in distribution with *P. (A.) poseidon* and differs from *P. poseidon* by chromosome number and karyotype structure (Lukhtanov and Dantchenko 2002b). *P. (A.) poseidon* has relatively low haploid chromosome number (from $n=19$ on the south and east of the distributional range to $n=21$ in the north), all the chro-

mosomes form a gradient size row with no especially large or small chromosomes (de Lesse 1963, Kandul and Lukhtanov 1997). Chromosome numbers $n=22$ and $n=23$ were also found in the northern population as intraindividual occasional deviations from the basic $n=21$ (de Lesse 1963). *P. (A.) putnami* has higher chromosome numbers (from $n=24$ to $n=27$, with $n=26$ as a distinct mode). Its karyotype is asymmetrical and includes chromosomes of two distinct classes: class of large chromosomes and class of small chromosomes (Lukhtanov and Dantchenko 2002b). Currently *P. (A.) putnami* is treated as a distinct species (Lukhtanov and Dantchenko 2002b, Wiemers 2003, Wiemers and Fiedler 2007) or a subspecies of *P. (A.) poseidon* (Tshikolovets 2011).

The taxon *P. (A.) deebe* (Larsen, 1974) discovered in Lebanon and Syria is often considered as a subspecies of *P. (A.) poseidon* (e.g. Tshikolovets 2011), however, it differs in chromosome number ($n=17$, Larsen 1975) and may represent a different species (Eckweiler and Häuser 1997). The taxon *P. (A.) damocles krymaeus* (Sheljuzhko, 1928) was also considered as subspecies of *P. (A.) poseidon* (Hesselbarth et al. 1995), however, with respect to mitochondrial genes *COI* and *COII* it is very distant from *P. (A.) poseidon* and was shown to be a subspecies of *P. (A.) damocles* (Herrich-Schäffer, [1844]) (Lukhtanov et al. 2005, Kandul et al. 2007).

Males of *P. (A.) poseidon* have plesiomorphic (Kandul et al. 2004, Lukhtanov et al. 2005) blue colouration of the upper side of the wings with no specific morphological characters. Therefore their morphological discrimination from phenotypically similar *P. (A.) caeruleus* (Staudinger, 1871), *P. (A.) damocles* and *P. (A.) damonides* (Staudinger, 1899) is difficult. With respect to *COI* barcodes, *P. (A.) poseidon* is indistinguishable from *P. (A.) hopfferi* (Herrich-Schäffer, [1851]) and *P. (A.) putnami* (Wiemers & Fiedler, 2007). As it was stated above, the chromosome number varies within *P. (A.) poseidon* (de Lesse 1963, Kandul and Lukhtanov 1997, Lukhtanov and Dantchenko 2002b) and thus overlap with chromosome numbers found in *P. (A.) elbursicus* (Forster, 1956), *P. (A.) cyaneus* (Staudinger, 1899), *P. (A.) ectabanensis* (de Lesse, 1963), *P. (A.) hamadanensis* (de Lesse, 1959), *P. (A.) alcestis* (Zerny, 1932), *P. (A.) altivagans* (Forster, 1956), *P. (A.) mithridates* (Staudinger, 1878), *P. (A.) shirkuhensis* ten Hagen




Figure 1. *Polyommatus (Agrodiaetus) poseidon* from Akhaltsikhe, Georgia. **a** male, upperside **b** male, underside.




Figure 2. Bayesian tree of the species close to *Polyommatus (Agrodiaetus) poseidon* inferred from *COI* sequences. Posterior probability values >50% are shown.




Figure 3. Male karyotype of *Polyommatus (Agrodiaetus) poseidon* from Georgia. **a** sample 2014VL57, metaphase I, n = 19 **b** sample 2014VL62, metaphase II, n = 19. Bar = 10 μm.

et Eckweiler, 2001 and *P. (A.) pierceae* (Lukhtanov & Dantchenko, 2002) (Kandul et al. 2007, Lukhtanov et al. 2014).

A population of blue butterflies which were morphologically similar to *P. (A.) poseidon* (Fig. 1) was discovered near Akhaltikhe in Georgia in 2013 by V.Tikhonov and I. Kostyuk. In 2014 the locality was visited again in order to collect material available for molecular and chromosomal study. Molecular analysis of this material revealed

that *COI* barcodes were completely identical or nearly identical (barcode gap from 0 to 0.6%) in population from Akhaltsikhe and other populations of *P. (A.) poseidon* and *P. (A.) putnami* (Fig. 2).

The haploid chromosome number $n=19$ was found in MI and MII cells of three studied individuals (2014VL57, 2014VL58, 2014VL62) (Fig. 3). All chromosome elements formed a gradient size row. The karyotype contained no exceptionally large or small chromosomes. In this respect, the population from Akhaltsikhe is indistinguishable from populations of *P. (A.) poseidon* from Amasya (de Lesse 1963) and Artvin (Kandul and Lukhtanov 1997), but differs from *P. (A.) putnami* ($n=26$) (Lukhtanov and Dantchenko 2002b).

Thus, although in the studied case neither the DNA barcodes nor chromosomal numbers are species-specific characters, their combination clearly indicates that the population from Akhaltsikhe should be identified as *P. (A.) poseidon*. This is the first evidence of *P. (A.) poseidon* for Georgia and for Caucasus region at whole.

Acknowledgements

We thank Igor Kostyuk (Kiev State University) for participation in the trip to Georgia. The financial support for this study was provided by the grant from the Russian Science Foundation N 14-14-00541 to Zoological Institute of the Russian Academy of Sciences.

References

- de Lesse H (1963) Variation chromosomique chez les *Agrodiaetus* [Lep. Lycaenidae]. Revue Française d'Entomologie 30(3): 182–189.
- Eckweiler W, Häuser C (1997) An illustrated checklist of *Agrodiaetus* Hübner, 1822, a subgenus of *Polyommatus* Latreille, 1804 (Lepidoptera, Lycaenidae). Nachrichten des Entomologischen Vereins Apollo, Supplement 16: 113–168.
- Hesselbarth G, Van Oorschot H, Wagener S (1995) Die Tagfalter der Türkei unter Berücksichtigung der angrenzenden Länder. Vol. 1-3. Selbstverlag S. Wagener, Bocholt, 1354 pp.
- Kandul NP, Lukhtanov VA (1997) Karyotype variability and systematics of blue butterflies of the species groups *Polyommatus (Agrodiaetus) poseidon* and *Polyommatus (Agrodiaetus) dama* (Lepidoptera, Lycaenidae). Zoologicheskij Zhurnal 76(1): 63–69.
- Kandul NP, Lukhtanov VA, Dantchenko AV, Coleman JWS, Sekercioglu CH, Haig D, Pierce NE (2004) Phylogeny of *Agrodiaetus* Hübner 1822 (Lepidoptera: Lycaenidae) inferred from mtDNA sequences of *COI* and *COII* and nuclear sequences of *EF1- α* : Karyotype diversification and species radiation. Systematic Biology 53(2): 278–298. doi: 10.1080/10635150490423692
- Kandul NP, Lukhtanov VA, Pierce NE (2007) Karyotypic diversity and speciation in *Agrodiaetus* butterflies. Evolution 61(3): 546–559. doi: 10.1111/j.1558-5646.2007.00046.x

- Larsen TB (1975) Chromosome numbers and notes on testicular morphology of some Lebanese Rhopalocera (Insecta: Lepidoptera). *Entomologica Scandinavica* 6(3-4): 253–260. doi: 10.1163/187631275X00091
- Lukhtanov VA (2014) Chromosome number evolution in skippers (Lepidoptera, Hesperiiidae). *Comparative Cytogenetics* 8(4): 275–291 doi: 10.3897/CompCytogen.v8i4.8789
- Lukhtanov VA, Dantchenko AV (2002a) Principles of highly ordered metaphase I bivalent arrangement in spermatocytes of *Agrodiaetus* (Lepidoptera). *Chromosome Research* 10(1): 5–20. doi: 10.1023/A:1014249607796
- Lukhtanov VA, Dantchenko AV (2002b) Descriptions of new taxa of the genus *Agrodiaetus* Hübner, [1822] based on karyotype investigation (Lepidoptera, Lycaenidae). *Aralanta* 33(1/2): 81–107, 224–225.
- Lukhtanov VA, Kandul NP, Plotkin JB, Dantchenko AV, Haig D, Pierce NE (2005) Reinforcement of pre-zygotic isolation and karyotype evolution in *Agrodiaetus* butterflies. *Nature* 436(7049): 385–389. doi: 10.1038/nature03704
- Lukhtanov VA, Shapoval NA, Anokhin BA, Saifitdinova AF, Kuznetsova VG (2015) Homoploid hybrid speciation and genome evolution via chromosome sorting. *Proceedings of the Royal Society B: Biological Sciences* 282(1807): 20150157. doi: 10.1098/rspb.2015.0157
- Lukhtanov VA, Shapoval NA, Dantchenko AV (2008) *Agrodiaetus shahkubensis* sp. n. (Lepidoptera, Lycaenidae), a cryptic species from Iran discovered by using molecular and chromosomal markers. *Comparative Cytogenetics* 2(2): 99–114.
- Lukhtanov VA, Shapoval NA, Dantchenko AV (2014) Taxonomic position of several enigmatic *Polyommatus* (*Agrodiaetus*) species (Lepidoptera, Lycaenidae) from Central and Eastern Iran: insights from molecular and chromosomal data. *Comparative Cytogenetics* 8(4): 313–322. doi: 10.3897/CompCytogen.v8i4.8939
- Lukhtanov VA, Vila R, Kandul NP (2006) Rearrangement of the *Agrodiaetus dolus* species group (Lepidoptera, Lycaenidae) using a new cytological approach and molecular data. *Insect Systematics and Evolution* 37(3): 325–334. doi: 10.1163/187631206788838563
- Przybyłowicz Ł, Lukhtanov V, Lachowska-Cierlik D (2014) Towards the understanding of the origin of the Polish remote population of *Polyommatus* (*Agrodiaetus*) *ripartii* (Lepidoptera: Lycaenidae) based on karyology and molecular phylogeny. *Journal of Zoological Systematics and Evolutionary Research* 52(1): 44–51. doi: 10.1111/jzs.12040
- Schurian KG, van Oorschoot H, van den Brink H (1992) *Polyommatus* (*Agrodiaetus*) *poseidon* (H.-S.) und *Polyommatus* (*Agrodiaetus*) *theresia* sp. nov. aus der Türkei (Lepidoptera: Lycaenidae). *Nachrichten des Entomologischen Vereins Apollo* 12(4): 217–232.
- Talavera G, Lukhtanov VA, Pierce NE, Vila R (2013) Establishing criteria for higher-level classification using molecular data: the systematics of *Polyommatus* blue butterflies (Lepidoptera, Lycaenidae). *Cladistics* 29: 166–192. doi: 10.1111/j.1096-0031.2012.00421.x
- Tshikolovets V (2011) Butterflies of Europe & Mediterranean Area. Tshikolovets Publication, Pardubice, Czech Republic.
- Vershinina AO, Anokhin BA, Lukhtanov VA (2015) Ribosomal DNA clusters and telomeric (TTAGG)_n repeats in blue butterflies (Lepidoptera, Lycaenidae) with low and high chromosome numbers. *Comparative Cytogenetics* 9(2): 161–171. doi: 10.3897/CompCytogen.v9i2.4715
- Vershinina AO, Lukhtanov VA (2010) Geographical distribution of the cryptic species *Agrodiaetus alcestis alcestis*, *A. alcestis karacetinae* and *A. demavendi* (Lepidoptera, Lycaenidae) revealed by cytogenetic analysis. *Comparative Cytogenetics* 4(1): 1–11. doi: 10.3897/compcytogen.v4i1.21

- Vila R, Lukhtanov VA, Talavera G, Gil-T F, Pierce NE (2010) How common are dot-like distribution ranges? Taxonomical oversplitting in Western European *Agrodiaetus* (Lepidoptera, Lycaenidae) revealed by chromosomal and molecular markers. *Biological Journal of the Linnean Society* 101: 130–154. doi: 10.1111/j.1095-8312.2010.01481.x
- Wiemers M (2003) Chromosome differentiation and the radiation of the butterfly subgenus *Agrodiaetus* (Lepidoptera: Lycaenidae: *Polyommatus*) a molecular phylogenetic approach. Ph.D. Dissertation, University of Bonn, Bonn, Germany, 203 pp. <http://hss.ulb.uni-bonn.de/2003/0278/0278.htm>
- Wiemers M, Fiedler K (2007) Does the DNA barcoding gap exist? – a case study in blue butterflies (Lepidoptera: Lycaenidae). *Frontiers in Zoology* 4: 8. doi: 10.1186/1742-9994-4-8