

The contribution of cytogenetics and flow cytometry for understanding the karyotype evolution in three *Dorstenia* (Linnaeus, 1753) species (Moraceae)

Paulo Marcos Amaral-Silva¹, Wellington Ronildo Clarindo¹, Tatiana Tavares Carrijo²,
Carlos Roberto Carvalho³, Milene Miranda Praça-Fontes¹

1 Laboratório de Citogenética, Departamento de Biologia, Centro de Ciências Agrárias, Universidade Federal do Espírito Santo. CEP: 29.500-000 Alegre – ES, Brazil **2** Laboratório de Botânica, Departamento de Biologia, Centro de Ciências Agrárias, Universidade Federal do Espírito Santo. CEP: 29.500-000 Alegre – ES, Brazil **3** Laboratório de Citogenética e Citometria, Departamento de Biologia Geral, Centro de Ciências Biológicas e da Saúde, Universidade Federal de Viçosa. CEP: 36.570-000 Viçosa – MG, Brazil

Corresponding author: Milene Miranda Praça-Fontes (milenemiranda@yahoo.com.br)

Academic editor: J.R. Daviña | Received 29 September 2015 | Accepted 18 November 2016 | Published 22 January 2016

<http://zoobank.org/EBC66E81-3A04-479B-96F0-D51CC2354AA1>

Citation: Amaral-Silva PM, Clarindo WR, Carrijo TT, Carvalho CR, Praça-Fontes MM (2016) The contribution of cytogenetics and flow cytometry for understanding the karyotype evolution in three *Dorstenia* (Linnaeus, 1753) species (Moraceae). *Comparative Cytogenetics* 10(1): 97–108. doi: 10.3897/CompCytogen.v10i1.6719

Abstract

Chromosome morphometry and nuclear DNA content are useful data for cytotaxonomy and for understanding the evolutionary history of different taxa. However, the chromosome number is the only karyotype aspect reported for the species of *Dorstenia* so far. In this study, the nuclear genome size of *Dorstenia arifolia* (Lamarck, 1786), *Dorstenia bonijesu* (Carauta & C. Valente, 1983) and *Dorstenia elata* (Hooker, 1840) was evaluated and their karyotype morphometry accomplished, with the aim of verifying the potential of those parameters to understand evolutionary issues. Mean nuclear 2C value ranged from 2C = 3.49 picograms (pg) for *D. elata* to 2C = 5.47 pg for *D. arifolia*, a variation of ± 1.98 pg. Even though showing a marked difference in 2C value, the three species exhibited the same $2n = 32$. Corroborating the flow cytometry data, differences in chromosome morphology were found among the karyotypes of the species investigated. Based on this and the only phylogeny proposed for *Dorstenia* thus far, structural rearrangements are related to the karyotype variations among the three species. Besides, the karyological analysis suggests a polyploid origin of the *Dorstenia* species studied here.

Keywords

Cytogenetics, euploidy, karyotype, flow cytometry, chromosome structure

Introduction

The pantropical genus *Dorstenia* Linnaeus, 1753 comprises about 105 species distributed in Asia, Africa and Neotropical regions (Carauta 1978, Berg 2001). Thirty-nine species occur in Brazil (Romaniuc-Neto et al. 2015), representing the sections *Dorstenia* Linnaeus, 1753, *Emigdioa* Carauta, 1976 and *Lecanium* Fischer & Meyer, 1846. Carauta (1978) and Berg (2001) have contributed much of the knowledge about the morphology of Neotropical species, while the phylogenetic reconstructions using molecular data (Misiewicz and Zerega 2012) consolidated the hypothesis about the genus' monophyly. Despite the potential information of cytogenetic data for taxonomic (Stace 2000) and phylogenetic studies (Melo and Guerra 2003), this marker has been little explored in previous works on *Dorstenia*.

According to the current knowledge, the basic chromosome number in African species of *Dorstenia* is $x = 12$ and $x = 13$, while in American species $x = 14$, 15 and 16 (Berg 2001) are found. Fagerlind (1944) reported *Dorstenia mannii* Hooker f., 1871 as a tetraploid ($2n = 48$). In addition, intraspecific variation in chromosome number have been reported for subspecies of *Dorstenia psilurus* Welwitsch, 1869 ($2n = 26$ and $2n = 40$) (Misiewicz and Zerega 2012) and *Dorstenia elata* Hooker, 1840 ($2n = 26$ or 32) (Krause 1931, Hoen 1983, unpublished data). These data indicate that the evolution of the karyotype in *Dorstenia* involved euploidy and aneuploidy events. However, it is still necessary to invest efforts in confirming and understanding the chromosome changes reported for some species in previous studies. Considering that chromosome alterations are a significant mechanism of diversification and speciation in Angiosperms (Stace 2000, Peer et al. 2009, Weiss-Schneeweiss et al. 2013), the investigation of this aspect in *Dorstenia* can bring light to the knowledge of speciation processes in the genus.

Cytogenetic approaches, which regard the chromosome number, morphometric measurements and karyotype analysis, contribute to the understanding of evolutionary processes in plants (Shan et al. 2003). The knowledge of these aspects in related species helps to elucidate issues related to diversification of a taxonomic group (Clarindo et al. 2012). Morphometric analysis of the chromosomes is a way to determine the karyotype changes that occurred throughout evolution, the processes that led to the diversification, and the direction taken by evolution (Gao et al. 2012).

Numeric and structural chromosome rearrangements have been reported as triggers of karyotype changes in several plant taxa. Therefore, nuclear genome size variation occurs between phylogenetically related species due to these alterations (Bonifácio et al. 2012, Raskina et al. 2008). For this reason, nuclear DNA content measurement has been increasingly employed in systematic approaches using flow cytometry (FCM). In addition to its practicality and reproducibility, FCM is useful to reveal differences between taxa (Stace 2000), especially in groups of species that exhibit conserved chromosome number (Mabuchi et al. 2005).

The cytogenetic and FCM approaches in *Dorstenia* could provide relevant information to sections and species taxonomy, as well as contribute to understanding the evolutionary history of the genus. The main goal of this study was therefore to meas-

ure the nuclear 2C value, determine the chromosome number and characterize the karyotype of Neotropical species of *Dorstenia*: *D. arifolia* Lamarck, 1786, *D. bonijesu* Carauta & C. Valente, 1983 and *D. elata* Hooker, 1840.

Material and methods

Plant samples – Specimens of *D. elata*, *D. bonijesu* and *D. arifolia* (Fig. 1 a, c, e, respectively) were collected in an Atlantic Rainforest remnant located in the city of Castelo – ES, Brazil. Voucher specimens were included in the herbarium VIES: T.T. Carrijo 1516 (*D. arifolia*); T.T. Carrijo 1682 (*D. bonijesu*) and T.T. Carrijo 1618 (*D. elata*).

FCM – Nuclear suspensions were obtained by chopping (Galbraith et al. 1983) of leaf fragments (1 cm²) excised from each *Dorstenia* species (sample) and *Solanum lycopersicum* Linnaeus, 1753 (internal standard, 2C = 2.00 picograms – Praça-Fontes et al. 2011). Samples and standard leaf fragments were simultaneously chopped with razor blade in a Petri dish containing 0.5 ml OTTO-I nuclear extraction buffer (Otto 1990) supplemented with 2 mM dithiothreitol and 50 µg ml⁻¹ RNase (Praça-Fontes et al. 2011). Afterwards, 0.5 ml of the OTTO-I buffer was added, the suspensions were filtered through 30-µm nylon mesh, placed into microtube and centrifuged at 100 × g for 5 min. The precipitate was resuspended in 100 µl OTTO-I buffer and incubated for 10 min (Praça-Fontes et al. 2011). The nuclei suspensions were stained with 1.5 ml of OTTO-I:OTTO-II (1:2 – Otto 1990, Loureiro et al. 2006) solution supplemented with 2 mM dithiothreitol, 50 µg ml⁻¹ propidium iodide and 50 µg ml⁻¹ RNase (Praça-Fontes et al. 2011). The nucleus suspensions were kept in the dark for 30 min, then filtered through 20-µm nylon mesh. The samples were analyzed in a flow cytometer Partec PAS II/III (Partec GmbH, Germany). Histograms were analyzed with the Partec Flow Max software tools to measure nuclear DNA content. The genome size of the *Dorstenia* species was calculated according to the formula:

$$2C_D = \left(\frac{C1}{C2} \right) \cdot 2C_S$$

wherein:

- 2C_D: value of 2C DNA content (pg) of each *Dorstenia* species,
- C1: average G₀/G₁ peak channel of the *Dorstenia* species,
- C2: average G₀/G₁ peak channel of *S. lycopersicum*,
- 2C_S: value of 2C DNA content of *S. lycopersicum* (2.00 pg).

Cytogenetics – Stems with length of approximately 15 cm exhibiting one leaf pair were excised and disinfected with 1% NaOCl₂ solution for 15 min. These propagules were maintained in hydroponic system for rooting. The system was oxygenated by a compressor coupled to a plastic hose, which was immersed in H₂O. The roots were treated with 3 µM or 4 µM amiprofos-methyl (APM) for 16 h or 18 h at 4 °C. The roots were washed in dH₂O for 20 min, fixed in methanol:acetic acid solution (3:1)

and stored at $-20\text{ }^{\circ}\text{C}$ (Carvalho et al. 2007). After 24 h, the roots were washed in dH_2O and macerated in pectinase solution 1:40, 1:45, 1:50, 1:55 or 1:60 (enzyme: dH_2O) for 1 h 45 min or 2 h at $34\text{ }^{\circ}\text{C}$. The material was washed in dH_2O , fixed again and kept at $-20\text{ }^{\circ}\text{C}$ until use. Using the macerated root meristems, slides were prepared by cell dissociation and air-drying techniques, then placed onto a hot plate at $50\text{ }^{\circ}\text{C}$ (Carvalho et al. 2007). The slides were stained with 5% Giemsa (Merck®) for 8 min, washed twice in dH_2O , and air-dried. All slides were analyzed under a Nikon Eclipse Ci-S microscope (Nikon). The capture of metaphase images was performed using 100× objective and a CCD camera (Nikon Evolution™) coupled to a Nikon 80i microscope (Nikon).

Morphometric analysis – The chromosomes of three *Dorstenia* species were characterized as to the total length, length of the long and short arms, and classes. The chromosome class was determined as proposed by Levan et al. (1964) and reviewed by Guerra (1986): $r = \text{length of the long arm} / \text{length of the short arm}$. The asymmetry of the karyotype was evaluated using the method proposed by Zarco (1986), using the formulae:

$$A_2 = \frac{s}{\bar{X}}$$

where A_2 = interchromosomal asymmetry index, s = standard deviation, and \bar{X} = average length of the chromosomes.

Results

Flow cytometry measurements

The nuclear suspensions resulted in histograms with G_0/G_1 peaks showing coefficient of variation below 3.45%. Thus, the isolation and staining procedures provided suspensions containing isolated, intact and stoichiometrically stained nuclei. Using the histograms, the mean 2C nuclear DNA content of *Dorstenia* species was measured for the first time. The values were $2C = 3.49\text{ pg} \pm 0.0035$ ($1C = 1.71\text{ bp} \times 10^9$) for *D. elata*; $2C = 4.05\text{ pg} \pm 0.014$ ($1C = 1.98\text{ bp} \times 10^9$) for *D. bonijesu*; and $2C = 5.47\text{ pg} \pm 0.002$ ($1C = 2.67\text{ bp} \times 10^9$) for *D. arifolia* (Fig. 1 b, d, f). The mean value for *D. arifolia* was 36.20% higher than for *D. elata*, and 26.00% greater than for *D. bonijesu*. Besides, *D. bonijesu* showed a DNA content 13.83% higher than that of *D. elata*. Based on these values, an interspecific variation of the nuclear genome size was identified between the species.

Cytogenetics

The rooting of vegetative propagules occurred after 40 days in hydroponic system. The disinfection of the propagules contributed for relatively rapid rooting due to absence of contamination. Owing to the lack of cytogenetic studies for the genus *Dorstenia*, we tested different treatments with microtubule inhibitor as well as distinct procedures of




Figure 1. Representative adult plants of *D. elata* (a), *D. bonijesu* (c) and *D. arifolia* (e) of the Atlantic Rainforest remnant located in the Castelo city (ES, Brazil). FCM histograms showing G_0/G_1 peaks generated by nuclei suspensions of *S. lycopersicum* (internal standard, channel 200, $2C = 2.00$ pg), and of *D. elata* (b channel 349, $2C = 3.49$ pg), *D. bonijesu* (d channel 405, $2C = 4.05$ pg) and *D. arifolia* (f channel 547, $2C = 5.47$ pg).

enzymatic maceration. Root meristems treated with 4 μ M APM for 16 h at 4 °C and macerated in 1:60 pectinase solution for 1 h 45 min at 34 °C resulted in metaphases adequate for karyotype characterization of the *Dorstenia* species (Fig. 2).

The differences chromatin compaction levels verified among the metaphases (Fig. 2 a, b) of the species allowed morphometric analysis (Table 1) and karyogram assembly (Fig. 2). All *Dorstenia* species showed a conserved number of 32 chromosomes.

Morphometric analysis of the chromosomes

The mean values for the sum of total length as well as short- and long-arm length differed among the species (Table 1). *D. arifolia* showed the highest total, short- and long-arm length in relation to other species, corroborating the FCM data. *D. bonijesu*, which exhibited an intermediate mean 2C value, presented the lowest total chromosome and short-arm length. In comparison to *D. elata*, *D. bonijesu* displayed a greater mean value for the long arm (Table 1). The A_2 index also varied between the species: *D. bonijesu* exhibited the most asymmetrical karyotype, with $A_2 = 0.16$, followed by *D. arifolia* ($A_2 = 0.14$) and *D. elata* ($A_2 = 0.13$).

Based on the morphometric data, the chromosome class was determined and the differences between the karyotypes for the three species were endorsed. *D. elata* showed twelve metacentric chromosome pairs (1, 2, 4, 5, 7, 8, 9, 10, 13, 14, 15 and 16), two submetacentric (3 and 6) and two acrocentric pair (11 and 12) (Fig. 2 a, b). *D. bonijesu* exhibited four metacentric chromosome pairs (1, 2, 3 and 15), ten submetacentric (4, 5, 6, 8, 9, 10, 11, 12, 13 and 14) and two acrocentric ones (7 and 16) (Fig. 2c). *D. arifolia* presented four metacentric chromosome pairs (1, 2, 5 and 10) and twelve submetacentric ones (3, 4, 6, 7, 8, 9, 11, 12, 13, 14, 15 and 16) (Fig. 2d).

Discussion

Despite exhibiting the same number of chromosomes ($2n = 32$), the species of *Dorstenia* studied here show distinct mean nuclear 2C values. The interspecific DNA content variation indicates that the karyotypes differ between the species (Gitaí et al. 2014). Considering that the *Dorstenia* species have the same chromosome number, the karyotype divergences are probably associated to chromosome structure.

The morphometric analysis revealed karyomorphological differences in the sum of the mean values for total chromosome length, and the short arm and long arm (Table 1). Besides, some chromosomes presented distinct classes among the species, such as in chromosomes 7 and 16, which are submetacentric in *D. arifolia*, acrocentric in *D. bonijesu* and metacentric in *D. elata* (Fig. 2, Table 1). In view of this, structural chromosomal rearrangements have occurred during the evolutionary history of the group. This is supported by the interchromosomal asymmetry index (A_2 , Zarco 1986), which also varied between the three *Dorstenia* species, despite the predominance of meta-




Figure 2. Karyograms assembled from mitotic chromosomes of *D. elata* (a, b), *D. bonijesu* (c) and *D. arifolia* (d). Note the distinct chromatin compact level (a, b) showed by the mitotic chromosomes of *D. elata*. a, b *D. elata* showed twelve metacentric chromosome pairs (1, 2, 4, 5, 7, 8, 9, 10, 13, 14, 15 and 16), two submetacentric (3 and 6) and two acrocentric pair (11 and 12) c *D. bonijesu* exhibited four metacentric chromosome pairs (1, 2, 3 and 15), ten submetacentric (4, 5, 6, 8, 9, 10, 11, 12, 13 and 14) and two acrocentric ones (7 and 16) d *D. arifolia* presented four metacentric chromosome pairs (1, 2, 5 and 10) and twelve submetacentric ones (3, 4, 6, 7, 8, 9, 11, 12, 13, 14, 15 and 16). Bar: 5 μ m.

Table 1. Morphometry of the metaphasic chromosomes of *D. elata* (2C = 3.49 pg, 2n = 32), *D. bonijesu* (2C = 4.05 pg, 2n = 32) and *D. arifolia* (2C = 5.47 pg, 2n = 32).

Chrom.	<i>D. elata</i>						<i>D. bonijesu</i>						<i>D. arifolia</i>					
	Total (µm)	Arms		r	Class	Size (%)	Total (µm)	Arms		r	Class	Size (%)	Total (µm)	Arms		r	Class	Size (%)
		Short	Long					Short	Long					Short	Long			
1	4.72	2.09	2.63	1.26	M	8.01	4.50	1.95	2.55	1.31	M	7.97	5.39	2.57	2.82	1.10	M	7.93
2	4.27	2.00	2.27	1.14	M	7.25	4.27	1.90	2.37	1.25	M	7.56	5.21	2.45	2.76	1.13	M	7.67
3	4.21	1.66	2.55	1.54	SM	7.15	4.04	1.81	2.23	1.23	M	7.15	4.87	1.48	3.39	2.29	SM	7.17
4	3.87	1.78	2.09	1.17	M	6.57	4.00	1.45	2.55	1.76	SM	7.08	4.81	1.39	3.42	2.46	SM	7.08
5	3.84	1.75	2.09	1.19	M	6.52	3.86	1.27	2.59	2.04	SM	6.83	4.69	2.27	2.42	1.07	M	6.90
6	3.81	1.48	2.33	1.57	SM	6.47	3.81	1.45	2.36	1.63	SM	6.75	4.36	1.42	2.94	2.07	SM	6.42
7	3.81	1.69	2.12	1.25	M	6.47	3.72	0.90	2.82	3.13	A	6.59	4.33	1.39	2.94	2.12	SM	6.37
8	3.72	1.57	2.15	1.37	M	6.32	3.59	1.22	2.37	1.94	SM	6.36	4.18	1.45	2.73	1.88	SM	6.15
9	3.66	1.54	2.12	1.38	M	6.21	3.59	1.22	2.37	1.94	SM	6.36	4.18	1.27	2.91	2.29	SM	6.15
10	3.59	1.75	1.84	1.05	M	6.10	3.54	1.00	2.54	2.54	SM	6.27	3.93	1.81	2.12	1.17	M	5.78
11	3.54	0.71	2.83	3.99	A	6.01	3.40	0.95	2.45	2.58	SM	6.02	3.93	1.36	2.57	1.89	SM	5.78
12	3.48	0.62	2.86	4.61	A	5.91	3.04	0.86	2.18	2.53	SM	5.38	3.84	1.39	2.45	1.76	SM	5.65
13	3.30	1.33	1.97	1.48	M	5.60	2.95	0.95	2.00	2.11	SM	5.22	3.78	1.21	2.57	2.12	SM	5.56
14	3.24	1.42	1.82	1.28	M	5.50	2.95	0.81	2.14	2.64	SM	5.22	3.63	1.18	2.45	2.08	SM	5.34
15	3.03	1.30	1.73	1.33	M	5.14	2.63	1.09	1.54	1.41	M	4.66	3.51	1.24	2.27	1.83	SM	5.16
16	2.81	1.36	1.45	1.07	M	4.77	2.59	0.59	2.00	3.39	A	4.59	3.33	1.21	2.12	1.75	SM	4.90
Sum	56.09	24.36	34.54	-	-	100.00	53.89	19.42	37.06	-	-	100.00	64.64	25.09	42.88	-	-	100.00

Chrom – chromosome; r – arm ratio (long/short); Size – % size in relation to sum of the mean values of total length; M – metacentric; SM – submetacentric; A – acrocentric; Sum – sum of the mean values.

centric and submetacentric chromosomes. Therefore, the FCM and cytogenetic data indicate that structural chromosome changes have occurred throughout the evolution of these *Dorstenia* species.

According to the more recent phylogeny for the genus, *D. arifolia* occupies a basal position in comparison to *D. elata* (Misiewicz and Zerega 2012). Given that *D. arifolia* has mean nuclear 2C value and total genome length (Fig. 1 and 2, Table 1) greater than *D. elata*, the structural chromosome alterations seem to promote loss of DNA sequences. Thus, deletions can be involved in the karyotype changes in *Dorstenia*.

Based on the morphometric analysis, groups of morphologically identical chromosome pairs were found for each *Dorstenia* species: 3–4, 6–7, 11–12 and 13–14 in *D. arifolia*; 5–6, 11–12 and 13–14 in *D. bonijesu*; and 7–8 and 14–15 in *D. elata* (Fig. 2, Table 1). Regarding this karyotype aspect, polyploidization events have occurred during the evolution of these species. As the three species produces reduced reproductive cells ($x = 16$), the dispoidy can also explain the evolutionary scenario of the *Dorstenia* karyotype. The polyploid origin of plant species has been shown by classical cytogenetics, such as for the genera *Psidium* Linnaeus, 1753 (Souza et al. 2015), *Claytonia* Linnaeus, 1753 (McIntyre 2012) and *Cardamine* Linnaeus, 1753 (Marhold et al. 2010). From the assembly of accurate karyograms using chromosomes with different levels of chromatin compaction, the type of polyploidy was also evidenced, with autopolyploidy being found in *Glycine max* (Linnaeus, 1753) Merrill, 1917 (Clarindo et al. 2007) and *Zephyranthes* Herbert, 1821 (Felix et al. 2011), and allopolyploidy in *Paullinia cupana* Kunth, 1821 (Freitas et al. 2007) and *Triticum aestivum* Linnaeus, 1753 (Kamel 2006). Polyploidy has played a key role in plant evolution, with estimates maintaining euploidy in the ascendancy of all angiosperms (Abbott et al. 2013, Soltis et al. 2014). In addition, 15% of speciation events in this taxon are directly involved with polyploidization (Abbott et al. 2013).

Conclusion

The nuclear 2C value and karyogram indicate changes covering chromosome number and structure that occurred during the karyotype evolution of *D. arifolia*, *D. bonijesu* and *D. elata*. The combination of FCM and classical cytogenetics revealed differences among the *Dorstenia* species that can be exploited in phylogenetic approaches, as the results support the current knowledge on the phylogeny of *Dorstenia*.

Acknowledgements

The authors are grateful to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brasília – DF, Brazil), Fundação de Amparo à Pesquisa do Espírito Santo (FAPES, Vitória – ES, Brazil), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG, Belo Horizonte – MG, Brazil), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brasília – DF, Brazil) for financial support.

References

- Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJE, Bierne N, Boughman J, Brelsford A, Buerkle CA, Buggs R, Butlin RK, Dieckmann U, Eroukhmanoff F, Grill A, Cahan SH, Hermansen JS, Hewitt G, Hudson AG, Jiggins C, Jones J, Keller B, Marczewski T, Mallet J, Martinez-Rodriguez P, Möst M, Mullen S, Nichols R, Nolte AW, Parisod C, Pfennig K, Rice AM, Ritchie MG, Seifert B, Smadja CM, Stelkens R, Szymura JM, Väinölä R, Wolf JBW, Zinner D (2013) Hybridization and speciation. *Journal of Evolutionary Biology* 26: 229–246. doi: 10.1111/j.1420-9101.2012.02599.x
- Berg CC (2001) *Moreae, Artocarpeae and Dorstenia* (Moraceae). With introductions to the family and *Ficus* and with additions and corrections to *Flora Neotropica Monograph 7*. *Flora Neotropica Monograph* 83: 1–346.
- Bonifácio EM, Fonsêca A, Almeida C, Santos KGB, Pedrosa-Harand A (2012) Comparative cytogenetic mapping between the lima bean (*Phaseolus lunatus* L.) and the common bean (*P. vulgaris* L.). *Theoretical and Applied Genetics* 124: 1513–1520. doi: 10.1007/s00122-012-1806-x
- Carauta JPP (1978) *Dorstenia* L. (Moraceae) do Brasil e países limítrofes. *Rodriguésia* 29: 53–223.
- Carvalho CR, Clarindo WR, Almeida PM (2007) Plant cytogenetics: still looking for the perfect mitotic chromosomes. *Nucleus* 50: 453–462.
- Clarindo WR, Carvalho CR, Alves BMG (2007) Mitotic evidence for the tetraploid nature of *Glycine max* provided by high quality karyograms. *Plant Systematics and Evolution* 265: 101–107. doi: 10.1007/s00606-007-0522-5
- Clarindo WR, Carvalho CR, Mendonça MAC (2012) Cytogenetic and flow cytometry data expand knowledge of genome evolution in three *Coffea* species. *Plant Systematics and Evolution* 298: 835–844. doi: 10.1007/s00606-012-0595-7
- Fagerlind F (1944) Die samenbildung und die zytology bei agamosper-mischen und sexuellen arten von *Elatostema* und einigen nahestehenden gattungen nebst gelechtung einiger damit zusammenhängender probleme. *Kung Svensk Vet Akad Handl* 21: 1–130.
- Felix WJP, Felix LP, Melo NF, Oliveira MBM, Dutilh JHA, Carvalho R (2011) Karyotype variability in species of the genus *Zephyranthes* Herb. (Amaryllidaceae–Hippeastreae). *Plant Systematics and Evolution* 294: 263–271. doi: 10.1007/s00606-011-0467-6
- Freitas DV, Carvalho CR, Filho FJN, Astolfi-Filho S (2007) Karyotype with 210 chromosomes in guaraná (*Paullinia cupana* ‘Sorbilis’). *Journal of Plant Research* 120: 399–404. doi: 10.1007/s10265-007-0073-4
- Galbraith DW, Harkins KR, Maddox JM, Ayres NM, Sharma DP, Firoozabady E (1983) Rapid flow cytometric analysis of the cell cycle in intact plant tissues. *Science* 220: 1049–1051. doi: 10.1126/science.220.4601.1049
- Gao YD, Zhou SD, He XJ, Wan J (2012) Chromosome diversity and evolution in tribe Liliaceae (Liliaceae) with emphasis on Chinese species. *Journal of Plant Research* 125: 55–69. doi: 10.1007/s10265-011-0422-1
- Gitai J, Paule J, Zizka G, Schulte K, Benko-Iseppon AM (2014) Chromosome numbers and DNA content in Bromeliaceae: additional data and critical review. *The Journal of the Linnean Society* 176: 349–368. doi: 10.1111/boj.12211

- Guerra MS (1986) Reviewing the chromosome nomenclature of Levan et al. *Revista Brasileira de Genética* 9: 741–743.
- Kamel EAR (2006) Karyotype characterization and polyploid variations in some genotypes of *Triticum aestivum* L. *Cytologia* 71: 169–174. doi: 10.1508/cytologia.71.169
- Krause O (1931) Zytologische Studien bei den Urticales unter besonderer Berücksichtigung der Gattung *Dorstenia*. *Planta* 13: 29–84. doi: 10.1007/BF01908980
- Levan A, Fredga K, Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201–220. doi: 10.1111/j.1601-5223.1964.tb01953.x
- Loureiro J, Rodriguez E, Doležel J, Santos C (2006) Comparison of four nuclear isolation buffers for plant DNA flow cytometry. *Annals of Botany* 98: 679–689. doi: 10.1093/aob/mcl141
- Mabuchi T, Kokubun H, Mii M, Ando T (2005) Nuclear DNA content in the genus *Hepatica* (Ranunculaceae). *Journal of Plant Research* 118: 37–41. doi: 10.1007/s10265-005-0191-9
- Marhold K, Kudoh H, Pak JH, Watanabe K, Španiel S, Lihová J (2010) Cytotype diversity and genome size variation in eastern Asian polyploid *Cardamine* (Brassicaceae) species. *Annals of Botany* 105: 249–264. doi: 10.1093/aob/mcp282
- McIntyre JP (2012) Cyto geography and genome size variation in the *Claytonia perfoliata* (Portulacaceae) polyploid complex. *Annals of Botany* 110: 1195–1203. doi: 10.1093/aob/mcs187
- Melo NF, Guerra M (2003) Variability of the 5S and 45S rDNA sites in *Passiflora* L. species with distinct base chromosome numbers. *Annals of Botany* 92: 309–316. doi: 10.1093/aob/mcg138
- Misiewicz TM, Zerega NC (2012) Phylogeny, biogeography and character evolution of *Dorstenia* (Moraceae). *Edinburgh Journal of Botany* 69: 413–440. doi: 10.1017/S096042861200025X
- Otto FJ (1990) DAPI staining of fixed cells for high-resolution flow cytometry of nuclear DNA. In: Darzynkiewicz Z, Crissman HA, Robinson JP (Eds) *Methods in cell biology*. San Diego Academic Press, San Diego, Ca, 105–110. doi: 10.1016/s0091-679x(08)60516-6
- Peer YV, Maere S, Meyer A (2009) The evolutionary significance of ancient genome duplications. *Nature Reviews Genetics* 10: 725–732. doi: 10.1038/nrg2600
- Praça-Fontes MM, Carvalho CR, Clarindo WR, Cruz CD (2011) Revisiting the DNA C-values of the genome size-standards used in plant flow cytometry to choose the “best primary standards”. *Plant Cell Reports* 30: 1183–1191. doi: 10.1007/s00299-011-1026-x
- Raskina O, Barber JC, Nevo E, Belyayev A (2008) Repetitive DNA and chromosomal rearrangements: speciation-related events in plant genomes. *Cytogenetic and Genome Research* 120: 351–377. doi: 10.1159/000121084
- Romaniuc-Neto S, Carauta JPP, Vianna-Filho MDM, Pereira RAS, Ribeiro JELS, Machado AFP, Santos A, Pelissari G, Pederneiras LC (2015) Moraceae in Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB10115> [accessed 22 June 2015]
- Shan F, Yan G, Plummer JA (2013) Karyotype evolution in the genus *Boronia* (Rutaceae). *The Journal of the Linnean Society* 142: 309–320. doi: 10.1046/j.1095-8339.2003.00163.x

- Soltis DE, Visger CJ, Soltis PS (2014) The polyploidy revolution then...and now: Stebbins revisited. *American Journal of Botany* 101: 1057–1078. doi: 10.3732/ajb.1400178
- Souza AG, Resende LV, Lima IP, Martins LSS, Techio VH (2015) Chromosome number and nuclear DNA amount in *Psidium* spp. resistant and susceptible to *Meloidogyne enterolobii* and its relation with compatibility between rootstocks and commercial varieties of guava tree. *Plant Systematics and Evolution* 301: 231–237. doi: 10.1007/s00606-014-1068-y
- Stace CA (2000) Cytology and cytogenetics as a fundamental taxonomic resource for the 20th and 21st centuries. *Taxon* 49: 451–477. doi: 10.2307/1224344
- Weiss-Schneeweiss H, Emadzade K, Jang TS, Schneeweiss GM (2013) Evolutionary consequences, constraints and potential of polyploidy in plants. *Cytogenetic and Genome Research* 140: 137–150. doi: 10.1159/000351727
- Zarco RC (1986) A new method for estimating karyotype asymmetry. *Taxon* 35: 526–530. doi: 10.2307/1221906