

Cytogenetic characterization of three Balistoidea fish species from the Atlantic with inferences on chromosomal evolution in the families Monacanthidae and Balistidae

Lorena Corina Bezerra de Lima, Pablo Ariel Martinez, Wagner Franco Molina

Department of Cell Biology and Genetics, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Campus Universitário, 59078 – 970 Natal, RN, Brazil

Corresponding author: *Wagner Franco Molina* (molinawf@pq.cnpq.br)

Academic editor: *Snejana Grozeva* | Received 16 February 2010 | Accepted 23 March 2010 | Published 5 May 2011

Citation: de Lima LCB, Martinez PA, Molina WF (2011) Cytogenetic characterization of three Balistoidea fish species from the Atlantic with inferences on chromosomal evolution in the families Monacanthidae and Balistidae. *Comparative Cytogenetics* 5(1): 61–69. doi: 10.3897/compcytogen.v5i1.1141

Abstract

The Tetraodontiformes are the most derived group of teleostean fish. Among other apomorphies, they are characterized by a high degree of fusions or significant bone loss in the head and body. In the early phylogenetic proposals presented for this order, the families Balistidae and Monacanthidae have been unanimously considered to be closely related. Although they have moderate species diversity, they are scarcely known in cytogenetic aspect and chromosomal pattern comparisons between these groups have yet to be established. The species *Cantherhines macrocerus* (Hollard, 1853), *C. pullus* (Ranzani, 1842) (Monacanthidae) and *Melichthys niger* (Bloch, 1786) (Balistidae) were cytogenetically analyzed using conventional (Ag-impregnation, C-banding, CMA₃- and DAPI-fluorescence) and molecular (FISH with an *18S rDNA* probe) cytogenetic protocols. The karyotypes of all three species were very similar possessing diploid chromosome numbers $2n = 40$ and composed exclusively of acrocentric chromosomes. Single NOR-bearing pair as well as positive heterochromatic blocks at pericentromeric regions were identified in the karyotypes of the three species studied. NOR-bearing sites were positively labeled after Ag-impregnation, C-banding, CMA₃-fluorescence and FISH with an *18S rDNA* probe but were negative after DAPI-fluorescence. Such remarkable shared conspicuous chromosomal characters corroborate either close phylogenetic relationship of these families, previously established by morphological and molecular data, or rather conservative nature of karyotype differentiation processes. The later hypothesis, however, appears less probable due to centric or *in tandem* fusions documented for another Balistoidea species.

Keywords

Balistoidea, fish cytogenetics, karyotype evolution, Tetraodontiformes

Introduction

The order Tetraodontiformes, which stands out among marine fish for its marked diversity, is composed of approximately 430 species distributed in nine families (Nelson 2006). This group is the most recent branch of Neoteleostean radiation, representing a post-Perciformes lineage (Elmerot et al. 2002) and although it is generally recognized as a monophyletic group, the relationships between its families and genera have yet to be defined (Holcroft 2004).

Among the Tetraodontiformes, the superfamily Balistoidea (leatherjackets) includes the families Balistidae (triggerfish) and Monacanthidae (filefish), with a fossil record that dates back to the Early Eocene and probably to the Late Cretaceous (Frickhinger 1995, Santini and Tyler 2003). Based on morphological similarities (e.g., osteological and myological characters), molecular studies using RAG1 gene sequences and DNA content data (Winterbottom 1974, Brainerd et al. 2001, Holcroft 2004, 2005), and more recently analysis of complete mitochondrial genomes (mitogenome) (Yamanoue et al. 2008), members of families Balistidae and Monacanthidae are considered monophyletic sister groups.

To date nearly 60 species of Tetraodontiformes have been cytogenetically studied (Sá-Gabriel and Molina 2005). Cytogenetic analyses have been carried out in 15 Balistidae species, most from the Pacific region, and only few from the Western Atlantic. A total of ten Monacanthidae species are karyotyped, being one from the Brazilian coast.

In this work, we revise cytogenetic data for *Melichthys niger* (Bloch, 1786) (Balistidae) and describe the karyotype and other chromosomal characteristics for *Cantherhines macrocerus* (Hollard, 1853) and *C. pullus* (Ranzani, 1842) (Monacanthidae), to compare the chromosomal patterns of the families Balistidae and Monacanthidae, estimating their divergence level.

Material and methods

We analyzed 42 specimens of *Melichthys niger* and 14 of *Cantherhines macrocerus*, collected in the Saint Peter and Saint Paul's Archipelago (00°55'15" N, 029°20'60" W), 1010 km from the Brazilian northeastern coast and about 1.824 km from the African coast, and two specimens of *Cantherhines pullus* collected in the coastal region of Salvador (12° 58'S, 38° 31'W), Bahia state, northeastern Brazil (Fig. 1).

The specimens were subjected to mitotic stimulation as proposed by Molina (2001) for a period of 24–48 hours. Mitotic chromosomes were obtained following the protocol developed by Gold et al. (1990). Cell suspensions obtained from fragments of the anterior portion of specimens' kidney were spread onto slides with a film of water heated at 60°C. The active nucleolus organizer regions (NOR) were identified by silver nitrate staining as described by Howell and Black (1980), while the heterochromatic regions were visualized by C-banding (Sumner 1972). The FISH technique (according to Pinkel et al. 1986) was performed using an 18S rDNA probe from *Prochilodus*




Figure 1. Map showing the geographic collection points of the species *Melichthys niger*, *Cantherhines macrocerus* and *C. pullus*. SPSPA – Saint Peter and Saint Paul's Archipelago; BA – Bahia state.

argenteus, Agassiz, 1829 (Hatanaka and Galetti 2004), labeled with biotin-14-dATP by nick translation according to the manufacturer's instructions (BioNick Labeling System; Invitrogen, Carlsbad, CA, U.S.A.). The hybridization signal was detected by the streptavidin-fluorescein isothiocyanate conjugate. Sequential staining with AT-specific 4'-6-diamidino-2-phenylindole (DAPI) and GC-specific Chromomycin A₃ (CMA₃) fluorochromes was performed as described by Schweizer (1980). The metaphases were photographed by a DP70 digital image capture system coupled to an Olympus BX50 epifluorescence microscope. About thirty metaphases of each specimen were analyzed to determine the modal number of mitotic chromosomes and the karyotype. The chromosomes were classified according to centromere position and organized into decreasing size order, as proposed by Levan et al. (1964).

Results

The specimens of *Melichthys niger* showed $2n=40$ chromosomes and a karyotype consisting of 20 pairs of acrocentric (a) chromosomes (NF=40) (Fig. 2a). The presence of a conspicuous secondary constriction was observed in the interstitial position on the long arm of the chromosome pair No. 2, corresponding to the nucleolus organizer regions (NORs), identified by Ag-NOR sites and by *in situ* hybridization with an *18S rDNA* ribosomal probes (Fig. 2, upper boxes). The heterochromatic blocks were reduced in size and dispersed in the pericentromeric regions in most of the chromo-




Figure 2. Karyotypes of *Melichthys niger* (a, b), *Cantherhines macrocerus* (c, d) and *Cantherhines pullus* (e, f), arranged from Giemsa stained (a, c, e) and C-banded chromosomes (b, d, f). In the center highlighted are the NOR-bearing pairs of analyzed species (2nd, 7th and 5th, respectively) after Ag-NOR staining, *in situ* hybridization with an 18S rDNA probe, CMA₃ and DAPI fluorescence. Bar = 5µm.

some pairs (Fig. 2b). The NORs were heterochromatic and CMA₃ positive and DAPI negative (Fig. 2, upper boxes).

The specimens of *Cantherhines macrocerus* (Monacanthidae) had 2n=40 chromosomes and karyotype composed of a acrocentric chromosomes (Fig. 2c). Ag-NOR

sites were located in the pair No. 7 in interstitial region, near the centromere (Figure 2, middle box). C-banding revealed heterochromatic blocks distributed in the pericentromeric region in most of the chromosome pairs (Figure 2d) and more intensively stained on the secondary constriction of the NOR-bearing pair. In this species, experiments using FISH probes and fluorochrome staining were unsuccessful.

The specimens of *C. pullus* showed $2n=40$ chromosomes and karyotype composed entirely of acrocentric chromosomes (Fig. 2e). The NORs were identified at the pericentromeric position of pair No. 5, as revealed by Ag-NOR-staining and FISH with an *18S rDNA* probes (Fig. 2, lower box). The heterochromatic regions were distributed in centromeric and pericentromeric positions in most of the chromosomes. The NORs sites were heterochromatic (Fig. 2f), and CMA positive and DAPI negative (Fig. 2, lower box). None of the karyotypes displayed sex-related chromosome heteromorphism.

Discussion

Six out of the ten karyotyped species in the family Monacanthidae have diploid numbers ranging from $2n = 33/34$ to 36 chromosomes. Such low $2n$ numbers have been a noticeable characteristic for Monacanthidae species. The present data for *C. macrocerus* and *C. pullus* increase the range of the highest $2n$ for representatives of this family. Surveys involving a larger number of genera may confirm a possible basal karyotype with 40 chromosomes for this family, showing on average lower diploid values than those of the Balistidae. Based on chromosomal number, *C. macrocerus* and *C. pullus* would be placed in the family Balistidae.

The karyotype of the individuals of *C. macrocerus* from the Saint Peter and Saint Paul's Archipelago was similar to those described for specimens from the coast of Rio de Janeiro (Pauls 1993), even though these populations were 2000 km apart. It is not clear whether the common karyotype found in *C. macrocerus* populations is maintained due to gene flow by the transport of their pelagic larvae by ocean currents, as has been commonly identified for a number of reef species (Rocha 2003, Feitoza et al. 2005), or if they were isolated too recently to accumulate observable chromosomal differences.

In this study, the data obtained for *Melichthys niger* corroborate results described for the species earlier (Sá-Gabriel and Molina 2005). However, the use of a larger number of individuals of this species, observation of lower chromatid condensation and the use of complementary cytogenetic protocols, such as CMA₃/DAPI fluorochromes and FISH with an *18S rDNA* probes, allow more precise localization of the NOR-bearing pair to the 2nd largest pair of the karyotype.

The chromosomal characteristics observed in *C. macrocerus*, *C. pullus* and *M. niger*, like the presence of single NOR and the pericentromeric heterochromatin blocks as reported in other Tetraodontiformes (Grützner et al. 1999, Mandrioli 2000, Fischer et al. 2000), Perciformes (Caputo et al., 2001; Molina, 2007), Mugiliformes (Nirchio et al. 2009), Beryciformes (Bacurau and Molina 2004), corroborate the hypothesis that

these are ancestral characteristics for each of these clades and therefore not exclusive to the families Monacanthidae and Balistidae.

In the karyotypes of species under our study, as well as in other representatives of Balistidae and Monacanthidae (Sá-Gabriel and Molina 2004), extra-pericentromeric heterochromatic regions were present only when adjacent to or associated with the major ribosomal sites. Given that the constitutive heterochromatin blocks in fish chromosomes are often associated with karyotype diversification (Mantovani et al. 2000; Molina and Bacurau 2006), it is possible that these regions present preferentially in peri-centromeric position in chromosomes of Balistoidea species and they are involved in the karyotypic differentiation of this group. The presence of heterochromatin associated with NORs in adjacent or interspersed regions (Pendás et al. 1993), as shown in the NOR-bearing pair in *M. niger*, may contribute to the occurrence of structural rearrangements involving NOR-bearing pairs (Vicari et al. 2003). Indeed, chromosome fusions have been commonly identified as the main mechanism of the karyotype diversification within this clade (Kitayama and Ojima 1984). Although a physical mapping of telomeric sequences is not yet available, the lower $2n$ in both Balistidae and Monacanthidae corroborate that centric and/or *in tandem* fusion events, followed by pericentric inversions, seem to have been important mechanisms in the karyotype differentiation of these post-Perciformes group.

Repetitive sequences and transposition elements are closely related to the heterochromatic regions, and although account for less than 10% of the genome of the Tetraodontiformes studied (Brenner et al. 1993), they may be effective promoters of chromosomal breaks, deletions, inversions and amplifications (Lim and Simmons 1994, Fischer et al. 2000). Knowledge regarding aspects of the Balistoidea genome is still limited, although it is known that, like Tetraodontidae, this clade is composed of species with compact genomes that evolved independently (Brainerd et al. 2001). The Tetraodontidae genome has been extensively studied (Crolius et al. 2000), revealing that the small amount of repetitive sequences is located in the centromeres and arms of a chromosomes. This physical disposition seems to be present in the chromosomes of Balistoidea and could help to explain why centromere and telomere regions would be prone to occurrence of *in tandem* or centric fusions rearrangements. In addition to the aforementioned data, a large number of acrocentric chromosomes and FN near or slightly higher than the diploid chromosome values were observed. These characteristics are not common in other families of the same order, such as the Tetraodontidae (Sá-Gabriel and Molina 2005), where pericentric inversions seem to have occurred more frequently.

Among the Perciformes, the presence of single interstitial NORs is common, representing an ancestral condition (Affonso et al. 2001). Similarly, the presence of heterochromatin restricted to the centromeric regions is also a typical cytogenetic trait of this group (Molina and Galetti 2002). The location and frequency of NORs sites and heterochromatic regions in *C. macrocerus*, *C. pullus* and *M. niger* are consistent with the general pattern found in Perciformes species, demonstrating the apparent conservativeness of these characters.

The set of cytogenetic characters already available in Monacanthidae and Balistidae species indicate a greater karyotypic similarities and common tendencies of karyotype evolution with other groups of the order Tetraodontiformes, corroborating previous analyses based on morphological and molecular data (Winterbottom 1974; Brainerd et al. 2001, Holcroft 2005; Yamanoue et al. 2008).

Acknowledgments

We would like to thank Universidade Federal do Rio Grande do Norte (UFRN) and Comissão Interministerial para os Recursos do Mar (CIRM) for providing the facilities and proper conditions for the accomplishment of the present survey and José Garcia Jr for species identification. This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq – Proc. 556793/2009-9).

References

- Affonso PRAM, Guedes W, Pauls E, Galetti Jr PM (2001) Cytogenetic analysis of coral reefs fishes from Brazil (Families Pomacanthidae and Chaetodontidae). *Cytologia* 66: 379–384.
- Bacurau TOF, Molina WF (2004) Karyotypic diversification in two Atlantic species of Holocentridae (Pisces, Beryciformes). *Caryologia* 57: 300–304.
- Brainerd EL, Slutz SS, Hall EK, Phillis RW (2001) Patterns of genome size evolution in tetraodontiform fishes. *Evolution* 55: 2363–2368.
- Brenner S, Elgar G, Sandford R, Macrae A, Venkatesh B, Aparicio S (1993) Characterization of the pufferfish (*Fugu*) genome as a compact model vertebrate genome. *Nature* 336: 265–268. doi:10.1038/366265a0
- Caputo V, Machella N, Nisi-Cerioni P, Olmo E (2001) Cytogenetics of nine species of Mediterranean blennies and additional evidence for an unusual multiple sex-chromosome system in *Parablennius tentacularis* (Perciformes, Blenniidae). *Chromosome Res.* 9: 3–12. doi:10.1023/A:1026779314932
- Crollius HR, Jaillon O, Dasilva C, Ozouf-Costaz C, Fizames C, Fischer C, Bouneau L, Billaud A, Quetier F, Saurin W, Bernot A, Weissenbach J (2000) Characterization and repeat analysis of the compact genome of the freshwater pufferfish *Tetraodon nigroviridis*. *Genome Res.* 10: 939–949. doi:10.1101/gr.10.7.939
- Elmerot C, Arnason U, Gojobori T, Janke A (2002) The mitochondrial genome of the pufferfish, *Fugu rubripes*, and ordinal teleostean relationships. *Gene* 295: 163–172. doi:10.1016/S0378-1119(02)00688-1
- Feitoza BM, Rosa RS, Rocha LA (2005) Ecology and zoogeography of deep reef fishes in northeastern Brazil. *B. Mar. Sci.* 76: 725–742.
- Fischer C, Ozouf-Costaz C, Crollius HR, Dasilva C, Jaillon O, Bouneau L, Bonillo C, Weissenbach J, Bernot A (2000) Karyotype and chromosomal location of characteristic tan-

- dem repeats in the pufferfish *Tetraodon nigroviridis*. *Cytogenet. Cell Genet.* 88: 50–55. doi:10.1159/000015484
- Frickhinger KA (1995) Fossil Atlas Fish. Blacksburg Tetra Press. 1088 p.
- Gold JR, Li C, Shipley NS, Powers PK (1990) Improved methods for working with fish chromosomes with a review of metaphase chromosome banding. *J. Fish Biol.* 37: 563–575. doi:10.1111/j.1095-8649.1990.tb05889.x
- Grützner F, Lütjens G, Rovira C, Barnes DW, Ropers H-H, Haaf T (1999) Classical and molecular cytogenetics of the pufferfish *Tetraodon nigroviridis*. *Chromosome Res.* 7: 655–662. doi:10.1023/A:1009292220760
- Hatanaka T, Galetti Jr PM (2004) Mapping of the 18S and 5S ribosomal genes in the fish *Prochilodus argenteus* Agassiz, 1829 (Characiformes, Prochilodontidae). *Genetica* 122: 239–244. doi:10.1007/s10709-004-2039-y
- Holcroft NI (2004) A molecular test of alternative hypotheses of tetraodontiform (Acanthomorpha: Tetraodontiformes) sister group relationships using data from the RAG1 gene. *Mol. Phylogenet. Evol.* 32: 749–760. doi:10.1016/j.ympev.2004.04.002
- Holcroft NI (2005) A molecular analysis of the interrelationships of tetraodontiform fishes (Acanthomorpha: Tetraodontiformes). *Mol. Phylogenet. Evol.* 34: 525–544. doi:10.1016/j.ympev.2004.11.003
- Howell WM, Black DA (1980) Controller silver staining of nucleolus organizer region with protective colloidal developer: a 1 – step method. *Experientia* 36: 1014–1015. doi:10.1007/BF01953855
- Kitayama E, Ojima Y (1984) Preliminary report on the phylogeny of five balistid fish in terms of several chromosome banding techniques in combination with a statistical analysis. *P. Jpn. Acad. B: Phys. Biol. Sci.* 60(3): 58–61. doi:10.2183/pjab.60.58
- Levan A, Fredga K, Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas.* 52: 201–220. doi:10.1111/j.1601-5223.1964.tb01953.x
- Lim JK, Simmons MJ (1994) Gross chromosome rearrangements mediated by transposable elements in *Drosophila melanogaster*. *Bioessays* 16: 269–75. doi:10.1002/bies.950160410
- Mandrioli M (2000) Cytogenetic analysis of the pufferfish *Tetraodon fluviatilis* (Osteichthyes). *Chromosome Res.* 8: 237–242. doi:10.1023/A:1009257131091
- Mantovani M, Abel LDS, Mestriner CA, Moreira-Filho O (2000) Accentuated polymorphism of heterochromatin and nucleolar organizer regions in *Astyanax scabripinnis* (Pisces, Characidae): tools for understanding karyotypic evolution. *Genetica* 109: 161–168. doi:10.1023/A:1017546601065
- Molina WF (2001) An alternative method for mitotic stimulation in fish cytogenetics. *Chromosome Sci.* 5: 149–152.
- Molina WF (2007) Chromosomal changes and stasis in marine fish groups. In: Pisano E, Ozouf-Costaz C, Foresti F, Kapoor BG (Eds) *Fish cytogenetics*. Enfield: Science publishers, 69–110.
- Molina WF, Galetti Jr PM (2002) Robertsonian rearrangements in the reef fish *Chromis* (Perciformes, Pomacentridae) involving chromosomes bearing 5S rRNA genes. *Genet. Mol. Biol.* 25: 373–377. doi:10.1590/S1415-47572002000400004

- Molina WF, Bacurau TOF (2006) Structural and numerical chromosomal variation in marine Perciformes (Priacanthidae and Gerreidae). *Cytologia*. 71: 237–242. doi:10.1508/cytologia.71.237
- Nelson JS (2006) *Fishes of the World*. New York. 601 p.
- Nirchio M, Oliveira C, Ferreira IA, Martins C, Rossi AR, Sola L (2009) Classical and molecular cytogenetic characterization of *Agonostomus monticola*, a primitive species of Mugilidae (Mugiliformes). *Genetica* 135(1): 1–5. doi:10.1007/s10709-008-9256-8
- Pauls E (1993) Estudos citogenéticos em peixes marinhos visando o melhoramento genético. Ph.D. Dissertation. Faculdade de Veterinária da Faculdade de Veterinária de UFF, Niterói, RJ, 234 p. [in Portuguese]
- Pendás AM, Moran P, Garcia-Vazquez E (1993) Multi chromosomal location of ribosomal RNA genes and heterochromatin association in brown trout // *Chromosome Res.* 1: 63–67. doi:10.1007/BF00710608
- Pinkel D., Straume T., Gray J. 1986. Cytogenetic analysis using quantitative, high sensitivity, fluorescence hybridization. *Proc. Natl. Acad. Sci. USA*. 83: 2934–2938. doi:10.1073/pnas.83.9.2934
- Rocha LA (2003) Patterns of distribution and processes of speciation in Brazilian reef fishes. *J. Biogeogr.* 30 (8): 1161–1171. doi:10.1046/j.1365-2699.2003.00900.x
- Sá-Gabriel LG, Molina WF (2004) Sex chromosomes in *Stephanolepis hispidus* (Monacanthidae, Tetraodontiformes). *Cytologia* 69: 447–452. doi:10.1508/cytologia.69.447
- Sá-Gabriel LG, Molina WF (2005) Karyotype diversification in fishes of the Balistidae, Diodontidae and Tetraodontidae (Tetraodontiformes). *Caryologia* 58: 229–237.
- Santini F, Tyler JC (2003) A phylogeny of the families of fossil and extant tetraodontiform fishes (Acanthomorpha, Tetraodontiformes), Upper Cretaceous to Recent. *Zool. J. Linn. Soc-Lond.* 139: 565–617.
- Schweizer D (1980) Simultaneous fluorescent staining of R-bands and specific heterochromatic regions (DA/DAPI bands) in human chromosomes. *Cytogenet Cell Genet.* 27: 190–193. doi:10.1159/000131482
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Exp. Cell Res.* 75: 304–306. doi:10.1016/0014-4827(72)90558-7
- Vicari MR, Artoni RF, Bertollo LAC (2003) Heterochromatin polymorphism associated with 18S rDNA. A differential pathway among *Hoplias malabaricus* fish populations. *Cytogenet. Genome Res.* 101: 24–28. doi:10.1159/000073413
- Winterbottom R (1974) The familial phylogeny of the Tetraodontiformes (Acanthopterygii: Pisces) as evidenced by their comparative myology. *Smithsonian Cont. Zool.* 155: 1–201. doi:10.5479/si.00810282.155
- Yamanoue Y, Miya M, Matsuura K, Katoh M, Sakai H, Nishida M (2008) A new perspective on phylogeny and evolution of tetraodontiform fishes (Pisces: Acanthopterygii) based on whole mitochondrial genome sequences: Basal ecological diversification? *BMC Evol. Biol.* 8: 212. doi:10.1186/1471-2148-8-212