

Comparative cytogenetic study of three *Macrolophus* species (Heteroptera, Miridae)

Ana Maria Jauset¹, Eva Edo-Tena¹, Cristina Castañé², Nuria Agustí²,
Oscar Alomar², Snejana Grozeva³

1 University of Lleida, Department of Crop and Forest Sciences. Alcalde Rovira Roure, 177, 25198 Lleida (Spain)
2 IRTA, Ctra. Cabriels Km 2, 08348 Cabriels (Barcelona), Spain **3** Institute of Biodiversity and Ecosystem Research,
Bulgarian Academy of Sciences, 1 Tsar Osvoboditel Blvd., 1000 Sofia, Bulgaria

Corresponding author: Snejana Grozeva (sgrozeva@gmail.com)

Academic editor: S. Nokkala | Received 30 June 2015 | Accepted 19 August 2014 | Published 29 September 2015

<http://zoobank.org/DD12EC27-7BC0-4C6B-A3BB-4E553FBBB0F5>

Citation: Jauset AM, Edo-Tena E, Castañé C, Agustí N, Alomar O, Grozeva S (2015) Comparative cytogenetic study of three *Macrolophus* species (Heteroptera, Miridae). *Comparative Cytogenetics* 9(4): 613–623. doi: 10.3897/CompCytogen.v9i4.5530

Abstract

Macrolophus pygmaeus (Rambur, 1839) (Insecta, Heteroptera, Miridae) is a predator of key vegetable crop pests applied as a biocontrol agent in the Mediterranean region. *M. pygmaeus* and *M. melanotoma* (A. Costa, 1853) are cryptic species with great morphological similarity which results in their misidentification and negative consequences for the conservation of their populations on greenhouse and outdoor crops. In order to find out specific markers for their separation we studied the karyotype, male meiosis and heterochromatin composition of these species and additionally of a third species (as a reference one), *M. costalis* Fieber, 1858. We demonstrate here that all the three species share achiasmate male meiosis and sex chromosome pre-reduction. On the other hand, the species differ in karyotype, with $2n=28$ ($26+XY$) in *M. pygmaeus*, $2n=27$ ($24+X_1X_2Y$) in *M. costalis*, and $2n=34$ ($32+XY$) in *M. melanotoma*, and heterochromatin distribution and composition. In addition, the species differ in sperm morphology: sperm cells of *M. costalis* are significantly longer with longer head and tail than those of *M. melanotoma* and *M. pygmaeus*, whereas sperm cells of *M. melanotoma* have a longer tail than those of *M. pygmaeus*. All these characters can be used as markers to identify the species, in particular the cryptic species *M. melanotoma* and *M. pygmaeus*.

Keywords

Macrolophus, Miridae, Heteroptera, karyotype, sex chromosomes, achiasmate meiosis, sex chromosome pre-reduction, sperm morphology

Introduction

The Miridae are the largest family of true bugs (Heteroptera, Cimicomorpha) with approximately 10000 species described (Schuh 1995). Cytogenetical data are presently available for about 200 species (Ueshima 1979, Nokkala and Nokkala 1986, Grozeva 2003, Grozeva et al. 2006, 2007, Grozeva and Simov 2008a, b, 2009, Kuznetsova et al. 2011). The mirid bugs share some cytogenetic characteristics with all the Heteroptera: they possess holokinetic (or holocentric) chromosomes and most of them are characterized by an inverted sequence of reductional and equational division of the sex chromosomes (post-reduction) in male meiosis (Ueshima 1979). On the other hand, they have some unique chromosomal characteristics. Chiasmata are absent in male meiosis, the achiasmatic meiosis being of a collochore type (Nokkala and Nokkala 1986, Kuznetsova et al. 2011). In the three hitherto studied *Macrolophus* Fieber, 1858 species, *M. costalis* Fieber, 1858, *M. pygmaeus* (Rambur, 1839) and *M. geranii* Josifov, 1961, both autosomes and sex chromosomes divide pre-reductionally during the achiasmatic male meiosis (Grozeva et al. 2006, 2007).

The species from the present study, *M. costalis*, *M. melanotoma* (A. Costa, 1853), and *M. pygmaeus*, occur on a variety of plant species in the Mediterranean region. *M. pygmaeus* is an efficient predator of several key vegetable crop pests in Europe produced commercially and used widely as a biocontrol agent (Alomar et al. 2002, 2006, van Lenteren 2003, Messelink et al. 2014). *M. pygmaeus* and *M. melanotoma* are cryptic species with great morphological similarity which results in their misidentification and negative consequences for the conservation of their populations on greenhouse and outdoor crops. In order to find specific markers for their separation we here studied the karyotype, male meiosis and heterochromatin composition of these species and additionally of a third species, *M. costalis*. The species have recently been separated based on differences of their genetic profiles, cuticular hydrocarbon composition and on the fact that interspecies crosses do not produce viable progeny (Martinez-Cascales et al. 2006, Gemeno et al. 2012, Castañé et al. 2013). *M. costalis* can be easily distinguished morphologically from *M. pygmaeus* or *M. melanotoma* by the black dot on the scutellum, but it was included in our study as a reference species. In earlier cytogenetic studies (Grozeva et al. 2006, 2007), karyotype of two of the three species here examined was reported. Such characters, as highly asymmetric karyotype ($2n=24+X_1X_2X_3Y$) with two extra-large autosome pairs and interstitial distribution of C-heterochromatin in them (Grozeva et al. 2006), provide excellent cytogenetic markers to distinguish *M. costalis* from other *Macrolophus* species. The karyotype of *M. pygmaeus* ($2n=26+XY$) is asymmetric, as in *M. costalis*, but with different number of autosomes and a simple XY sex chromosome system (Grozeva et al. 2007). The species share sex chromosome pre-reduction, but can easily be differentiated by their karyotype and pattern of C-heterochromatin distribution.

Sperm morphology is significant in fertilization (Evans and Simmons 2008). Franco et al. (2011) have recently shown that the species in which sperm competition occurs also displayed the longest sperm length.

With the aim of distinguishing between the cryptic *Macrolophus* species, both karyotype and male meiosis were studied for the first time in *M. melanotoma* and reinvestigated in *M. pygmaeus* and (as a reference species) in *M. costalis* using standard chromosome staining and fluorochromes DAPI and CMA₃. In addition, morphology of sperm cells was examined in each of the three species.

Material and methods

Insects

Males and females of *M. costalis*, *M. pygmaeus* and *M. melanotoma* were collected in Catalonia, NE of Spain, in the vicinity of Mataró (Barcelona) (41.556 North, 2.475 East) from *Cistus albidus* Linnaeus, 1753, commercial tomato fields and *Dittrichia viscosa* (Linnaeus) Greuter, respectively. Colonies from collected individuals were set-up under controlled conditions ($25 \pm 1^\circ\text{C}$, $70 \pm 10\%$ RH and L16:D8 photoperiod) on tobacco plants (*Nicotiana tabacum* Linnaeus, 1753) with *Ephestia kuehniella* Zeller, 1879 (Lepidoptera, Pyralidae) eggs as a prey (Agusti and Gabarra 2009a, b). All *Macrolophus* specimens were preliminarily identified following Josifov (1992). However, due to morphological similarity between *M. melanotoma* and *M. pygmaeus*, their identification was additionally tested by conventional PCR using methodology and specific primers described in Castañe et al. (2013).

Karyotype

The abdomen of 20 *M. pygmaeus*, 23 *M. melanotoma* and 13 *M. costalis* males were placed in 3:1 fixative (96% ethanol-glacial acetic mixture) and the thorax in 70% ethanol for later species identification by DNA analysis (Castañe et al. 2013). Dissected gonads were squashed in a small drop of 45% acetic acid. The cover slips were removed by the dry ice technique. Slides were dehydrated in fresh fixative (3:1) and air dried. Part of the preparations was stained using Schiff-Giemsa method of Grozeva and Nokkala (1996) to check the number of chromosomes and their behaviour in meiosis. For other slides, DNA-binding fluorochromes, GC-specific chromomycin A₃ (CMA₃) and AT-specific 4'-6'-diamino-2-phenylindole (DAPI) were applied following Schweizer (1976) and Donlon and Mafenis (1983), with minor modifications as described in Kuznetsova et al. (2001).

Chromosomes were analyzed using light/fluorescent microscopy (Axio Scope A1 – Carl Zeiss Microscope) at 100× magnification and documented with a ProgRes MFcool – Jenoptik AG digital camera. All cytogenetic preparations and remains of the specimens are stored at the Institute of Biodiversity and Ecosystem Research, BAS in Sofia.

Sperm morphology

In every species, sperm cells from 10 males (other than those used for the karyotype analysis) were measured following Franco et al. (2011). On a slide, one drop of Beadle saline solution (128.3 mM NaCl, 4.7 mM KCl, 23 mM CaCl₂) was added to the male abdomen. The seminal vesicle was extracted and opened in 20 µl of saline solution to allow the sperm out. The sperm were diluted carefully with the aid of a fine needle, and then one drop was transferred and smeared across a microscope slide, allowed to dry and rinsed. Sperm cells were analyzed using a Leica DM 4000 light microscope. Twelve sperm cells per individual were measured (head and tail length) at 400x under dark field using the QWin 6.0 (Leica Microsystems, Germany) software package. In order to reduce measurement variation, each component was measured five times for each sperm cell. Data were analyzed by a one-way ANOVA and means separation by Tukey multiple range test.

Results


Karyotype

Macrolophus costalis, 2n=27 (24+X₁X₂Y)

Published data: 2n=28 (24+X₁X₂X₃Y) (Grozeva et al. 2006)

Spermatogonial metaphases consisted of 5 large (incl. Y) and 22 similar in size chromosomes (incl. two X) (Fig. 1). In meiosis, condensation stage was most abundant and showed 12 autosomal bivalents plus a positively heteropycnotic sex chromosome body (Fig. 2). Size differences between the bivalents were observed. The complement included two extremely large bivalents, four to five times the size of the other 10 similar size bivalents. Bivalents consisted of parallel-aligned homologous chromosomes without chiasmata, i.e. the male meiosis was achiasmatic. After the Schiff-Giemsa staining, it was easy to see that conspicuous interstitial heterochromatic bands in both large bivalents divide them into the three almost equal parts. At metaphase I (MI), the sex chromosomes were seen either as a trivalent (Fig. 3a, b) or as a bivalent (Fig. 3c). They clearly segregated at anaphase I (Fig. 4a) resulting in two types of metaphase II, with 12 autosomes plus two X chromosomes (Fig. 4b) and with 12 autosomes plus the Y (Fig. 4c). Male meiosis was hence pre-reductional both for autosomes and sex chromosomes. After second division, every cell produced daughter cells possessing either 2X or Y chromosome respectively (Fig. 5). Thus, the chromosome formula of male *M. costalis* was determined as 2n=27 (24+X₁X₂Y) in contrast to 2n=28 (24+X₁X₂X₃Y) earlier reported in Grozeva et al. (2006).

After staining with fluorochromes, bright DAPI- and CMA₃- positive bands were observed in the same locations on the larger autosomal bivalents and sex chromosomes (Fig. 6a–d).



Figures 1–6. Male meiosis in *Macrolophus costalis*. (1–5 conventional staining 6 fluorochrome staining: 6a, c CMA₃ 6b, d DAPI) 1 spermatogonial metaphase 2 condensation stage 3a–c metaphase I 4a–c metaphase II 5 telophase 6a, b condensation stage 6c, d metaphase I. Sex-chromosomes are indicated by arrowheads. Heterochromatin blocks are indicated by arrows. Bar = 10 µm.

***Macrolophus pygmaeus*, 2n=28 (26+XY)**

Published data: 2n=28 (26+XY) (Grozeva et al. 2007)

Spermatogonial metaphases consisted of 5 large (incl. X) and 23 similar size chromosomes (incl. Y) (Fig. 7). In meiosis, condensation stage showed 13 autosomal bivalents and a positively heteropycnotic sex chromosome body (Fig. 8). The complement included two extremely large bivalents (~five times the size of the others), and eleven bivalents similar in size. The bivalents consisted of parallel-aligned homologous chromosomes; chiasmata were absent and the male meiosis was achiasmatic. MI was nonradial (i.e., the autosomes did not form a ring), and the sex chromosomes formed a pseudobivalent (Fig. 9). The sex chromosomes segregated at AI resulting in two MII cells each with 14 chromosomes (13 + X or Y) (Fig. 10). Male meiosis was hence pre-reductional both for autosomes and sex chromosomes. Thus, the chromosome formula of *M. pygmaeus* was confirmed as 2n=28 (26+XY) in line with that earlier reported in Grozeva et al. (2007).

After staining with fluorochromes, bright DAPI- and CMA₃-positive bands were observed on both sex chromosomes (Fig. 11a–d). In addition, a weak DAPI- positive / CMA₃-negative signal was registered in a telomere of one of the larger bivalents (Fig. 11a, b).

***Macrolophus melanotoma* 2n=34 (32+XY)**


Published data: absent

At spermatogonial metaphase, there were 34 chromosomes (Fig. 12) gradually decreasing in size and the sex chromosomes were difficult to distinguish. At meiotic condensation stage, autosomal bivalents consisted of parallel lying homologs, and the sex chromosomes appeared as a heteropycnotic body (Fig. 13). At MI, there were 16 autosomal bivalents and X and Y chromosomes (Fig. 14a–c). Note that both sex chromosomes were occasionally placed in the center of a ring formed by autosomal bivalents (Fig. 14b). The autosomal bivalents constituted a decreasing size series and the X was more than twice the size of the Y. As a result of pre-reductional division of sex chromosomes at AI (Fig. 15), two types of MII (Fig. 16) raised, both with 17 chromosomes while with X or Y chromosome respectively.


After staining with fluorochromes, bright DAPI- and CMA₃ bands were observed on the sex chromosomes (Fig. 17).

Sperm morphology

Sperm cells of the species studied were of similar shape, with a long and filiform head (Fig. 18). However, species are different in the total length (mean ± SE) of the sperm cells ($F_{2,27}=15.53$; $P<0.0001$), in the length of head ($F_{2,27}=38.89$; $P<0.0001$) and tail ($F_{2,27}=25.43$; $P<0.01$) (Table 1).



Figures 7–11. Male meiosis in *Macrolophus pygmaeus*. (7–10 conventional staining 11 fluorescence staining: 11a, c CMA₃ 11b, d DAPI) 7 spermatogonial metaphase 8 condensation stage 9 metaphase I 10 metaphase II 11a, b metaphase I 11c, d anaphase. Sex-chromosomes are indicated by arrowheads. CMA₃/DAPI signals are indicated by arrows. Bar = 10 μ m.



Figures 12–17. Male meiosis in *Macrolophus melanotoma*. (12–16 conventional staining 17 fluorescence staining: 17a CMA₃ 17b DAPI) 12 spermatogonial metaphase 13 condensation stage 14a–c metaphase I 15 anaphase I 16 metaphase II 17 condensation stage. Sex-chromosomes are indicated by arrowheads. Bar = 10 μ m.

Table I. Sperm cells length (means \pm standard error) of *M. costalis*, *M. melanotoma* and *M. pygmaeus*. Within each row, means followed by the same letter do not differ significantly ($P \leq 0.005$).

Sperm length (μ m)	<i>M. costalis</i>	<i>M. melanotoma</i>	<i>M. pygmaeus</i>
Total	236 \pm 0.9 a	220.4 \pm 0.9 b	215 \pm 1.4 c
Head	61.3 \pm 0.4 a	51.3 \pm 0.4 b	50.6 \pm 0.4 b
Tail	174.7 \pm 0.8 a	169.1 \pm 1 b	164.6 \pm 1.2 c

Discussion

The genus *Macrolophus* belongs to the tribe Dicyphini of the subfamily Bryocorinae (Heteroptera, Miridae). In Bryocorinae, besides the modal for the Miridae chromosome number $2n=34$ ($32+XY$), some higher ($2n=36+XY$ and $2n=46+XY/X_1X_2Y$) and




Figure 18. Sperm morphology of *Macrolophus* species. The arrow shows the end of sperm head.

lower ($2n=16-26+XY$) chromosome numbers have also been described (Ueshima 1979, Grozeva 2003, Grozeva et al. 2006, 2007, 2008a). The chromosome formula of the reference species *M. costalis* was reported earlier as $2n=28 (24+X_1X_2X_3Y)$ for a Bulgarian population collected from the tobacco plants, which was the first report of three X chromosomes in the family (Grozeva et al. 2006). In contrast, the chromosome formula of the NE Spanish population here studied appeared to be different, with two instead of three X chromosomes. As the specimens come from different geographic regions we could speculate that they represent chromosomal races within a species. At taxonomic level, they may probably be considered as subspecies, but to clarify this hypothesis a chromosomal analysis of individuals (males and females) from natural populations of this species over the whole distribution range will be necessary. *M. pygmaeus* showed $2n=28 (26+XY)$ both in Bulgaria (Grozeva et al. 2007) and Spain (present study). The third species, *M. melanotoma*, studied here for the first time, appeared to differ from *M. costalis* and *M. pygmaeus* in karyotype. Besides difference in the chromosome number, *M. melanotoma* lacks two large autosome pairs characteristic of the two other species allowing for the cryptic *M. melanotoma* and *M. pygmaeus* to be reliably differentiated.

Differences in molecular organisation of chromatin revealed after fluorochrome staining, suggest an additional chromosome marker to differentiate *Macrolophus* species. In *M. costalis*, bright DAPI/CMA₃ bands were observed in the same locations on the large autosomal bivalents and sex chromosomes whereas in two remaining species bright fluorescent bands were observed only on the sex chromosomes. In turn, *M. pygmaeus* differed from *M. melanotoma* in that it showed some additional weak DAPI-positive signals in a telomeric region of a larger bivalent.

Franco et al. (2011) have recently reported data on sperm morphology in *M. pygmaeus* and another dicyphine species *Nesidiocoris tenuis* Reuter, 1895. *M. pygmaeus* males were shown to have significantly smaller sperm cells (213.18 μm), with longer (50.94 μm) and wider heads and shorter tails (162.94 μm) than *N. tenuis*. In our study, the data on *M. pygmaeus* sperm cell size were confirmed. On the other hand, we found that *M. costalis* males have significantly longer sperm cells, with a longer head and tail compared to *M. melanotoma* and *M. pygmaeus*. In turn, *M. melanotoma* males have significantly longer sperm tails compared to *M. pygmaeus* (Table 1).

Conclusion

As mentioned in Introduction, the cryptic species *M. pygmaeus* and *M. melanotoma* can be differentiated from each other based on the cuticular hydrocarbon profiles and specific molecular primers (Gemeno et al. 2012, Castañé et al. 2013). In our study, we provide some alternative characters, such as karyotype (number and size of chromosomes, sex chromosome system, and amount and distribution of heterochromatin) and sperm cells' morphology, allowing for reliable identification of *M. pygmaeus*, *M. melanotoma* and *M. costalis*.

Acknowledgments

This study has been funded by the Spanish Ministry of Economy and Competitiveness (MINECO) (Project AGL2011-24349), and by a travel grant from the University of Lleida - Fundació "La Caixa" to Dr Grozeva. The chromosome analysis was performed using microscope Axio Scope A1 – Carl Zeiss Microscopy upgraded by the project WETLANET (FP7 CSA – SUPPORT ACTION, GA 229802). We thank cordially Prof. Dr V. Kuznetsova for the valuable advices to improve the manuscript.

References

- Agustí N, Gabarra R (2009a) Puesta a punto de una cría masiva del depredador polífago *Dicyphus tamaninii* Wagner (Heteroptera: Miridae). Boletín de Sanidad Vegetal (PLAGAS) 35: 205–218.
- Agustí N, Gabarra R (2009b) Effect of adult age and insect density of *Dicyphus tamaninii* Wagner (Heteroptera: Miridae) on progeny. Journal of Pest Science 82(3): 241–246. doi: 10.1007/s10340-009-0245-1
- Alomar O, Goula M, Albajes R (2002) Colonisation of tomato fields by predatory mirid bugs (Hemiptera: Heteroptera) in northern Spain. Agriculture, Ecosystems and Environment 89: 105–115. doi: 10.1016/S0167-8809(01)00322-X

- Alomar O, Riudavets J, Castañe C (2006) *Macrolophus caliginosus* in the biological control of *Bemisia tabaci* on greenhouse melons. *Biological Control* 36(2): 154–162. doi: 10.1016/j.biocontrol.2005.08.010
- Castañe C, Agustí N, Arnó J, Gabarra R, Riudavets J, Comas J, Alomar O (2013) Taxonomic identification of *Macrolophus pygmaeus* and *Macrolophus melanotoma* based on morphometry and molecular markers. *Bulletin of Entomological Research* 103(2): 204–215. doi: 10.1017/S0007485312000545
- Donlon TA, Mafenis RE (1983) Methyl green is a substitute for distamycin A in the formation of distamycin A/DAPI C-bands. *Human Genetics* 65: 144–146. doi: 10.1007/BF00286651
- Evans JP, Simmons LW (2008) The genetic basis of traits regulating sperm competition and polyandry: can selection favour the evolution of good- and sexy-sperm? *Genetica* 134: 5–19. doi: 10.1007/s10709-007-9162-5
- Franco K, Jauset A, Castañe C (2011) Monogamy and polygamy in two species of mirid bugs: a functional-based approach. *Journal of Insect Physiology* 57: 307–315. doi: 10.1016/j.jinsphys.2010.11.020
- Gemeno C, Laserna N, Riba M, Valls J, Castañe C, Alomar O (2012) Cuticular hydrocarbons discriminate cryptic *Macrolophus* species (Hemiptera: Miridae). *Bulletin of Entomological Research* 102: 624–631. doi: 10.1017/S0007485312000193
- Grozeva S (2003) Karyotype of three endemic Mediterranean Miridae species (Heteroptera) from Bulgaria. *Acta Zoologica Bulgarica* 10 (3): 19–27.
- Grozeva S, Nokkala S (1996) Chromosomes and their meiotic behaviour in two families of the primitive infraorder Dipsocoromorpha (Heteroptera). *Hereditas* 125: 31–36. doi: 10.1111/j.1601-5223.1996.t01-1-00031.x
- Grozeva S, Nokkala S, Simov N (2006) First evidence of sex chromosomes pre-reduction in male meiosis in the Miridae bugs (Heteroptera). *Folia Biologica (Kraków)* 54 (1-2): 9–12. doi: 10.3409/173491606777919166
- Grozeva S, Simov N, Josifov M (2007) Karyotaxonomy of some European *Macrolophus* species (Heteroptera, Miridae). *Mainzer Naturwissenschaftliches Archiv/Beiheft* 31: 81–87.
- Grozeva S, Simov N (2008a) Cytogenetic Studies of Bryocorinae Baerensprung, 1860 true bugs (Heteroptera, Miridae). *Acta Zoologica Bulgarica*, Suppl 2: 61–70.
- Grozeva S, Simov N (2008b) Cytotaxonomy of two *Cremnocephalus* species (Heteroptera, Miridae). In: Grozeva S, Simov N (Eds) *Advances in Heteroptera Research. Festschrift in Honour of 80th Anniversary of Michail Josifov*, 171–179.
- Grozeva S, Simov N (2009) Cytogenetic study of two species of *Dryophilocoris*: *D. luteus* and *D. flavoquadrimaculatus* (Insecta, Heteroptera, Miridae). *Genetics and Breeding* 38(1): 41–45.
- Josifov M (1992) Zur Taxonomie der paläarktischen *Macrolophus*-Arten (Insecta, Heteroptera: Miridae). *Reichenbachia* 29(1): 1–4.
- Kuznetsova VG, Westendorff M, Nokkala S (2001) Patterns of chromosome banding in the sawfly family Tenthredinidae (Hymenoptera, Symphyta). *Folia biologica (Kraków)* 54(3): 227–233. doi: 10.1080/00087114.2001.10589230

- Kuznetsova VG, Grozeva S, Nokkala S, Nokkala C (2011) Cytogenetics of the true bug infraorder Cimicomorpha (Hemiptera, Heteroptera): a review. *ZooKeys* 154: 31–70. doi: 10.3897/zookeys.154.1953
- Martinez-Cascales JI, Cenis JL, Cassis G, Sánchez JA (2006) Species identity of *Macrolophus melanotoma* (Costa 1853) and *Macrolophus pygmaeus* (Rambur 1839) (Insecta: Heteroptera: Miridae) based on morphological and molecular data and bionomic implications. *Insect Systematics and Evolution* 37: 385–404. doi: 10.1163/187631206788831470
- Messelink GJ, Bloemhard CMJ, Hoogerbrugge H, Van Schelt J, Ingegno BL, Tavella L (2014) Evaluation of mirid predatory bugs and release strategy for aphid control in sweet pepper. *Journal of Applied Entomology* 139(5): 333–341. doi: 10.1111/jen.12170
- Nokkala S, Nokkala C (1986) Achiasmatic male meiosis of collochore type in the heteropteran family Miridae. *Hereditas* 105: 193–197. doi: 10.1111/j.1601-5223.1986.tb00661.x
- Schuh RT (1995) Plant bugs of the world (Insecta: Heteroptera: Miridae). Systematic Catalog Distributions, Host list, and Bibliography. The New York Entomological Society, xii+1329 pp.
- Schweizer D (1976) Reverse fluorescent chromosome banding with chromomycin and DAPI. *Chromosoma* 58: 307–324. doi: 10.1007/BF00292840
- Ueshima N (1979) Insecta 6. Hemiptera II. Heteroptera. In: John B. Animal Cytogenetics. Gebrüder Borntraeger, Berlin, Stuttgart, 1–118.
- van Lenteren JC (2003) Commercial availability of biological control agents. In: van Lenteren JC (Ed.) *Quality Control and Production of Biological Control Agents, Theory and Testing Procedures*. Oxon, UK, CABI Publishing, 167–179. doi: 10.1079/9780851996882.0167