

Karyotype analysis of four jewel-beetle species (Coleoptera, Buprestidae) detected by standard staining, C-banding, AgNOR-banding and CMA3/DAPI staining

Gayane Karagyan¹, Dorota Lachowska², Mark Kalashian¹

¹ Institute of Zoology of Scientific Center of Zoology and Hydroecology, National Academy of Sciences of Armenia, P. Sevak 7, Yerevan 0014, Armenia ² Department of Entomology, Institute of Zoology Jagiellonian University, Ingardena 6, 30-060 Krakow, Poland

Corresponding author: Gayane Karagyan (gaykaragyan@yahoo.com)

Academic editor: Robert Angus | Received 15 February 2012 | Accepted 17 April 2012 | Published 27 April 2012

Citation: Karagyan G, Lachowska D, Kalashian M (2012) Karyotype analysis of four jewel-beetle species (Coleoptera, Buprestidae) detected by standard staining, C-banding, AgNOR-banding and CMA3/DAPI staining. *Comparative Cytogenetics* 6(2): 183–197. doi: 10.3897/CompCytogen.v6i2.2950

Abstract

The male karyotypes of *Acmaeodera pilosellae persica* Mannerheim, 1837 with $2n=20$ ($18+neoXY$), *Sphenoptera scovitzii* Faldermann, 1835 ($2n=38-46$), *Dicerca aenea validiuscula* Semenov, 1895 – $2n=20$ ($18+Xy_p$) and *Sphaerobothris aghababiani* Volkovitsh et Kalashian, 1998 – $2n=16$ ($14+Xy_p$) were studied using conventional staining and different chromosome banding techniques: C-banding, AgNOR-banding, as well as fluorochrome Chromomycin A₃ (CMA₃) and DAPI. It is shown that C-positive segments are weakly visible in all four species which indicates a small amount of constitutive heterochromatin (CH). There were no signals after DAPI staining and some positive signals were discovered using CMA₃ staining demonstrating absence of AT-rich DNA and presence of GC-rich clusters of CH. Nucleolus organizing regions (NORs) were revealed using Ag-NOR technique; argentophilic material mostly coincides with positive signals obtained using CMA₃ staining.

Keywords

Coleoptera, Buprestidae, karyotypes, Ag-banding, C-banding, CMA₃-staining, DAPI-staining

Introduction

The family Buprestidae (Coleoptera) is one of the largest groups of Polyphagan beetles containing over than 14500 nominal species worldwide (Bellamy 2008a-d, 2009). Until now, karyotypes have been published for 88 species (34 from Armenia) of jewel-

beetles belonging to 22 genera and 14 tribes of the subfamilies Julodinae, Polycestinae, Chrysochroinae, Buprestinae and Agrilinae (Smith and Virkki 1978; Karagyan and Kuznetsova 2000; Karagyan 2001; Karagyan et al. 2004; Karagyan and Lachowska 2007; Moura et al. 2008). In the works listed the data were obtained by study of gonads of imagos. Only one work was based on the study of hemocytes of the larva of *Chalcophora mariana* Linnaeus, 1758 (Chrysochroinae, Chrysochroini) with diploid chromosome number $2n=22$ (Baragaño 1978; this work was omitted in all previous reviews). The diploid chromosome numbers ($2n$) in the family Buprestidae range between 12 and 46. The modal number is $2n=20$ found in 16 species, 8 genera, 6 tribes and 4 subfamilies. The most frequent sex chromosome system is XX/XY. The XY system of males is diverse (X_{yp} , X_{yr} , “XY”, neo-XY and multiple X- and Y- sex chromosomes). The X_{yp} “parachute” type is the most common and occurs in 64 species, 15 genera, 10 tribes and 4 subfamilies.

Up till now, cytogenetic studies on jewel-beetles have been carried out using conventional staining techniques mainly, with few exceptions. Karagyan (2001) reported the first data on AgNOR- staining in the karyotypes of two species of the genus *Sphenoptera* Dejean, 1833 and four species of *Acmaeoderella* Cobos, 1955, and Moura et al. (2008) studied karyotype of *Euchroma gigantea* Linnaeus, 1758 using Ag-banding, C-banding and fluorescent *in situ* hybridization (FISH) with a rDNA probe.

The present paper is dedicated to study of karyotypes of *Acmaeodera pilosellae persica* Mannerheim, 1837, *Sphenoptera scovitzii* Faldermann, 1835, *Dicercia aenea validiuscula* Semenov, 1895 and *Sphaerobothris aghababiani* Volkovitsh & Kalashian, 1998 using conventional and differential staining techniques. AgNO₃-banding was used to reveal the nucleolus organizing chromosomes and to locate the NORs in them. C-banding method was used to study the distribution of constitutive heterochromatin (CH). To characterize the molecular composition of the CH chromosomes were stained with DNA-specific fluorochromes DAPI and CMA₃ which selectively stain AT-rich and GC-rich DNA fragments, respectively. In the present study the fluorochromes CMA₃ and DAPI were applied for the first time to jewel-beetles chromosomes.

Material and methods

The buprestid beetles were collected in 2006 in Southern Armenia (Vayotsdзор and Syunik provinces). Males' gonads were dissected in several drops of hypotonic 0.9% sodium citrate solution containing 0.005% colchicine and incubated for 30–45 min at room temperature. Then the gonads were fixed in 3:1 ethanol-acetic acid mixture. Karyological slides were made according to the method proposed by Rožek (1994) with minor modifications (Rožek and Lachowska 2001). Slides were examined with phase contrast optics and the best of them with well spread chromosomes were chosen for further treatments. Depending on the specific task, various staining techniques were used. First, to determine the number and morphology of chromosomes, karyo-

types were examined using 4% Giemsa solution in phosphate buffer (pH 6.8). Then a series of chromosome banding techniques was applied as follows:

The Ag-banding method (Howell and Black 1980)

The method is based on selective silver staining of NORs containing clusters of functional rRNA genes. The slides were subjected to hydrolysis in 2N formic acid for 10 min, rinsed in running water and dried. Then the slides were treated by putting of 4-5 drops of 50% aqueous silver nitrate (AgNO_3) solution and 2 drops of colloidal developer solution (0.2 g gelatin, 10 ml distilled water and 0.1 ml concentrated formic acid – HCOOH). The slides were covered with a coverslip and incubated on a hotplate for 3-4 min at 60°C in a moist chamber (warmed beforehand). After rinsing in distilled water, the slides were dried and stained by 4% Giemsa solution in phosphate buffer (pH 6.8) for 10-20 sec.

The C-banding method (Rožek 2000)

The slides were treated for 1-3 min. in 0.2N HCL at room temperature then rinsed in distilled water. Thereafter the slides were placed in 5% Ba(OH)_2 solution at 20°C for approximately 3 min, then carefully rinsed in streaming and distilled water, and for 1 min in 2xSSC solution (0.3 M sodium chloride containing 0.03 M tri-sodium citrate). Then the slides were incubated in 2xSSC solution at 50°C for 1 hr and rinsed in distilled water. Some of these slides were stained with 4% Giemsa solution in phosphate buffer (pH 6.8) and the others were dyed with DNA-specific fluorochromes DAPI and CMA_3 .

To characterize the molecular composition of the constitutive heterochromatin, we stained the chromosomes with DNA-specific fluorochromes CMA_3 (the antibiotic chromomycin A_3) and **DAPI** (4'-6-diamidino-2-phenylindol) that selectively stain GC-rich and AT-rich DNA fragments, respectively (Schweizer 1976; Donlon and Magenis 1983). Staining with fluorochromes was performed after C-banding procedure as described above. After 2xSSC treatment slides were subsequently rinsed in distilled water and dried. Then slides were immersed in McIlvaine buffer (pH 7.0) for 5 min. After this, slides were stained with CMA_3 at a final concentration of 5 $\mu\text{g}/\text{ml}$ in 10 mM McIlvaine buffer at pH 7.0 (25 μl of 96% methanol was added to 500 μl of final staining solution) for 30 min, rinsed in buffer and stained with DAPI at a final concentration of 0.4 $\mu\text{g}/\text{ml}$ in 10 mM McIlvaine buffer (pH 7.0) for 5 min. After staining, slides were rinsed in distilled water and mounted in anti-fade medium consisting of 1% n-propylgallate in a 10 M McIlvaine buffer (pH 7.0) solution with 70% glycerol. To improve the fluorochrome staining, 0.5% methanol was added to the fluorescent dye (Kuznetsova et al. 2001).

The slides were analyzed and photographed with a Nikon Eclipse 400 light microscope and CCD DS-U1 camera using the software Lucia Image 5.0.

Data resources

The data underpinning the analyses reported in this paper are deposited in the Dryad Data Repository at doi: 10.5061/dryad.hc52qb12.

Results

Subfamily Polycestinae

Tribe Acmaeoderini

Acmaeodera pilosellae persica, $2n=20$, $n \text{ ♂} = 9 + \text{neo-XY}$.

In prometaphase I/metaphase I nine autosomal bivalents formed a series of gradually decreasing sizes and large heteromorphic neo-XY sex-bivalent were observed (Fig. 1A).




Figure 1A–D. *Acmaeodera pilosellae persica*, $n=9\text{AA} + \text{neo-XY}$. **A** prometaphase I/metaphase I **B** late diakinesis (Ag-banding) **C** prometaphase I (DAPI staining) **D** (CMA₃ staining). Bar = 10 μm .

In late diakinesis / prometaphase I the 5–6 rod-shaped bivalents have most likely a terminal chiasma and 3–4 ring-shaped bivalents – two chiasmata (Fig. 1B). The X-chromosome seems to be submetacentric, Y-chromosome is most probably acrocentric and similar in size to the shorter arm of X-chromosome.

The C-banding technique revealed a very small amount of constitutive heterochromatin in the majority of the chromosomes (most probably, in pericentromeric regions) which do not form any distinct blocks. The DAPI staining did not reveal any positive signal and therefore absence of AT-rich clusters of DNA was demonstrated (Fig. 1C). Application of CMA₃ staining revealed small positive signals on telomeric regions of both homologues of two or sometimes three rod-shaped autosomal bivalents. Besides, there is small ring-shaped bivalent which is brightly visible in the majority of photographs (Fig. 1D). These signals correspond to argentophilic material revealed by the AgNOR-banding technique (Fig. 1B) and probably associated with NORs. Besides, the Y-chromosome of neo-XY bivalent is dyed argentophilic.

Subfamily Chrysochroinae

Tribe Sphenopterini

Sphenoptera scovitzii, 2n=38–46.

The karyotype of *Sph. scovitzii* studied using Ag-banding was published earlier (Karagyan 2001) and the data on karyotype of this species obtained in that study were confirmed. Unfortunately, until now, the male diploid karyotype of the species can not be determined with certainty. Thus, it seems that karyotype consists of 38–46 chromosomes, most probably of 46; the sex chromosomes could not be identified.

Some of chromosomes have a very small amount of constitutive heterochromatin weakly visible in pericentromeric regions and do not form distinct blocks. The DAPI staining of chromosomes did not reveal any positive signal (Fig. 2A), yet fluorescence after CMA₃ staining was discovered (Fig. 2B). In metaphase I, three or sometimes four rod-shaped bivalents showed distinct CMA₃ positive signals on telomeric regions of both homologues. These signals are quite stable in the largest and in two of the middle-sized bivalents and correspond to Ag-positive material revealed by the AgNOR-banding technique (Fig. 2C). Weak CMA₃ positive signals were also at times visible in some other small bivalents.

Tribe Dicerini

Dicerca aenea validiuscula, 2n=20 (18+X_y_p).

The male mitotic metaphase displayed 20 chromosomes including 9 autosomal pairs and X- and Y- sex chromosomes (Fig. 2D). All autosomes were biarmed: one pair large




Figure 2A–F. **A–C** *Sphenoptera scovitzii*, n=19-23, metaphase I. **A** DAPI staining; **B** CMA₃ staining; **C** Ag-banding **D–F** *Dicerca aenea validiuscula*, n=9AA+X_{yp}, **D** mitotic metaphase (CMA₃ staining) **E** mitotic metaphase (DAPI staining) **F** diakinesis/prometaphase I (Ag-banding). Bar = 10 μm.

and 5 pairs of middle-sized metacentrics, 2 pairs of middle-sized submetacentrics and 1 pair of middle-sized subtelocentric. The X-chromosome was middle-sized and acrocentric, Y-chromosome was dot-like with unclear morphology. In mitotic metaphase

CMA₃ positive signals were found in the pericentromeric region of long arm of middle-sized homologous pair of subtelocentric autosomes (Fig. 2D). The DAPI staining did not reveal any positive signal (Fig. 2E).

In diakinesis/prometaphase I nine autosomal bivalents and heteromorphic sex-bivalent most probably of “parachute” Xy_p type were observed (Figs 2F, 3A). The autosomal bivalents formed a series of gradually decreasing sizes. There were 5–6 ring-shaped autosomal bivalents with two chiasmata, 1–2 rod-shaped bivalents possessed most likely one terminal chiasma and 1–2 were cross-shaped with an interstitial chiasma. The heterovalent Xy_p was rather small. In the middle-sized rod-shaped bivalent (which, most likely, formed by subtelocentric autosomes) the CMA₃ positive signals were visible in terminal regions of both homologues (Fig. 3A). At this stage staining by AgNOR-banding revealed Ag-positive signals on the same bivalent, besides the sex bivalent was nearly homogenously argentophilic (Fig. 2F).

Metaphase II showed 10 chromosomes of which 7 chromosomes were meta- and submetacentric, one of chromosome was subtelocentric (which showed CMA₃ positive signal on pericentromeric region), whereas the morphology of other chromosomes was vague (Figs 3B, 3C). The C-banding revealed weakly visible constitutive heterochromatin localized in the pericentromeric region of the same subtelocentric chromosome (Fig. 3C).

Subfamily Buprestinae

Tribe Chrysobothrini

Sphaerobothris aghababiani, 2n=16 (14+Xy_p).

In diakinesis/prometaphase I seven autosomal bivalents and a sex chromosome heterovalent of the Xy_p type were observed (Fig. 3D). The bivalents gradually decreased in size. The Xy_p sex heterovalent was smallest element in the set. The majority of autosomal bivalents were rod-shaped, but in some cells one or two ring-shaped autosomal bivalents were observed.

The C-banding revealed a small amount of constitutive heterochromatin most probably in the pericentromeric regions and do not form distinct blocks. The DAPI staining did not reveal any positive signals (Fig. 3E). In metaphase I bright and distinct CMA₃ positive signal was visible on “parachute” sex heterovalent, probably connected with the Y-chromosome (Fig. 3F). In addition, weak CMA₃ positive signals were observed, as a rule they were situated in telomeric or near telomeric regions of some of rod-shaped autosomal bivalents. Ag-banding revealed the bright argentophilic material connected with whole “parachute” type sex heterovalent (Fig. 3D). Small and weak argentophilic blocks were visible in telomeric or near telomeric regions of some autosomal bivalents.




Figure 3A–F. **A–C** *Dicercia aenea validiuscula*, **A** diakinesis/prometaphase I (CMA₃ staining) **B** metaphase II (CMA₃ staining) **C** metaphase II (C-banding) **D–F** *Sphaerobothris aghababiani*, $n=7AA+X_{yp}$ **D** diakinesis/prometaphase I (Ag-banding) **E** metaphase I (DAPI staining) **F** metaphase I (CMA₃ staining). Bar = 10 μ m.

Discussion

In total, the karyotypes of 92 species of jewel-beetles belonging to 23 genera, 14 tribes of 5 subfamilies are presently described. Generalization of data including new ones shows that the modal diploid chromosome number in Buprestidae is $2n=20$ (9AA + X- and Y- sex chromosome heterovalent in males) so far found in 18 species belonging

to 8 genera, 6 tribes, 4 subfamilies. The Xy_p type of sex chromosome heterovalent is modal and occurs in 66 species, 16 genera, 10 tribes, 4 subfamilies. The most common karyotype $2n=20$ ($19+Xy_p$) can be considered as modal. The new data confirm modality of this karyotype within the family. This karyotype occurs in a large number of beetles from different families (Smith and Virkki 1978; James and Angus 2007; Schneider et al. 2007; Cabral-de-Mello et al. 2008; et al.).

Application of C-banding technique showed that the studied species of jewel-beetles are characterized by a small amount of heterochromatin localized, most probably, in pericentromeric regions of chromosomes, as in most studied species of the order Coleoptera (Smith and Virkki 1978; Rožek et al. 2004; Lachowska et al. 2009; Arcanjo et al. 2009; et al.). On the other hand, in *Euchroma gigantea* large C-blocks were observed in multiple sex chromosomes ($X_1X_2X_3-Y_1Y_2Y_3$) (Moura et al. 2008). The presence of large C-blocks in chromosomes is uncommon in Coleoptera as a whole and has been observed in a few species only (Juan et al. 1991; Rožek 1992; Rožek and Rudek 1992; Plohl et al. 1993; Rožek and Lachowska 2001, 2003; Rožek et al. 2004; Bione et al. 2005a; Cabral-de-Mello et al. 2010).

Until now, the karyotypes of only seven species of jewel-beetles studied by using Ag-NOR-staining technique have been published (Karagyan 2001; Moura et al. 2008). In the present paper Ag-banding was applied for study of karyotypes of another three species.

Even this restricted material showed noticeable variability of distribution of argentophilic material (probably NOR) in the karyotypes of jewel-beetles. The argentophilic material is located on:

- 1) the autosomes: *Sphenoptera scovitzii* ($2n=38-46$), *Sph. mesopotamica* Marseul, 1865 ($2n=24$, Xy_p).
- 2) both on the sex chromosomes and on the autosomes: *Acmaeoderella villosula* Steven, 1830 (described as *Acmaeoderella boryi* Brullé, 1832 in Karagyan 2001) ($2n=18$, Xy_p), *Acmaeodera pilosellae persica* ($2n=20$, neo-XY), *Sphaerobothris aghababiani* ($2n=16$, Xy_p), *Dicerca aenea validiuscula* ($2n=20$, Xy_p).
- 3) sex chromosomes only, situating either on one of the sex chromosomes or localized between sex chromosomes of bivalent: *Acmaeoderella flavofasciata* (Piller & Mitterpacher, 1783) ($2n=18$, Xy_p), *A. gibbulosa* Ménétériés, 1832 ($2n=18$, Xy_p), *A. vetusta* (Ménétériés, 1832) ($2n=18$, Xy_p). In *Euchroma gigantea* ($2n=32$, 36 , $X_1X_2X_3Y_1Y_2Y_3$ – Moura et al. 2008) argentophilic material labeled the multiple sex chromosomes chain.

As commonly understood, in Coleoptera NORs may be located in some autosomal pair and/or sex chromosomes (Almeida et al. 2000; Moura et al. 2003; Bione et al. 2005b; Holecová et al. 2008). The most common pattern in Coleoptera is the location of the nucleolus organizer region in one autosomal pair (Virkki 1983; Virkki et al. 1984; Vitturi et al. 1999; Colomba et al. 2000; Moura et al. 2003; Almeida et al. 2006; Schneider 2007; et al.). On the other hand, the argentophilous body has been repeatedly observed between the sex-chromosomes of Xy_p type (Postiglioni and Brum-Zorrilla 1988; Virkki et al. 1990, 1991; Maffei et al.

2001; Moura et al. 2003; et al.). Among Buprestidae, the whole Xy_p sex bivalent is brightly argentophilic and probably bears NOR only in *Sphaerobothris aghababiani* and *Dicerca aenea validiuscula*. However, the lack of relationship between nucleolus and sex chromosome system of Xy_p has also been demonstrated for some Coleoptera including one species of jewel-beetles, *Sphenoptera mesopotamica* (Karagyan 2001). According to some authors, the “parachute” can be strongly marked by the silver nitrate during different phases of meiosis independent of whether or not the NORs are located in Xy_p bivalent (Virkki et al. 1991; Moura et al. 2003; Bione et al. 2005b; Mendes-Neto et al. 2010). This phenomenon is probably related to the presence of argentophilic substance (proteins) that theoretically facilitates the configuration and segregation of the sex chromosomes of the Xy_p system (Virkki et al. 1990, 1991; Juan et al. 1993; Petitpierre 1996; Moura et al. 2003; Bione et al. 2005a, b; Schneider et al. 2007).

Chromosome staining by DNA base specific fluorochromes has been used in cytogenetic studies of Coleoptera (Juan and Petitpierre 1989; Vitturi et al. 1999; Colomba et al. 2000; Moura et al. 2003; Schneider et al. 2007; Lachowska 2008; Mendes-Neto et al. 2010; et al) but has never before been applied to Buprestidae.

The correlation between NORs and CMA_3 bands is quite common in some insects, including beetles (Camacho et al. 1991; Vitturi et al. 1999; Colomba et al. 2000; Maffei et al. 2001; Kuznetsova et al. 2001; Araújo et al. 2002; Brito et al. 2003; Nechayeva et al. 2004). Silver staining mainly reveals transcriptionally active NORs (Sumner 1990), as opposed to fluorochrome CMA_3 staining which labels NORs independently of their activity (Schmid and Guttenbach 1988). In this study, the fluorescent signals after CMA_3 staining were positive in all four species of jewel-beetles. While in *Sphenoptera scovitzii* and *Sphaerobothris aghababiani* they were nearly fully correlated with argentophilic material observed on silver dyed chromosomes, in *Acmaeodera pilosellae persica* CMA_3 signals correlated with argentophilic blocks on 2–3 autosomal bivalents but not with Y-chromosome which was dyed argentophilic as well. Meanwhile in *Dicerca aenea validiuscula* CMA_3 signal was correlated with one of autosomal pairs when argentophilic material was revealed in the same autosomal pair as well as on sex bivalent.

Fluorochrome DAPI staining in all studied species of jewel-beetles did not reveal any particular bright regions on chromosomes which allows for preliminary suggestion that CH have no distinct AT-rich DNA clusters. In general, there is variable distribution of AT- or GC-rich clusters of CH among beetles studied by fluorochromes staining. For instance, positive CMA_3 and negative DAPI signals were found in some Elateridae (Schneider et al. 2006, 2007), and most of studied Scarabaeoidea (Vitturi et al. 1999; Moura et al. 2003; Bione et al. 2005b; Cabral-de-Mello et al. 2010). Positive DAPI signals were found only in some other Scarabaeoidea (Moura et al. *l. c.*) and in majority of the Curculionidae studied (Lachowska 2008; Lachowska et al. 2008). Rarely both DAPI and CMA_3 positive signals were revealed in Scarabaeidae (Bione et al. 2005b), in a few Curculionidae (Lachowska 2008) and Chrysomelidae (Almeida et al. 2006).

In conclusion, data of the present study offer important insights into the karyotypes characteristics of jewel-beetles which may be useful in elucidation of relationships both among the species of the family itself as well as between jewel-beetles and the representatives of other coleopteran families.

Acknowledgements

This study was supported by Józef Mianowski Fund (Poland). We would like to express our greatest gratitude to our colleagues from the Institute of Systematics and Evolution of Animals, Polish Academy of Sciences Prof. dr hab. Elżbieta Warchałowska-Śliwa, Prof. dr hab. Anna Maryańska-Nadachowska, Dr. Maria Rożek and Prof. dr hab. Adam Nadachowski for their invaluable support of this study. We gratefully acknowledge to Dr. Alexander Danchenko (Moscow State University, Moscow, Russia) for his kind help in collecting some of the specimens studied.

References

- Almeida MC, Zacaro AA, Cella DM (2000) Cytogenetic analysis of *Epicauta atomaria* (Meloidae) and *Palembus dermestoides* (Tenebrionidae) with Xy_p sex determination system using standard staining, C-bands, NOR and synaptonemal complex microspreading techniques. *Hereditas* 133: 147–157. doi: 10.1111/j.1601-5223.2000.00147.x
- Almeida MC, Campaner C, Cella DM (2006) Karyotype characterization, constitutive heterochromatin and nucleolus organizer regions of *Paranaita opima* (Coleoptera, Chrysomelidae, Alticinae). *Genetics and Molecular Biology* 29(3): 475–481. doi: 10.1590/S1415-47572006000300013
- Arcanjo AP, Cabral-de-Mello DC, Silva AEB, Moura RC (2009) Cytogenetic characterization of *Eurysternus caribaeus* (Coleoptera: Scarabaeidae): evidence of sex-autosome fusion and diploid number reduction prior to species dispersion. *Journal of Genetics* 88(2): 177–182. doi: 10.1007/s12041-009-0025-y
- Araújo SMR, Pompolo SG, Perfectti F, Camacho JPM (2002) Genetic load caused by variation in the amount of rDNA in a wasp. *Chromosome Research* 10: 607–613. doi: 10.1023/A:1020970820513
- Baragaño R (1978) Nuevo método para el estudio de cromosomas en Coleoptera a partir de hemocitos de estados larvarios. *Boletín del Servicio de Plagas* 4: 23–33.
- Bellamy CL (2008a) A world catalogue and bibliography of the jewel beetles (Coleoptera: Buprestoidea). Volume 1: Introduction; Fossil Taxa; Schizopodidae; Buprestidae; Julodinae - Chrysochroinae: Poecilonotini. Pensoft Publishers, Moscow-Sofia, 606 pp.
- Bellamy CL (2008b) A world catalogue and bibliography of the jewel beetles (Coleoptera: Buprestoidea). Volume 2: Chrysochroinae: Sphenopterini through Buprestinae: Stigmoderini. Pensoft Publishers, Moscow-Sofia, 614 pp.

- Bellamy CL (2008c) A world catalogue and bibliography of the jewel beetles (Coleoptera: Buprestoidea). Volume 3: Buprestinae: Pterobothrini through Agrilinae: Rhaeboscelina. Pensoft Publishers, Moscow-Sofia, 632 pp.
- Bellamy CL (2008d) A world catalogue and bibliography of the jewel beetles (Coleoptera: Buprestoidea). Volume 4: Agrilinae: Agrilina through Trachyini. Pensoft Publishers, Moscow-Sofia, 722 pp.
- Bellamy CL (2009) A world catalogue and bibliography of the jewel beetles (Coleoptera: Buprestoidea). Volume 5: Appendices, Bibliography, Indices. Pensoft Publishers, Moscow-Sofia, 582 pp.
- Bione E, Camparoto ML, Simões ZLP (2005a) A study of the constitutive heterochromatin and nucleolus organizer regions of *Isocopris inhiata* and *Diabroctis mimas* (Coleoptera: Scarabaeidae, Scarabaeinae) using C-banding, AgNO₃ staining and FISH techniques. Genetics and Molecular Biology 28(1): 111–116. doi: 10.1590/S1415-47572005000100019
- Bione E, Moura RC, Carvalho R, Souza MJ (2005b) Karyotype, C- and fluorescence banding pattern, NOR location and FISH study of five Scarabaeidae (Coleoptera) species. Genetics and Molecular Biology 28(3): 376–381. doi: 10.1590/S1415-47572005000300006
- Brito RM, Caixerio APA, Pompolo SG, Azevedo GG (2003) Cytogenetic data of *Partamona peckolti* (Hymenoptera, Apidae, Meliponini) by C-banding and fluorochrome staining with DA/CMA₃ and DA/DAPI. Genetics and Molecular Biology 26(1): 53–57. doi: 10.1590/S1415-47572003000100009
- Cabral-de-Mello DC, Oliveira SG, Ramos IC, Moura RC (2008) Karyotype differentiation patterns in species of the subfamily Scarabaeinae (Scarabaeidae, Coleoptera). Micron 39: 1243–1250. doi: 10.1016/j.micron.2008.04.002
- Cabral-de-Mello DC, Moura RC, Souza MJ (2010) Amplification of repetitive DNA and origin of a rare chromosomal sex bivalent in *Deltochilum (Calhyboma) verruciferum* (Coleoptera, Scarabaeidae). Genetica 138: 191–195. doi: 10.1007/s10709-009-9409-4
- Camacho JPM, Cabrero J, Viseras E (1991) G Banding in two species of grasshoppers and its relationships to C, N, and fluorescence banding techniques. Genome 34: 638–643. doi: 10.1139/g91-097
- Colomba MS, Vitturi R, Zunino M (2000) Karyotype analyzes, banding, and fluorescent *in situ* hybridization in the Scarab beetle *Gymnopleurus sturmi* McLeady (Coleoptera, Scarabaeoidea, Scarabaeidae). The Journal of Heredity 91: 260–264. doi: 10.1093/jhered/91.3.260
- Donlon TA, Magenis RE (1983) Methyl green is a substitute for distamycin A in the formation of distamycin A/DAPI C-bands. Human Genetics 65: 144–146. doi: 10.1007/BF00286651
- Holecová M, Rožek M, Lachowska D (2008) The first cytogenetic report on *Laena reitteri* Weise, 1877 (Coleoptera, Tenebrionidae, Lagriinae) with notes on karyotypes of darkling beetles. Folia biologica (Kraków) 56(3–4): 213–217.
- Howell WM, Black DA (1980) Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-srep method. Experientia 36: 1014–1015. doi: 10.1007/BF01953855
- James LV, Angus RB (2007) A chromosomal investigation of some British Cantharidae (Coleoptera). Genetica 130: 293–300. doi: 10.1007/s10709-006-9106-5

- Juan C, Gonsálves J, Mezzanotte R, Petitpierre E (1991) Cytological and biochemical characterization of the *in situ* endonuclease digestion of fixed *Tenebrio molitor* chromosomes. *Chromosoma* 100: 432–438. doi: 10.1007/BF00364553
- Juan C, Petitpierre E (1989) C-banding and DNA content in seven species of Tenebrionidae (Coleoptera). *Genome* 32: 834–839. doi: 10.1139/g89-519
- Juan C, Pons J, Petitpierre E (1993) Localization of tandemly repeated DNA sequences in beetle chromosomes by fluorescent *in situ* hybridization. *Chromosome Research* 1: 167–174.
- Karagyan G (2001) Ag-banded karyotypes in six species of jewel-beetles (Buprestidae, Coleoptera). *Folia biologica (Kraków)* 49(1–2): 42–48. doi: 10.1007/BF00710770
- Karagyan GH, Kuznetsova VG (2000) Chromosome numbers and sex chromosome systems in buprestid beetles (Coleoptera, Buprestidae). *Entomological Review* 80(1): 38–49. Translated from *Entomologicheskoe Obozrenie* 79(1): 96–106.
- Karagyan G, Kuznetsova VG, Lachowska D (2004) New cytogenetic data on Armenian buprestids (Coleoptera, Buprestidae) with a discussion of karyotype variation within the family. *Folia biologica (Kraków)* 52(3–4): 151–158. doi: 10.3409/1734916044527601
- Karagyan GH, Lachowska D (2007) New karyotypes of five species of jewel-beetles and presumptive ancestral state of karyotypes of the subfamilies Polycestinae, Chrysochroinae and Buprestinae (Coleoptera: Buprestidae). *Comparative Cytogenetics* 1(2): 121–127. (www.zin.ru/journals/compcyt)
- Kuznetsova VG, Westendorff M, Nokkala S (2001) Patterns of chromosome banding in the sawfly family Tenthredinidae (Hymenoptera, Symphyta). *Caryologia* 54(3): 227–233.
- Lachowska D (2008) Karyotypes and chromosome rearrangements in two tribes of weevils (Coleoptera, Curculionidae: Sciaphilini and Brachydrini). *Folia biologica (Kraków)* 56(3–4): 219–225. doi: 10.3409/fb.56_3-4.219-225
- Lachowska D, Rožek M, Holecová M (2008) Cytotaxonomy and karyology of the tribe Otiorhynchini (Coleoptera: Curculionidae). *European Journal of Entomology* 105: 175–184.
- Lachowska D, Rožek M, Holecová M (2009) Chromosomal similarities and differences among three sibling species of the *Acalles echinatus* group (Coleoptera, Curculionidae, Cryptorhynchinae). *Zootaxa* 1985: 63–68.
- Maffei EMD, Pompolo SG, Campos LAO, Petitpierre E (2001) Sequential FISH analysis with rDNA genes and Ag-NOR banding in the lady beetle *Olla v-nigrum* (Coleoptera, Coccinellidae). *Hereditas* 135: 13–18. doi: 10.1111/j.1601-5223.2001.00013.x
- Mendes-Neto EO, Vicari MR, Campaner C, Nogaroto V, Artoni RF, Almeida MC (2010) Cytogenetic analysis of *Astylus antis* (Perty, 1830) (Coleoptera, Melyridae): karyotype, heterochromatin and location of ribosomal genes. *Genetics and Molecular Biology* 33(2): 237–243. doi: 10.1590/S1415-47572010005000050
- Moura RC, Souza MJ, Melo NF, Lira-Neto AC (2003) Karyotypic characterization of representatives from Melolonthinae (Coleoptera, Scarabaeidae): Karyotypic analysis, banding and fluorescent *in situ* hybridization (FISH). *Hereditas* 138: 200–206. doi: 10.1034/j.1601-5223.2003.01611.x
- Moura RC, Melo NF, Souza MJ (2008) High levels of chromosomal differentiation in *Euchroma gigantea* L., 1735 (Coleoptera, Buprestidae). *Genetics and Molecular Biology* 31(2): 431–437. doi: 10.1590/S1415-47572008000300007

- Nechayeva GA, Kuznetsova VG, Nokkala S (2004) New data on the karyotype of *Pseudococcus viburni* (Sign.) (Homoptera, Coccinea). Entomological Review 84(4): 393–400.
- Petitpierre E (1996) Molecular cytogenetics and taxonomy of insects, with particular reference to the Coleoptera. International Journal of Insect Morphology and Embryology 25: 115–134. doi: 10.1016/0020-7322(95)00024-0
- Plohl M, Lucijanic-Justic V, Ugarkovic D, Petitpierre E, Juan C (1993) Satellite DNA and heterochromatin of the flour beetle *Tribolium confusum*. Genome 36: 467–475. doi: 10.1139/g93-064
- Postiglioni A, Brum-Zorrilla N (1988) Non-relationship between nucleolus and sex chromosome system Xy_p in *Chelymorpha variabilis* Boheman (Coleoptera, Chrysomelidae). Genetica 77: 137–141. doi: 10.1007/BF00057764
- Rożek M (1992) Karyological analysis of *Acupalpus (Stenolophus) teutonius* (Schränk) and *Pogonus presicus* (Chaud.) (Coleoptera, Carabidae). Folia biologica (Kraków) 40(3–4): 113–118.
- Rożek M, Rudek Z (1992) Karyotype analysis and C-banding pattern in two species of Carabid (Coleoptera, Carabidae). Folia biologica (Kraków) 40: 47–52.
- Rożek M (1994) A new chromosome preparation technique for Coleoptera (Insecta). Chromosome Research 2: 76–78. doi: 10.1007/BF01539458
- Rożek M (2000) The C-banding technique for Coleoptera (Insecta). Folia biologica (Kraków) 48(1–2): 29–31.
- Rożek M, Lachowska D (2001) C-bands on chromosomes of four beetle species (Coleoptera: Carabidae, Silphidae, Elateridae, Scarabaeidae). Folia biologica (Kraków) 49: 179–182.
- Rożek M, Lachowska D (2003) C-banded chromosomes of genus *Bembidion* Latr. (Coleoptera, Carabidae). Folia biologica (Kraków) 51: 143–146.
- Rożek M, Lachowska D, Petitpierre E, Holecová M (2004) C-bands on chromosomes of 32 beetle species (Coleoptera: Elateridae, Cantharidae, Oedemeridae, Cerambycidae, Chrysomelidae and Curculionidae). Hereditas 140: 1–10. doi: 10.1111/j.1601-5223.2004.01810.x
- Schmid M, Guttenbach M (1988) Evolutionary diversity of reverse (R) fluorescent chromosome bands in vertebrates. Chromosoma 77: 101–114. doi: 10.1007/BF00327367
- Schneider MC, Almeida MC, Rosa SP, Costa C, Cella DM (2006) Evolutionary chromosomal differentiation among four species of *Conoderus* Eschscholtz, 1829 (Coleoptera, Elateridae, Agrypninae, Conoderini) detected by standard staining, C-banding, silver nitrate impregnation, and CMA₃/DA/DAPI staining. Genetica 128: 333–346. doi: 10.1007/s10709-006-7101-5
- Schneider MC, Rosa SP, Almeida MC, Costa C, Cella DM (2007) Chromosomal similarities and differences among four Neotropical Elateridae (Conoderini and Pyrophorini) and other related species, with comments on the NOR patterns in Coleoptera. Journal of Zoological Systematics and Evolutionary Research 45(4): 308–316. doi: 10.1111/j.1439-0469.2006.00398.x
- Schweizer D (1976) Reverse fluorescent chromosome banding with chromomycin and DAPI. Chromosoma (Berl.) 58: 307–324. doi: 10.1007/BF00292840
- Smith SG, Virkki N (1978) Animal cytogenetics. 3. Insecta 5. Coleoptera. Berlin-Stuttgart, 366 pp.
- Sumner AT (1990) Chromosome banding. Unwyn Hyman, London, XIV + 434 pp.

- Virkki N (1983) Banding of *Oedionychna* (Coleoptera, Alticinae) chromosomes: C- and Ag-bands. *Journal of Agriculture of the University of Puerto Rico* 67: 221–225.
- Virkki N, Flores M, Escudero J (1984) Structure, orientation and segregation of the sex trivalent in *Pyrophorus luminosus* (Coleoptera, Elateridae). *Canadian Journal of Genetics and Cytology* 26: 326–330.
- Virkki N, Mazzella C, Denton A (1990) Staining of substances adjacent to the Xy_p sex bivalent of some weevils (Coleoptera: Curculionidae). *Journal of Agriculture of the University of Puerto Rico* 74: 405–418.
- Virkki N, Mazzella C, Denton A (1991) Silver staining of the coleopteran Xy_p sex bivalent. *Cytobios* 67: 45–63.
- Vitturi R, Colomba MS, Barbieri R, Zunino M (1999) Ribosomal DNA location in the scarab beetle *Thorectes intermedius* (Costa) (Coleoptera, Geotrupidae) using banding and fluorescent *in-situ* hybridization. *Chromosome Research* 7: 255–260. doi: 10.1023/A:1009270613012