

Multiple sex chromosome systems in howler monkeys (Platyrrhini, *Alouatta*)

Eliana Ruth Steinberg¹, Mariela Nieves¹, Marta Dolores Mudry¹

I Grupo de Investigación en Biología Evolutiva (GIBE) – Departamento de Ecología, Genética y Evolución – Facultad de Ciencias Exactas y Naturales – Universidad de Buenos Aires – IEGEBA (CONICET-UBA) – Ciudad Universitaria – Pab. II – 4º piso - Labs 43–46 - (C1428EGA) – Buenos Aires – Argentina

Corresponding author: *Eliana Ruth Steinberg* (steinberg@ege.fcen.uba.ar)

Academic editor: *A. Polyakov* | Received 29 November 2013 | Accepted 25 January 2014 | Published 25 February 2014

Citation: Steinberg ER, Nieves M, Mudry MD (2014) Multiple sex chromosome systems in howler monkeys (Platyrrhini, *Alouatta*). *Comparative Cytogenetics* 8(1): 43–69. doi: 10.3897/CompCytogen.v8i1.6716

Abstract

In light of the multiple sex chromosome systems observed in howler monkeys (*Alouatta* Lacépède, 1799) a combined cladistic analysis using chromosomal and molecular characters was applied to discuss the possible origin of these systems. Mesoamerican and South American howlers were karyologically compared. FISH analysis using the chromosome painting probes for the #3 and #15 human chromosomes was applied to corroborate the homeology of the sexual systems. We found that the HSA3/15 syntenic association, present in the sex chromosome systems of South American Howlers, is not present in those of Mesoamerican ones. The autosomes involved in the translocation that formed the sexual systems in the Mesoamerican and South American species are different, thus suggesting an independent origin. Parsimony analysis resolved the phylogenetic relationships among howler species, demonstrating utility of the combined approach. A hypothesis for the origin of the multiple sex chromosome systems for the genus is proposed.

Keywords

Multiple sex chromosome systems, combined phylogenetic analysis, FISH, Neotropical Primates, cytochrome b

Introduction

Howler monkeys (genus *Alouatta* Lacépède, 1799 of the family Atelidae) exhibit one of the widest geographic distributions recorded to date for Neotropical Primates. Their distribution extends from southern Mexico to northern Argentina (Crockett and Eisenberg 1987, Rylands 2000). They inhabit a diverse range of environments, including tropical rain forests, flood forests, gallery forests, patches of forest and deciduous and semideciduous seasonal environments (Crockett and Eisenberg 1987, Zunino et al. 2001). There remains a lack of consensus regarding both the number of species within the genus, which, depending on the author, ranges from 9 to 14 species (Rylands 2000, Groves 2001, 2005, Gregorin 2006, Rylands and Mittermeier 2009), and the phylogenetic relationships among them. This shows the complexity of the taxonomy of *Alouatta* and highlights the importance of including a larger number of variables for a more accurate characterization of the species in the genus. We adhere to the classification proposed by Groves (2001, 2005) in recognizing 10 species (*Alouatta belzebul* Linnaeus, 1766, *A. seniculus* Linnaeus, 1766, *A. sara* Elliot, 1910, *A. macconnelli* Linnaeus, 1766, *A. caraya* Humboldt, 1812, *A. palliata* Gray, 1849, *A. pigra* Lawrence, 1933, *A. guariba* Humboldt, 1812, *A. nigerrima* Lönnberg, 1941, *A. coibensis* Thomas, 1912), since it considers both morphological and genetic information.

In this context, and to contribute to the description of the phylogenetic relationships in the genus, several authors have proposed that chromosomal data can also be used as phylogenetic markers, since they are inherited as mendelian characters and are conserved within species (Sankoff 2003, Dobigny et al. 2004, Stanyon et al. 2008). Following the Maximum Parsimony criterion, karyological comparisons allow the identification of chromosomal forms shared by common ancestry.

In primates, different researchers in the last three decades have proposed chromosomal speciation as a probable evolutionary mechanism to explain the diversity observed in living species (de Grouchy et al. 1972, Seuánez 1979, Dutrillaux and Couturier 1981, Clemente et al. 1990, Stanyon et al. 2008, de Oliveira et al. 2012). In howler monkeys, species exhibit diploid numbers (2N) ranging from 44 in *Alouatta seniculus* to 58 in *Alouatta pigra*, and in a large number of species, multiple sex chromosome systems in males originated from Y-autosome translocations have been described (Table 1). The chromosomes involved in the Y-autosome translocations in *Alouatta caraya*, *Alouatta macconnelli*, *Alouatta guariba guariba* Humboldt, 1812, *Alouatta guariba clamitans* Cabrera, 1940, *Alouatta sara* and *Alouatta seniculus arctoidea* Cabrera, 1940, are homeologous to the same regions of human chromosomes #3 and #15 (Consigliere et al. 1996, 1998, Mudry et al. 2001, de Oliveira et al. 2002).

The phylogenies proposed so far for *Alouatta* have used either molecular markers (γ^1 -globin (Meireles et al. 1999), Mt ATP synt 8 and 6, Mt cyt b, CAL and RAG1 (Cortés-Ortiz et al. 2003)) or chromosomal characters (de Oliveira et al. 2002). However, the combination of different variables can improve the phylogenetic sig-

Table 1. Cytogenetic characteristics of howler monkeys (*Alouatta*).

Species	2N	Sex Chromosome Systems	References
<i>A. belzebul</i>	♀50 ♂49	X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y	Armada et al. 1987 [§]
<i>A. s. seniculus</i>	♀♂47 to 49†	XY	Yunis et al. 1976
<i>A. s. stramineus</i>	♀♂47 to 49†	X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y ₁ Y ₂	Lima and Seuánez 1991 [§]
<i>A. s. arctoidea</i>	♀44 ♂45‡	X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y ₁ Y ₂	Stanyon et al. 1995
<i>A. sara</i>	♀♂ 48 to 51†	X ₁ X ₁ X ₂ / X ₁ X ₂ Y	Minezawa et al. 1985
	♀♂50	X ₁ X ₁ X ₂ / X ₁ X ₂ Y ₁ Y ₂	Stanyon et al. 1995
<i>A. macconnelli</i>	♀♂ 47 to 49†	X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y ₁ Y ₂	Lima et al. 1990
<i>A. caraya</i>	♀♂52	XX/XY	Egozcue and De Egozcue 1966, Mudry et al. 1984, 1994
		X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y ₁ Y ₂	Rahn et al. 1996 [§] , Mudry et al. 1998 [§] , 2001 [§]
<i>A. palliata</i>	♀♂56	XX/XY	Torres and Ramírez 2003
	♀54 ♂53	X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y	Ma et al. 1976 Solari and Rahn 2005 [§]
<i>A. pigra</i>	♀♂58	X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y ₁ Y ₂	Steinberg et al. 2008 [§]
<i>A. guariba guariba</i>	♀50 ♂49	XX/XY	Koiffmann and Saldanha 1974
	♂49	X ₁ X ₂ Y	de Oliveira et al. 1995
	♀50♂49	X ₁ X ₁ X ₂ X ₂ X ₃ / X ₁ X ₂ X ₃ Y ₁ Y ₂	de Oliveira et al. 2002
<i>A. guariba clamitans</i>	♀46 ♂45	XX/XY	de Oliveira et al. 1995
		X ₁ X ₁ X ₂ X ₂ /X ₁ X ₂ Y	de Oliveira et al. 1998
		X ₁ X ₁ X ₂ X ₂ X ₃ / X ₁ X ₂ X ₃ Y ₁ Y ₂	de Oliveira et al. 2002
<i>A. nigerrima</i>	♀50	XX	Armada et al. 1987
<i>A. coibensis</i>	ND	ND	---

†These differences are due to the presence of microchromosomes (1 to 3 per nuclei); ‡Differences due to a variation in microchromosome number between sexes. ND: not yet cytogenetically characterized.

[§]Meiotic studies performed to corroborate the sex chromosome system.

nal due to the possible common shared history of different datasets. At the same time, this combination can increase the support of a tree, since different characters evolve at particular rates and will support different parts of the tree (Kluge 1989, Whittaker et al. 2007).

In the present contribution, howler species were karyologically compared and FISH analyses were carried out to corroborate the homeology of the sex chromosome systems among them. Using these data and molecular data obtained from the literature, a phylogenetic analysis combining them in a single matrix was performed.

Methods

Sampled specimens: A total of 29 adult specimens of both sexes of four species of howlers, both from captivity as well as from the wild within their natural geographical distribution, were analyzed: *Alouatta caraya* (9 ♂ and 6 ♀), *A. guariba clamitans* (1 ♂), *A. pigra* (6 ♂ and 5 ♀) and *A. palliata* (2 ♂).

The origin of the animals was as follows:

Argentina

A. caraya, 1 ♂ from Corrientes Zoo, Corrientes; 1 ♂ from Ecological Park “El Puma”, Misiones; 1 ♂ and 2 ♀ from Mendoza Zoo, Mendoza; 6 ♂ and 4 ♀ from the Black Howler Monkey Reeducational Center, La Cumbre, Córdoba.

A. g. clamitans, 1 ♂ from “Güira-Oga”, Misiones.

Mexico

A. pigra: 4 ♂ and 4 ♀ were sampled in the wild in Campeche, Yucatán Península; 2 ♂ and 1 ♀ from San Juan de Aragón Zoo, Mexico City.

A. palliata: 1 ♂ from San Juan de Aragón Zoo, Mexico City; 1 ♂ from Chapultepec Zoo, Mexico City.

Classical cytogenetic analysis

Chromosome preparation: Peripheral blood samples were collected from all animals with previously heparinized disposable syringes. Lymphocytes were cultured for 72 h at 37 °C following Mudry (1990). At least 50 metaphases were analyzed to determine the diploid number (2N) at 1000×. Metaphase spreads were treated with G-Wright banding (Steinberg et al. 2007). At least 10 G-banded metaphases with the species diploid number (2N) were photographed with a Leica DFC 340 FX camera. Chromosomes were arranged according to previously described karyotypes using Photoshop CS (Adobe) and the species assignment of each specimen was corroborated.

Analysis of homeologies: For *A. caraya* and *A. g. clamitans*, the homeologies with human chromosomes and the homeologies with the other South American howlers are well known (Consigliere et al. 1998, Mudry et al. 2001, de Oliveira et al. 2002, Stanyon et al. 2011). The G-banded chromosomes of *A. pigra* and *A. palliata* were first compared with those of *A. caraya* and *A. g. clamitans*. We took *A. caraya*'s karyotype as the reference for the comparisons with Mesoamerican howlers (Mudry et al. 2001, Szapkievich and Mudry 2003). To compare homeologies, the G-banded metaphases obtained for *A. caraya*, *A. g. clamitans*, *A. pigra* and *A. palliata* were also compared with those published for *A. g. guariba* (de Oliveira et al. 2002, Stanyon et al. 2011), *A. macconnelli* (de Oliveira et al. 2002), *A. s. arctoidea* (Consigliere et al. 1996), *A. belzebul* (Armada et al. 1987, Consigliere et al. 1998) and *A. sara* (Consigliere et al. 1996).

Cytomolecular study

FISH analysis with human chromosome painting probes #3 and #15 was used as a tool to confirm the identity of the sex chromosome systems in howlers. Whole chromosome painting probes for human chromosomes #3 (red), #15 (green), #21 (green), X (green) and Y (red) (PCT3 Cy3, PCT15 FITC, PCT21 FITC, PCTX FITC, PCTY

Cy3, LEXEL S.R.L., Buenos Aires, Argentina) were used for FISH analysis on the metaphases of *A. pigra*, *A. caraya*, *A. g. clamitans* and *A. palliata*. *Homo sapiens* (HSA) metaphases were used as a positive control of hybridization. The HSA3/21 syntenic association, considered ancestral in mammals and conserved in most primate species (Müller et al. 2000), was analyzed simultaneously as a control of synteny conservation. Human X and Y chromosomes were also tested.

FISH was performed according to the supplier's instructions (LEXEL S.R.L., Buenos Aires, Argentina). Slides were counterstained with DAPI (Sigma) and analyzed with a Leica DMLB fluorescence microscope. Chromosome images were obtained with a Leica DFC 340 FX camera. Images were processed with Image Pro-Plus 4.5 (Media Cybernetics Inc.).

Our results were compared with those previously described (Consigliere et al. 1996, 1998, Mudry et al. 2001, de Oliveira et al. 2002, Stanyon et al. 2011).

Phylogenetic analysis

Chromosomal dataset: We used data obtained from the comparisons of G-banding patterns and the analysis of chromosomal syntenic associations, both from the present study and from previous reports (Consigliere et al. 1996, 1998, García et al. 2001, 2002, Mudry et al. 2001, de Oliveira et al. 2002, Amaral et al. 2008, Stanyon et al. 2001, 2011). We considered the structural changes as characters. The pattern observed before and after their occurrence, i.e. their presence or absence, was considered as the character states. The matrix (see Appendix 1) was produced taking into consideration the characters proposed by Neusser et al. (2001) and modified for howlers by de Oliveira et al. (2002). These authors used an abbreviated nomenclature for ancestral Platyrrhini chromosome forms with a correspondence in the human karyotype. In the present contribution, for the character nomenclature, we refer directly to the human G band ideogram (Table 2). New characters were obtained from our karyological comparisons and introduced in the chromosomal dataset.

Molecular dataset: The sequences available in GenBank for the same species used in the G-banding pattern and FISH comparisons were taken into to choose the molecular marker. The only molecular marker that fulfilled all the requirements was cyt b. The sequences used were (Genbank Accession Numbers): *A. belzebul* (AY374348.2), *A. caraya* (AY374359.2), *A. s. arctoidea* (AY065886.1), *A. sara* (AY065887.1), *A. macconnelli* (AY065888.1), *A. g. guariba* (AY065899.1), *A. g. clamitans* (DQ679782.1), *A. pigra* (AY065884.1), *A. palliata* (AY065879.1) (Cortés-Ortiz et al. 2003, Harris et al. 2005, Lorenz et al. 2005, Nascimento et al. 2005, Casado et al. 2010). *Cebus apella* Linnaeus, 1758 (FJ529102.1) and *Lagothrix lagotricha* Humboldt, 1812 (AY671799.1) were used as outgroups. *C. apella*, from the Cebidae family, was taken as an outgroup species, since it is accepted that this species presents the most ancestral karyotype within Platyrrhini (Clemente et al. 1990, García et al. 2000). *Lagothrix lagotricha*, also a member of the Atelidae family, was chosen as the second outgroup to test the monophyly of the group. All sequences were aligned using CLUSTALW (Thompson et al. 1994).

Table 2. Human chromosome syntenic association considered as characters and used to construct the binary matrix of chromosomal homeologies among howler monkeys (modified from Neusser et al. 2001, de Oliveira et al. 2002).

1. 1p21-pter/1p12-21	36. 15q11 q13; q25 qter/Y	69. (10q/16p) ₂
2. 5q31.3-qter/ 7p22; q11 q21	37. 2pter q12	70. 11/(10q/16p) ₂
3. 5pter-q31.2/5q31.3-qter	38. 16q	71. 10q/16p
4. 2pter q12/16q	39. 3p24 q21; q13 q26/ 15q11 q13; q25 qter	72. 10q/16p/4pter-q22
5. 4q31.3-qter/4q23-q31.2	40. 11/5pter-q31.2	73. 10p/10q/16p/4pter-q22
6. 4q23-q31.2/4pter-q22	41. 5pter-q31.2/7p22; q11; q21	74. 10p/10q/16p
7. (10q/16p) ₃	42. 12/9	75. 19/13
8. 6	43. 1p21-pter/2pter q12	76. 19/22
9. 8p/18	44. 16q/4pter-q22	77. 22/1p21-pter
10. 15q21.3-q24/15q13-q21.2	45. 22/14	78. 19/22/1p21-pter
11. 15q11 q13; q25 qter	46. 2q13 qter/20	79. 2q13 qter/4q23-q31.2
12. 7p21 p11; q11 q21; q22 qter	47. 2q13 qter/4q23-q31.2	80. 6/1p12-21
13. 8q	48. 8q/2q13 qter	81. Y/15q11 q13; q25 qter/ 3p24 q21; q13 q26
14. 12	49. 7p22; q11; q21/8q	82. 1p12-21/5pter-q31.2/7p22; q11; q21/5q31.3-qter/ 7p22; q11 q21
15. 11	50. 7p22; q11; q21/8q	83. 9/22
16. 13	51. 17/2pter q12/12	84. 17/11
17. 9	52. 2pter q12/12	85. 3pter p24; p21 p12; q12 q13; q27 qter/8p
18. 3pter p24; p21 p12; q12 q13; q27 qter	53. 1q32 qter/11	86. 15q21.3-q24/1q32 qter
19. 3p24 q21; q13 q26	54. 1q32 qter/(11/5pter-q31.2) ₂	87. 16q/15q21.3-q24/1q32 qter
20. 1q32 qter	55. (11/5pter-q31.2) ₂	88. 4pter-q22/1p12-21
21. 1q21 q31	56. 18/14	89. 14
22. 3p12/21	57. 3pter p24; p21 p12; q12 q13; q27 qter/15q21.3-q24	90. 2pter q12/4pter-q22
23. 10p	58. 3pter p24; p21 p12; q12 q13; q27 qter/15q21.3-q24/16q	91. 15q13-q21.2/7p22; q11; q21/5q31.3-qter/ 7p22; q11; q21
24. 22	59. 15q21.3-q24/16q	92. 6/15q21.3-q24
25. X	60. 17/10p	93. 14/1p12-21
26. Y	61. 17/10p/19	94. 6/15q21.3-q24/14/1p12-21
27. 5pter-q31.2	62. 10p/19	95. 17/8p/18
28. (5q31.3-qter/7p22; q11 q21) ₂	63. 22/20	96. 22/10q/16p
29. 1p21-pter	64. 22/20/1q21 q31	97. 2q13 qter/11
30. 1p12-21	65. 20/1q21 q31	98. (10q/16p) ₂ /1q21 q31/20
31. 4q31.3-qter	66. (11/5pter-q31.2) ₃	99. Y/7
32. 4q23-q31.2/15q13-q212	67. 1p12-21/8p	
33. 4pter-q22	68. 7p21 p11; q11 q21; q22 qter /14/15q21.3-q24	
34. 14/15q21.3-q24		
35. (10q/16p) ₂ /(10q/16p) ₁		

/: separates the chromosomal segments that constitute an association. ()_n: n = number of repeats in the segment

Phylogeny: A Maximum Parsimony phylogeny using the exhaustive search option was obtained with PAUP 4.0 software (Phylogenetic Analysis Using Maximum Parsimony, (Swofford 2002)), for each separate partition and the combination of both the chromosomal and molecular datasets. All characters had the same weight, based on the premise that chromosome rearrangements occur by equal chance (de Oliveira et al. 2002, Dobigny et al. 2004). The relative stability of nodes was assessed by bootstrap estimates (Felsenstein 1985) based on 200 iterations. Each bootstrap replicate involved a heuristic parsimony search with 10 random taxon additions and tree-bisection reconnection (TBR) branch swapping.

Results

Classical cytogenetic analysis

Karyological analysis: The cytogenetic characterization of the *Alouatta* specimens showed diploid numbers, sex chromosome systems and G-banding patterns in agreement with the ones previously described for each species. Figures 1a and 1b show all the comparisons performed.

Chromosomal homeologies between A. caraya and A. palliata: The chromosomal rearrangements that could explain the homeologies were grouped in two categories: 1) *A. palliata* chromosomes with no rearrangements with respect to *A. caraya* chromosomes: 3, 5, 6, 7, 8, 9, 10, 13, 14, 16, 17, 19, 20, 21, 22, 24, 25 and X₁; 2) *A. palliata* chromosomes with more than one rearrangement with respect to *A. caraya* chromosomes: 1, 2, 4, 11, 12, 15, 18 and 23. No homeologies were allocated for *A. palliata* chromosome 26 and chromosome arms 4q and 2p using the level of resolution of the classical cytogenetic techniques applied. The rearrangements detected between the *A. caraya* and *A. palliata* karyotypes included at least seven fissions/fusions, two paracentric inversions and one deletion. *A. caraya* chromosome 7 (X₂ in males) shares homeology with two *A. palliata* chromosome pairs, 23 and 18, which are not the ones involved in the sex chromosome system in *A. palliata*. The *A. palliata* chromosomal pair 19 (X₂ in males) shares homeology with chromosome 14 of *A. caraya*.

Chromosomal homeologies between A. caraya and A. pigra: The chromosomal rearrangements that could explain the homeologies were grouped in two categories: 1) *A. pigra* chromosomes with no rearrangements with respect to *A. caraya* chromosomes: 2, 5, 6, 7, 8, 10, 15, 16, 17, 20, 22, 23, 25, 28 and X₁; 2) *A. pigra* chromosomes with more than one rearrangement with respect to *A. caraya* chromosomes: 1, 3, 4, 9, 11, 13, 18, 19, 24, 26 and 27. No homeologies were allocated for *A. pigra* chromosomes 4p_{prox}, 12, 14 and 21 using the level of resolution of the classical cytogenetic techniques applied. The rearrangements detected between the *A. caraya* and *A. pigra* karyotypes included at least 12 fissions/fusions, two paracentric inversions, two translocations and one deletion. *A. caraya* chromosome 7 (X₂ in males) shares homeology with two *A. pigra* chromosome pairs, 26 and 19, which are not the ones involved in the sex chromosome system in *A. pigra*. *A. pigra* chromosome 17 (X₂ in males) shares homeology with chromosome 14 of *A. caraya* (which in turn has homeology with HSA7).

Chromosomal homeologies among all howlers: The chromosomal homeologies found among all howlers are shown in Table 3 and Figures 1a and 1b. Results show that Mesoamerican howlers share several human chromosomal syntenic associations with South American ones: HSA15q13-q21.2/4q23-q31.2 and HSA16p/10q, shared with all howlers; HSA15q21.3-q24/14, shared with all howlers except *A. s. arctoidea* and *A. macconnelli*, and HSA8p/18, shared with all howlers except *A. s. arctoidea*. Two new chromosomal syntenic associations, HSA4p_{ter}-q22/9/11 and HSA15q21.3-q24/14/21q, were found for *A. pigra* in chromosomes 1 and 4q, respectively.




Figure 1. Comparison of *Homo sapiens* (HSA), *Alouatta caraya* (ACA), *A. pigra* (API), *A. palliata* (APA) and *A. guariba clamitans* (AGUc) G-banded chromosomes, taking *A. caraya*'s karyotype as reference. On the left, human chromosomal bands with homeology for its corresponding ACA chromosome segment are indicated. The boxes highlight the homeologies of the autosomes involved in the sex chromosome systems in these species **a** Comparison for ACA chromosomes #1 to #13 **b** Comparison for ACA chromosomes #14 to X₁

Table 3. Chromosomal homeologies between howlers, obtained from data both from this contribution and from previous reports. ACA: *A. caraya*; API: *A. pigra*; APA: *A. palliata*; AGU: *A. guariba*; ASEa: *A. s. arctoidea*; AMA: *A. macconnelli*; ASA: *A. sara*; ABE: *A. belzebul*.

Human Chromosomal associations†	ACA	API	APA	AGU	ASEa	AMA	ASA	ABE
1p12-p21	1	3p _{prox}	4p	2q _{ter}	9q _{ter}	18q _{ter}	16q _{ter}	23
5q _{prox} -q31.2				13	3q _{ter}	13	1q _{ter}	1q
7p22; q11; q21		13q		1q _{ter}	12q _{prox}	12q _{ter}	13q _{ter}	
5q31.3-q _{ter}					8q _{ter}		7q _{ter}	
7p22; q11; q21		24q			1q _{ter}		4q _{ter}	
6	2	2	3	2p _{ter} -q _{prox}	4	18p _{ter} -q _{prox}	5	4
						8		
22q	3	9	11	9p	9q _{prox}	5p _{prox}	8q _{prox}	6p _{ter}
9q		1q _{prox}	15	3p	13	15	11	2q
16q	4	1p	5	12q	6q _{ter}	3q	9p _{ter}	5
4p _{ter} -q22				1q _{prox}	11		14	
16p	5	6	7	9q	10	3p _{prox}	21	7
10q				20		2p	19	
16p								
10q								
8p	6	5	6	4p _{ter} -q _{prox}	15q _{ter}	6	2q _{ter}	8
18					5q _{prox}			
15q11-q13	7 (X ₂)	26q	23	7 (X ₂)	X ₂	X ₂	X ₂	24
15q25-q _{ter}								
3p24-p21		19q	18					17 (X ₂)
3q13-q26								
17	8	27q	8	4q _{ter}	7q _{prox}	7	1p-1q _{prox}	9
					2p			
4q31.3-q _{ter}	9	8	10	11	18	10	16p-q _{prox}	11
19	10	7	9	12	7q _{ter}	5p _{ter}	15p-q _{prox}	10
						4p		
2q13-q _{ter}	11	11	12	5p	2p _{prox}	11q _{prox}	3q _{prox}	12
20				17	9q _{int}	16q _{prox}	8q _{int}	
12	12	10	13	3q	2	14	6	2p
11	13	1q _{ter}	1q _{prox}	5q	3q _{prox}	2q	1q _{int}	1p
					12q _{ter}		13q _{prox}	
7 q11-q21	14	17 (X ₂)	19 (X ₂)	6	8	1q	7	13
7q22-q _{ter}								
3p _{ter} -p24	15	16	16	16	6q _{prox}	19	2q _{prox}	15
3p21-p12								
3q13								
3q27-q _{ter}								
8q	16	3q	--	19	1q _{prox}	12q _{prox}	4q _{prox}	16
13	17	18	2q	14	16	4q	12	14
2p _{ter} -q12	18	15	14	1p	2q _{prox}	17	10	3q
1p21-p _{ter}	19	20	22	15	14	5q	17	3p

Human Chromosomal associations†	ACA	API	APA	AGU	ASEa	AMA	ASA	ABE
15q21.3-q24	20	4q _{int}	17	2q _{int}	6q _{int}	1p	9q _{prox}	6p _{prox} -q _{ter}
14					5q _{ter}		18	
3p12	21	4p _{ter}	1p _{ter}	18	17	9	20	18q _{ter}
21q								
4q23-q31.2	22	22	21	1q _{int}	1p _{ter}	1q _{ter}	3q _{ter}	20
15q13-q21.2								
1q32-q _{ter}	23	23	20	X ₃	15q _{prox}	21	8q _{ter}	19
1q21-q31	24	25	25	8	3p	16q _{ter}	9q _{ter}	21
10p	25	28	24	22	7q _{int}	3p _{ter}	15q _{ter}	22
X	X ₁	X ₁	X ₁	X ₁	X ₁	X ₁	X ₁	X ₁

†from pter to qter.

Cytomolecular study

In the *Homo sapiens* metaphases, the hybridization signals on chromosomes HSA3, HSA21, HSA15, HSAX and HSAY for chromosome painting probes #3 (red), #21 (green), #15 (green) (Figure 2a), X and Y (data not shown) were corroborated.

In *A. g. clamitans*, the signal for HSA21 was observed in 18q_{ter}, the signal for HSA3 was observed in 18q_{prox} (thus corroborating the HSA3/21 synteny in *A. g. clamitans*), 16q, 7q (X₂ in males) and Y₂, and the signal for HSA15 was observed in 1_{int}, 2_{int}, 7p (X₂) and Y₁. This corroborates the HSA3/15 syntenic association to the multiple sex chromosome system X₁X₁X₂X₂X₃X₃/X₁X₂X₃Y₁Y₂ of this species (Figures 2b and 3b).

In *A. pigra*, the signal for HSA3 was observed in 16q and 19q, while that for HSA21 hybridized in 4p_{ter}, thus indicating that the HSA3/21 synteny is not present in *A. pigra* (Figures 2c and 3c). The probe for HSA15 hybridized in *A. pigra* metaphases in 4q_{ter}, 22q_{ter} and 26q, showing that the HSA3/15 syntenic association is also absent. None of these *A. pigra* chromosomes is involved in the sex chromosome system of this species (Figures 2d and 3c).

In *A. caraya*, the signal for HSA21 was observed in 21q_{ter}, whereas that for HSA3 was observed in 21q_{prox}, thus confirming the conservation of the HSA3/21 synteny. HSA15 hybridized in 7p (X₂ in males) and Y_{1ter}, and HSA3 in 7q and Y_{1prox}, exhibiting the HSA3/15 syntenic association in the sex chromosome system X₁X₁X₂X₂/X₁X₂Y₁Y₂ (Figure 3a).

A. palliata showed a pattern similar to that of *A. pigra* (therefore Figure 2 illustrates only the latter). HSA3 hybridized in 16q and 18q, HSA21 hybridized in 1p_{ter} and HSA15 in 2q_{ter}, 21q_{ter} and 23q (Figure 3d). Both the HSA3/21 and HSA3/15 syntenic associations are absent in *A. palliata* and chromosomes with homeology to HSA3 and HSA15 are also not involved in the sex chromosome system of this species.

The probe for the human X chromosome showed positive hybridization signal in X₁ of all the species analyzed. The probe for the human Y chromosome did not hybridize in any of the howler species (data not shown).




Figure 2. Analysis of the conservation of the HSA3/21 and HSA3/15 syntenic chromosomal associations in howlers (bar=10 μ m). The arrows indicate the chromosomes with positive FISH signal **a** *Homo sapiens* partial metaphase hybridized with probes HSA21 (green) and HSA3 (red) (control of the hybridization). Inset: *Homo sapiens* partial metaphase hybridized with HSA15 (green) **b** AGUc metaphase hybridized with HSA15 (green) and HSA3 (red) **c** API metaphase hybridized with HSA3 (red) and HSA21 (green) **d** API metaphase hybridized with HSA15 (green).

Phylogenetic analysis

The data obtained from the G-banding pattern and FISH homeologies, together with cyt b sequences obtained from previous reports, were used as the basis to perform a cladistic analysis. The HSAY/7 association, corresponding to the Y-autosome translocation that gave rise to the multivalents observed in *A. pigra* and *A. palliata*, was added




Figure 3. Howler monkeys G-banded chromosomes with positive signal for the human chromosome painting probes analyzed. On the right, the hybridization pattern of human chromosomes #3, #21 and #15. **a** *A. caraya* **b** *A. guariba clamitans* **c** *A. pigra* **d** *A. palliata*.

as an extra character to the original list (de Oliveira et al. 2002). The syntenic associations HSA4p_{ter}-q22/9/11 observed in chromosome 1 of *A. pigra* and HSA15q21.3-q24/14/21q observed in chromosome arm 4q were not included in the analysis, because, as autopolymorphies for *A. pigra*, they are considered non-informative.

Three data matrices were obtained: one including only chromosomal data, another including only molecular data and the last one including both types of characters (chromosomal and molecular) in a single matrix (see Appendix 1).

Chromosomal partition: The analysis of chromosomal data resulted in 36 informative characters, 23 constant characters and 40 non-informative characters. After ana-




Figure 4. 50% majority consensus tree obtained by “bootstrap” for the combined analysis. Next to the name of each species, the diploid number (2N) and sex chromosome system is described. (m)=microchromosomes.

lyzing 704 trees, PAUP retained the two most parsimonious trees (Appendix 2: Figures Sa and Sb), both with a length of 87 ($L = 87$). The analysis using only the partition of chromosomal data did not resolve the node (*(A. palliata, A. pigra)*, *(A. caraya, A. belzebul)*, *(A. g. clamitans, A. g. guariba)*, *(A. macconnelli (A. sara, A. s. arctoidea))*), since it was established in a polytomy (Appendix 2: Figure Sc).

Molecular partition: Heuristic analysis of cyt b gene sequences, made from a total of 800 characters, produced 109 informative characters, 551 constant characters and 140 non-informative characters. After analyzing 916 trees, PAUP retained a single most parsimonious tree (Appendix 2: Figure Sd), with a length of $L = 366$. The analysis using only molecular data did not resolve the node *(A. sara, A. macconnelli, A. s. arctoidea, A. caraya)*, which was established as a polytomy different from that described from chromosomal data.

Combined analysis: The heuristic analysis of the combined data showed a total of 899 characters, 145 of which were informative, 180 non-informative and 574 constant. After analyzing 684 trees, PAUP retained only one, with a length of $L=460$ (Figure 4). This type of analysis allowed us to solve all the nodes, resulting in a fully resolved tree.

Discussion

We present the first phylogenetic study using a combined analysis of chromosomal and molecular characters in Ceboidea to contribute to the characterization of the specio-genic processes in howler monkeys. The homoplasy distribution is likely to be different

in each dataset because these are subject to different constraints. Therefore, when different datasets are analyzed simultaneously, the signal common to all of them is more likely to overwhelm the homoplasy signal on the data (Kluge 1989).

In primates, few studies have compared and taken into account more than one type of character. Bonvicino et al. (2001) superimposed chromosomal information on the phylogeny obtained from molecular characters. Villalobos et al. (2004) used numerical and metric values that describe the karyotype, such as diploid number (2N) and fundamental number (FN), in a combined phylogenetic analysis with morphological characters. However, these values (2N, FN, etc) can be identical simply by chance and, if interpreted in a phylogenetic context, may be spurious indicators of relatedness (Dobigny et al. 2004). Our encoding strategy (using the rearrangements as characters) is quite similar to that used for morphological data but in cytogenetics one can retrieve information on the mutational event itself, something that is clearly not available to morphologists. As such, chromosomal mutations that accumulate along the tree are comparable to transitions, transversions, and insertions/deletions in molecular phylogenies (Dobigny et al. 2004). Our combined phylogeny evidences the accuracy of this encoding strategy.

In all the above-mentioned contributions, the $X_1X_1X_2X_2/X_1X_2Y_1Y_2$ sex chromosome system was proposed as the ancestral condition for the genus. However, as discussed by Solari and Rahn (2005), the $X_1X_1X_2X_2/X_1X_2Y$ sex chromosome system is simpler and is present in other genera of Neotropical Primates, such as *Aotus* Illiger, 1811, *Callimico* Miranda Ribeiro, 1912, and *Cacajao* Lesson, 1840 (Ma et al. 1976, Seuánez et al. 1989, Moura-Pensin et al. 2001). The $X_1X_1X_2X_2/X_1X_2Y$ sex chromosome system as an ancestral state appears to be a more parsimonious hypothesis. Moreover, since Mesoamerican howlers (*A. pigra* and *A. palliata*) were poorly karyologically characterized at the time, data on these howlers are missing in all previous contributions.

Homeology analysis

The karyotypes of *A. pigra* and *A. palliata* share more syntenic associations with those of *A. caraya* and *A. belzebul* than with those of the “*A. seniculus* group” (*A. s. arctoidea*, *A. sara*, *A. macconnelli*, denominated as such because they were once all subspecies of *A. seniculus* together with *A. s. seniculus* Linnaeus, 1766, and *A. s. stramineus* Hill, 1962). This supports the basal grouping of the *A. pigra*-*A. palliata* Mesoamerican clade and the basal grouping of *A. belzebul* among South American howlers.

The chromosomal comparisons showed that *A. pigra* and *A. palliata* conserved the HSA8/18 and HSA14/15 syntenies, considered ancestral for Platyrrhini (Stanyon et al. 2008), as well as the HSA10/16/10/16 syntenic association, ancestral for Atelidae (de Oliveira et al. 2002), but lost the HSA3/21 syteny, ancestral for mammals (Müller et al. 2000, Müller 2006).

According to our combined phylogeny, the HSA2/20 and HSA5/7/5/7 syntenic associations, previously considered as synapomorphies of the *A. caraya*-*A. belzebul* group (de Oliveira et al. 2002), would be homoplasies (parallelism). The HSA16/4

syntenic association would be ancestral for the genus and might either be absent in *A. pigra*, *A. guariba* and the *A. seniculus* group or might constitute a parallelism among *A. palliata*, *A. caraya* and *A. belzebul*.

Like the HSA3/21 synteny, the HSA3/15 syntenic association, involved in the sex chromosome systems in South American howlers, is not present in Mesoamerican ones. This syntenic association of human 3/15 chromosomal segments has been described in other Atelidae species such as *Ateles geoffroyi* Kuhl, 1820 and *Ateles belzebul hybridus* Geoffroy, 1829, although not associated with the sex chromosome system (Morescalchi et al. 1997), but not observed in *Lagothrix* Geoffroy, 1812, and *Brachyteles* Spix, 1823 (Stanyon et al. 2001, de Oliveira et al. 2005). This association has not been observed in other genera of Neotropical Primates such as *Cebus libidinosus* Spix, 1823, or *Saimiri boliviensis boliviensis* Geoffroy & Blainville, 1834 (Mudry et al. 2001). Therefore, the HSA3/15 syntenic association either could be interpreted as the ancestral condition for the family Atelidae, where the association with multiple sex chromosomes would be an evolutionary novelty (apomorphy) in howlers and the loss of the association a apomorphy for the *Lagothrix* and *Brachyteles* group, or could have appeared independently in *Alouatta*, involved in the Y-autosome translocation, and in *Ateles*, not involved in the sex chromosome system (de Oliveira et al. 2005). However, our results suggest that the HSA3/15 syntenic association is not an ancestral condition for *Alouatta*, since the most basal species (see Figure 4) *A. pigra* and *A. palliata* (this contribution) and *A. belzebul* (Consigliere et al. 1996) do not possess this association.

Possible origin of the multivalents

Taking into consideration the data obtained, a hypothesis can be proposed regarding the origin of the sex chromosome systems in the genus. Within the family Atelidae, with the exception of *Alouatta*, all genera have an XX/XY sex chromosome system. Therefore, it can be considered that the *Alouatta* ancestor possessed a chromosomal sex determination XX/XY, prior to the biogeographic separation of Mesoamerican and South American groups (see below). After this separation, both groups independently acquired the multiple sex chromosome systems currently observed through independent Y-autosome translocations.

The sex chromosome system $X_1X_1X_2X_2/X_1X_2Y$ would have arisen independently in the lineages of Meso and South American howlers by a Y-autosome translocation (Figure 5a). In males, two fissions, one in $Y_{p_{ter}}$ and another in $q_{p_{prox}}$ of the autosomal pair involved ($Aq_{p_{prox}}$), followed by translocation of $Aq_{p_{prox}}$ to $Yq_{p_{prox}}$, formed the new chromosome Y_1 . The $Y_{p_{ter}}$ segment is lost and the proximal region of the fissioned autosome either is lost or, in certain howler species, could have given rise to microchromosomes (e.g.: *A. seniculus* (Yunis et al. 1976, Lima and Seuánez 1991, Torres and Leibovici 2001), *A. sara* (Minezawa et al. 1985) and *A. macconnelli* (Lima et al. 1990)). The homologous autosomal pair involved in the translocation is the one now denominated X_2 . In the case of South American howlers, the autosomal pair involved




Figure 5. a Possible origin for X₁X₁X₂X₂/X₁X₂Y sex chromosome system in the genus *Alouatta*. The ancestral X chromosome is shown in white, the ancestral Y chromosome in light gray and the autosomal pair (A) in dark gray. Two fissions occurs, one in Yp_{ter} and another in q_{prox} of the autosomal pair involved (Aq_{prox}). The translocation of Yq-p_{prox} to the Aq formed the new Y₁ chromosome and the homolog of the autosomal pair involved in the translocation is now denominated X₂. The Yp_{ter} acentric fragment is lost and the rest of the autosome (Ap and Aq_{prox}) could either be lost or remain as a microchromosome in some howlers **b** Possible origin for the X₁X₁X₂X₂/X₁X₂Y₁Y₂ sex chromosome systems from a X₁X₁X₂X₂/X₁X₂Y system. The ancestral X is shown in white, the ancestral Y in light gray, the autosomal pair involved in the first translocation (A) in dark gray and the autosomal pair (A') involved in the formation of this new sex chromosome system in black. Simultaneous breaks in X₂p_{prox} and A'q_{prox} followed by the translocation of the rest of the A'q to X₂p_{prox} give origin to the new X₂ chromosome. The X₂p_{ter} acentric fragment could be lost and the rest of the autosome (A') could either be lost or remain as a microchromosome in some howlers. The homolog to the autosomal chromosome in question is now identified as Y₂ **c** Simultaneous breaks in Y₁q and Y₂q and a translocation between Y₁ and Y₂ further explain the hybridization pattern observed in the sex chromosome systems of South American howlers. A *de novo* centromere arises in the remains of the old Y₁ (now Y₂). The remains of the old Y₂ could either be lost or remain as a microchromosome in some howlers **d** Hybridization pattern in South American howlers.

would share homeology with HSA3, whereas in the Mesoamerican species it would share homeology with HSA7.

From this $X_1X_1X_2X_2/X_1X_2Y$ sex chromosome system, an $X_1X_1X_2X_2/X_1X_2Y_1Y_2$ system could have arisen from a new translocation (Figure 5b). Under this hypothesis, simultaneous breaks in $X_{2p_{prox}}$ and q_{prox} of another autosome ($A'q_{prox}$), followed by the translocation of most of the $A'q$ arm to $X_{2p_{prox}}$, gave rise to the new X_2 chromosome. The $X_{2p_{prox}}$ acentric fragment is lost and the rest of the autosome (A') either is lost or could have remained as a microchromosome in some howler species (see above). The chromosome homologous to the autosome in question (A') became Y_2 . In the case of South American howlers, the new autosomal pair involved in the sex chromosome system would share homeology with HSA15. A further translocation between Y_1 and Y_2 (Figure 5c) would explain the hybridization pattern of the segments with homeology to human chromosomes 3 and 15 observed in the sex chromosome systems $X_1X_1X_2X_2/X_1X_2Y_1Y_2$ in South American howlers (Figure 5d).

On the other hand, in the Mesoamerican species, the $X_1X_1X_2X_2/X_1X_2Y_1Y_2$ sex chromosome system could have arisen either as described in Figure 5b (with the autosomal pair involved sharing homeology with a human chromosome not yet identified by G-banding pattern) or by a fission in Y_1 that would have given rise to two chromosomes, the new Y_1 (containing the segment corresponding to the ancestral Y chromosome) and Y_2 (containing a portion of the autosomal pair with homeology to HSA7). This last hypothesis would require a centromeric activation in Y_2 .

However, considering the observation of the independent origin of the multiple sex chromosome systems in these two groups of howlers, the possibility of an independent origin of the $X_1X_1X_2X_2/X_1X_2Y$ and $X_1X_1X_2X_2/X_1X_2Y_1Y_2$ sex chromosome systems within the Meso and South American groups cannot be ruled out until further studies.

It can be considered that multiple sex chromosome systems would be an extremely rare phenomenon due to complication in meiosis. Extreme cases are platypus and echidna, with a large number of sex chromosomes (Bick and Jackson 1967, Renz et al. 2007). In primates, multiple sex chromosome systems are even more infrequent. Moreover, *Alouatta* would be the first case where an independent origin of multiple sex chromosome systems is described. In other taxa, such as *Drosophila* (Flores et al. 2008), Erythrinidae fishes (Cioffi et al. 2013) and mole-rats (Deuve et al. 2006), a few cases have been observed, but these descriptions are still scarce.

Phylogeny of *Alouatta*

The chromosomal homeologies and FISH analysis were used to construct a data matrix for the phylogenetic analysis. For comparison purposes, independent phylogenetic reconstructions were performed with each type of partition (Appendix 2: Figure Sa, b, c and d), along with the combined analysis of the two datasets (Figure 4). The chromosome partition grouped *A. caraya* and *A. belzebul* as sister taxa, in agreement with that reported by de Oliveira et al. (2002), a relationship that was not observed in the other

two analyses, which grouped *A. caraya* with *A. sara*, *A. s. arctoidea* and *A. macconnelli* (although in the case of the molecular partition this relationship constituted a polytomy). This last species arrangement was also proposed by Nascimento et al. (2005) and Cortés-Ortiz et al. (2003) using molecular characters. In our molecular data partition (Figure Sd), *A. belzebul* was grouped with the clade of *A. guariba*, in agreement with that reported by Bonvicino et al. (2001) and Cortés-Ortiz et al. (2003). The three types of analyses agreed to place *A. sara*, *A. s. arctoidea* and *A. macconnelli* into a single group, although the molecular data partition did not resolve the relationships between them, as they formed a polytomy. The grouping of all those species (“*A. seniculus* group”) was observed in all phylogenetic studies performed so far (see above). Another coincidence was that the Mesoamerican species were placed as a separate clade from other South American species and, as expected, the two subspecies of *A. guariba* in one group. All sets of taxa analyzed in this new approach were solved without polytomies only with the combined analysis, demonstrating the usefulness of incorporating more than one source of data for a more accurate elucidation of the relationships among current taxa.

The grouping of South American species as a separate group of the Mesoamerican group coincides with previous phylogenetic analyses using only molecular characters (Cortés-Ortiz et al. 2003, Ellsworth and Hoelzer 2006) and with the hypothesis of monophyletic origin of the Mesoamerican howlers previously proposed by Smith (1970). Smith’s hypothesis holds that Mesoamerican howlers originated by an expansion of the geographic distribution of South American howlers after the formation of the Isthmus of Panama, estimated to be completed about 3 million years ago (Coates et al. 2004). However, other studies indicate that the rise of the isthmus was a process rather than an event (Knowlton and Weigt 1998), resulting in intermittent periods with connected and divided lands during the past 18 million years. Given this last fact, another hypothesis was postulated to explain the current geographic distribution of the species of the genus. Instead of a single colonization event, various founder events, either across the Isthmus of Panama during one of the periods in which the two Americas were connected or across islands in the Caribbean archipelago, could have occurred (Cortés-Ortiz et al. 2003, Ellsworth and Hoelzer 2006). Primate fossils have been found in Cuba and Jamaica, but the origin and relationships of these specimens with modern Platyrrhini are still under debate (Fleagle 1999, Gutiérrez Calvache and Jaimez Salgado 2007, Rosenberger et al. 2009, Cooke et al. 2011).

Independently of the biogeographic scenario under consideration, it is clear that the evolutionary history of Mesoamerican howlers is different from that of South American howlers, an assertion that would be supported by the evidence provided by our new data.

This contribution provides new useful information for the systematics of the genus *Alouatta*, while supporting the hypothesis of chromosomal evolution in primates as a specio-genic strategy. The combined analysis resolved the phylogenetic relationships between howler species of both American origins, as a first approach to the “Total Evidence” concept and towards clarifying the controversies related to the Taxonomy and Evolution of Ceboidea.

Acknowledgements

All research reported in this manuscript has met the appropriate national and institutional guidelines for the legal acquisition and use of laboratory animals and authorized study of wild animals. The authors also adhered to the Guide for Care and Use of Experimental Animals, as promulgated by the Canadian Council of Animal Care, and to the American Society of Primatologists (ASP) Principles for the Ethical Treatment of non-human Primates. The work which took place both in Argentina as in Mexico was done in accordance with the laws of these countries.

We would like to thank the head of the institutions that made possible the sampling of the specimens for the genetic studies presented here, the veterinarians who collaborated in the sampling of all specimens, and the caretakers of the animals. To Dr. Liliana Cortés-Ortiz (Dept. of Ecology and Evolutionary Biology & Museum of Zoology, University of Michigan), Dr. Domingo Canales-Espinosa; Vet. Javier Hermida, Sr. Francisco García-Orduña (Instituto de Neuroetología, Universidad Veracruzana, Xalapa, Veracruz, México) and all the field team for their collaboration and support in the obtention of the samples of mexican howlers in their natural geographic distribution. MDM PIP-CONICET 0744 and UBACyT X154 provided funding for this project.

References

- Amaral P, Finotelo L, de Oliveira E, Pissinatti A, Nagamachi C, Pieczarka J (2008) Phylogenetic studies of the genus *Cebus* (Cebidae-Primates) using chromosome painting and G-banding. *BMC Evolutionary Biology* 8: 169. doi: 10.1186/1471-2148-8-169
- Armada JLA, Barroso CML, Lima MMC, Muniz JAPC, Seuánez HN (1987) Chromosome studies in *Alouatta belzebul*. *American Journal of Primatology* 13: 283–296. doi: 10.1002/ajp.1350130306
- Bick YAE, Jackson WD (1967) Karyotype of the Monotremes *Ornithorhynchus anatinus* (Platypus) and *Tachyglossus aculeatus* (Echidna). *Nature* 214: 600–601. doi: 10.1038/214600a0
- Bonvicino CR, Lemos B, Seuánez H (2001) Molecular Phylogenetics of howler monkeys (*Alouatta*, Platyrrhini) a comparison with karyotypic data. *Chromosoma* 110(3): 241–246. doi: 10.1007/s004120000128
- Cabrera A (1940) Los nombres científicos de algunos monos americanos. *Ciencia* 9: 402–405.
- Casado F, Bonvicino CR, Nagle C, Comas B, Manzur TD, Lahoz MM, Seuánez HN (2010) Mitochondrial divergence between 2 populations of the hooded capuchin, *Cebus (Sapajus) cay* (Platyrrhini, Primates). *Journal of Heredity* 101(3): 261–269. doi: 10.1093/jhered/esp119
- Cioffi MB, Liehr T, Trifonov V, Molina WF, Bertollo LAC (2013) Independent Sex Chromosome Evolution in Lower Vertebrates: A Molecular Cytogenetic Overview in the Erythrinidae Fish Family. *Cytogenetic and Genome Research* 141(2–3): 186–194. doi: 10.1159/000354039

- Clemente IC, Ponsà M, García M, Egozcue J (1990) Evolution of the simiiformes and the phylogeny of human chromosomes. *Human Genetics* 84(6): 493–506. doi: 10.1007/BF00210798
- Coates AG, Collins LS, Aubry M-P, Berggren WA (2004) The Geology of the Darien, Panama, and the late Miocene-Pliocene collision of the Panama arc with northwestern South America. *Bulletin of Geological Society of America* 116(11/12): 1327–1344. doi: 10.1130/B25275.1
- Consigliere S, Stanyon R, Koehler U, Agoramoorthy G, Wienberg J (1996) Chromosome painting defines genomic rearrangements between red howler monkey subspecies. *Chromosome Research* 4(4): 264–270. doi: 10.1007/BF02263675
- Consigliere S, Stanyon R, Koehler U, Arnold N, Wienberg J (1998) In situ hybridization (FISH) maps chromosomal homologies between *Alouatta belzebul* (Platyrrhini, Cebidae) and other primates and reveals extensive interchromosomal rearrangements between howler monkey genomes. *American Journal of Primatology* 46(2): 119–133. doi: 10.1002/(SICI)1098-2345(1998)46:2<119::AID-AJP2>3.0.CO;2-Z
- Cooke SB, Rosenberger AL, Turvey S (2011) An extinct monkey from Haiti and the origins of the Greater Antillean primates. *Proceedings of the National Academy of Sciences* 108(7): 2699–2704. doi: 10.1073/pnas.1009161108
- Cortés-Ortiz L, Bermingham E, Rico C, Rodríguez-Luna E, Sampaio I, Ruiz-García M (2003) Molecular systematics and biogeography of the Neotropical monkey genus, *Alouatta*. *Molecular Phylogenetics and Evolution* 26(1): 64–81. doi: 10.1016/S1055-7903(02)00308-1
- Crockett C, Eisenberg J (1987) Howlers: variations in group size and demography. In: Smuts BB, Cheney DL, Seyfarth RM, Wrangham RW, Struhsaker TT (Eds) *Primate societies*. Chicago IL, USA, 54–68.
- de Grouchy J, Turleau C, Roubin M, Klein M (1972) Karyotypic evolution in man and chimpanzees. A comparative study of band topographies after controlled denaturation. *Annales de Génétique* 15(2): 79–84.
- de Oliveira EHC, Lima MMC, Sbalqueiro IJ (1995) Chromosomal variation in *Alouatta fusca*. *Neotropical Primates* 3: 181–183.
- de Oliveira EHC, de Lima MMC, Sbalqueiro IJ, Pissinati A (1998) The karyotype of *Alouatta fusca clamitans* from Rio de Janeiro, Brazil: Evidence for a y-autosome translocation. *Genetics and Molecular Biology* 21(3): 361–364. doi: 10.1590/S1415-47571998000300012
- de Oliveira EHC, Neusser M, Figueiredo WB, Nagamachi C, Pieczarka JC, Sbalqueiro IJ, Wienberg J, Müller S (2002) The phylogeny of howler monkeys (*Alouatta*, Platyrrhini): Reconstruction by multicolor cross-species chromosome painting. *Chromosome Research* 10(8): 669–683. doi: 10.1023/A:1021520529952
- de Oliveira EHC, Neusser M, Müller S (2012) Chromosome Evolution in New World Monkeys (Platyrrhini). *Cytogenetics and Genome Research* 137(2–4): 259–272. doi: 10.1159/000339296
- de Oliveira EHC, Neusser M, Pieczarka JC, Nagamachi C, Sbalqueiro IJ, Müller S (2005) Phylogenetic inferences of Atelinae (Platyrrhini) based on multi-directional chromosome painting in *Brachyteles arachnoides*, *Ateles paniscus paniscus* and *Ateles b. marginatus*. *Cytogenetics and Genome Research* 108(1–3): 183–90. doi: 10.1159/000080814

- Deuve J, Bennett N, O'Brien P, Ferguson-Smith M, Faulkes C, Britton-Davidian J, Robinson T (2006) Complex evolution of X and Y autosomal translocations in the giant mole-rat, *Cryptomys mehowi* (Batherygidae). *Chromosome Research* 14(6): 681–691. doi: 10.1007/s10577-006-1080-3
- Dobigny G, Ducroz JF, Robinson TJ, Volobouev V (2004) Cytogenetics and Cladistics. *Systematic Biology* 53(3): 470–484. doi: 10.1080/10635150490445698
- Dutrillaux B, Couturier J (1981) The ancestral karyotype of platyrrhine monkeys. *Cytogenetics and Cell Genetics* 30(4): 232–242. doi: 10.1159/000131614
- Egozcue J, De Egozcue V (1966) The Chromosome Complement of the Howler Monkey (*Alouatta caraya*, HUMBOLDT 1812). *Cytogenetics and Genome Research* 5(1): 20–27. doi: 10.1159/000129881
- Elliot DG (1910) Description of a new species of monkeys of the Genera *Galago*, *Alouatta* and *Cercopithecus*. *Annals and Magazine of Natural History* 18(5): 77–83. doi: 10.1080/00222931008692726
- Ellsworth JA, Hoelzer GA (2006) Genetic Evidence on the Historical Biogeography of Central American Howler Monkeys. In: Lehman SM, Fleagle JG (Eds) *Primate Biogeography*. New York, USA, 81–103. doi: 10.1007/0-387-31710-4_3
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39(4): 783–791. <http://www.jstor.org/stable/10.2307/2408678>, doi: 10.2307/2408678
- Fleagle JG (1999) *Primate adaptation and evolution*. Academic Press, San Diego, California, USA, 486 pp.
- Flores SV, Evans AL, McAllister BF (2008) Independent Origins of New Sex-Linked Chromosomes in the melanica and robusta Species Groups of *Drosophila*. *BMC Evolutionary Biology* 8(1): 33. doi: 10.1186/1471-2148-8-33
- García F, Nogués C, Ponsà M, Ruiz-Herrera A, Egozcue J, García Caldés M (2000) Chromosomal homologies between humans and *Cebus apella* (Primates) revealed by ZOO-FISH. *Mammalian Genome* 11(5): 399–401. doi: 10.1007/s003350010075
- García F, Ruiz-Herrera A, Egozcue J, Ponsà M, García M (2002) Chromosomal homologies between *Cebus* and *Ateles* (Primates) based on ZOO-FISH and G-banding comparisons. *American Journal of Primatology* 57(4): 177–188. doi: 10.1002/ajp.10047
- Geoffroy Saint-Hilaire E (1812) Tableau des quadrumanes ou des animaux composant le premier ordre de la classe des mammifères. *Annales du Muséum national d'histoire naturelle*, Paris 7: 260–273.
- Geoffroy Saint-Hilaire E (1829) *Cours de l'histoire naturelle des Mammifères*. Pichon et Didier, Paris, Leçon IX, 36 pp.
- Geoffroy Saint-Hilaire I, de Blainville (1834) Rapport sur les résultats scientifiques du voyage de M. Alcide d'Orbigny dans l'Amérique du Sud, pendant des années 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833. Partie Zoologique. *Nouvelles annales du Muséum d'Histoire Naturelle* 3: 84–115.
- Gray JE (1849) Description of two species of Mammalia from Caracas. *Proceedings of the Zoological Society of London* 848: 138.

- Gregorin R (2006) Taxonomia e variação geográfica das espécies do gênero *Alouatta* Lacépède (Primates, Atelidae) no Brasil. *Revista Brasileira de Zoologia* 23(1): 64–144. doi: 10.1590/S0101-81752006000100005
- Groves CP (2005) Order Primates. In: Wilson DE, Reeder DM (Eds) *Mammal species of the world: a taxonomic and geographic reference*. Johns Hopkins University Press, Baltimore, Maryland, USA, 111–185.
- Groves CP (2001) *Primate Taxonomy*. Smithsonian Institution Press, Washington DC, USA, 350 pp.
- Gutiérrez Calvache DA, Jaimez Salgado EJ (2007) Introducción a los primates fósiles de las Antillas. 120 años de Paleoprimatología en el caribe insular. Editora Universitaria - UASD, Santo Domingo, Cuba, 208 pp.
- Harris EE, Gifalli-Iughetti C, Braga ZH, Koiffmann CP (2005) Cytochrome b Sequences Show Subdivision between Populations of the Brown Howler Monkey (*Alouatta guariba*) from Rio de Janeiro and Santa Catarina, Brazil. *Neotropical Primates* 13(2): 16–21. doi: 10.1896/1413-4705.13.2.16
- Humboldt A de (1812) *Recueil d'observations de zoologie et d'anatomie comparée, faites dans l'océan Atlantique, dans l'intérieur du nouveau continent et dans la mer du sud pendant les années 1799, 1800, 1801, 1802 et 1803*. Vol 1. Paris, France, VIII+368 pp.
- Illiger JCW (1811) *Prodomus systematis mammalium et avium additis terminis zoographicis utriusque classis, eorumque versione germanica*. Berlim, Berolini, Sumptibus, C. Salfeld, XVIII+301 pp.
- Kluge AG (1989) A Concern for Evidence and a Phylogenetic Hypothesis of Relationships Among *Epicrates* (Boidae, Serpentes). *Systematic Zoology* 38(1): 7–25. doi: 10.1093/sysbio/38.1.7
- Knowlton N, Weigt LA (1998) New dates and new rates for divergence across the Isthmus of Panama. *Proceedings of the Royal Society B* 265(1412): 2257–2263. doi: 10.1098/rspb.1998.0568
- Koiffmann CP, Saldanha PH (1974) Cytogenetics of Brazilian monkeys. *Journal of Human Evolution* 3(4): 275–282. doi: 10.1016/0047-2484(74)90021-9
- Kuhl H (1820) *Beiträge zur Zoologie und Vergleichenden Anatomie*. Hermannschen Buchhandlung, Frankfurt, 152 pp.
- Lacépède BGE de la V (1799) *Tableau des divisions, sous-divisions, ordres et genres des Mammifères*. Plassan, Paris, 18 pp.
- Lawrence B (1933) Howler monkeys of the *palliata* group. *Bulletin of the Museum of Comparative Zoology* 75(8): 313–354.
- Lesson RP (1840) *Species de Mammifères: Bimanes et quadrumanes; suivi d'un mémoire sur les Oryctérope*. JB Baillière, Paris, VIII+291 pp.
- Linnaeus C (1758) *Systema Naturae per Regna tria naturae, classes, ordines, genera, species cum bracteribus, differentiis, synonymis, locis*. Regnum Animale. Estocolmo, Holmiae, 823 pp.
- Linnaeus C (1766) *Systema Naturae per Regna tria naturae, classes, ordines, genera, species cum bracteribus, differentiis, synonymis, locis*. Regnum Animale. Estocolmo, Holmiae, vol 1, 823 pp.

- Lima MMC, Sampaio MIC, Schneider MPC, Scheffrahn W, Schneider H, Salzano FM (1990) Chromosome and protein variation in red howler monkeys. *Revista Brasileira de Genetica* 13(4): 789–802.
- Lima MMC, Seuánez H (1991) Chromosome studies in the red Howler Monkey, *Alouatta seniculus stramineus* (Platyrrhini, Primates): description of an $X_1X_2Y_1Y_2/X_1X_1X_2X_2$ sex chromosome system and karyological comparisons with other subspecies. *Cytogenetics and Cell Genetics* 57(2-3): 151–156. doi: 10.1159/000133135
- Lönnberg E (1941) Notes on members of the genera *Alouatta* and *Aotus*. *Arkiv für Zoologi* 33A (10): 1–44.
- Lorenz JG, Jackson WE, Beck JC, Hanner R (2005) The Problems and Promise of DNA Barcodes for Species Diagnosis of Primate Biomaterials. *Philosophical Transactions of the Royal Society B* 360(1462): 1869–1877. doi: 10.1098/rstb.2005.1718
- Ma NSF, Elliott MW, Morgan L, Miller A, Jones TC (1976) Translocation of Y chromosome to an autosome in the Bolivian owl monkey, *Aotus*. *American Journal of Physical Anthropology* 45(2): 191–201. doi: 10.1002/ajpa.1330450205
- Meireles CM, Czelusniak J, Ferrari SF, Schneider MPC, Goodman M (1999) Phylogenetic relationships among Brazilian howler monkeys, genus *Alouatta* (Platyrrhini, Atelidae), based on gamma1-globin pseudogene sequences. *Genetics and Molecular Biology* 22(3): 337–344. doi: 10.1590/S1415-47571999000300009
- Minezawa M, Harada M, Jordan OC, Borda CJV (1985) Cytogenetics of Bolivian endemic red howler monkeys (*Alouatta seniculus sara*): accessory chromosomes and Y-autosome translocation-related numerical variations. *Research Report on New World Monkeys* 5: 7–16.
- Miranda Ribeiro A de (1912) Zwei neue Affen unserer Fauna. *Brasilianische Rundschau* 2(1): 21–23.
- Morescalchi M, Schempp W, Consigliere S, Bigoni F, Wienberg J, Stanyon R (1997) Mapping chromosomal homology between humans and the black-handed spider monkey by fluorescence in situ hybridization. *Chromosome Research* 5(8): 527–536. doi: 10.1023/A:1018489602312
- Moura-Pensin C, Pieczarka J, Nagamachi C, Pereira Carneiro Muniz JA, Do Carmo de Oliveira Brigido M, Pissinatti A, Marinho A, de Souza Barros RM (2001) Cytogenetic relations among the genera of the subfamily Pitheciinae (Cebidae, Primates). *Caryologia* 54(4): 385–391. doi: 10.1080/00087114.2001.10589250
- Mudry MD (1990) Cytogenetic variability within and across populations of *Cebus apella* in Argentina. *Folia Primatologica* 54(3-4): 206–216. doi: 10.1159/000156445
- Mudry MD, Labal de Vinuesa ML, Colillas OJ, Brieux de Salum S (1984) Banding patterns of *Alouatta caraya*. *Revista Brasileira de Genetica* 7: 373–379.
- Mudry MD, Ponsà M, Borrell A, Egozcue J, García M (1994) Prometaphase chromosomes of the howler monkey (*Alouatta caraya*): G, C, NOR, and restriction enzyme (RES) banding. *American Journal of Primatology* 33(2): 121–132. doi: 10.1002/ajp.1350330206
- Mudry MD, Rahn MI, Gorostiaga M, Hick A, Merani MS, Solari AJ (1998) Revised Karyotype of *Alouatta caraya* (Primates: Platyrrhini) Based on Synaptonemal Complex and Banding Analyses. *Hereditas* 128(1): 9–16. doi: 10.1111/j.1601-5223.1998.00009.x

- Mudry MD, Rahn MI, Solari AJ (2001) Meiosis and chromosome painting of sex chromosome systems in Ceboidea. *American Journal of Primatology* 54(2): 65–78. doi: 10.1002/ajp.1013
- Müller S (2006) Primate Chromosome Evolution. In: Lupski JR, Stankiewicz P (Eds) *Genomic Disorders*, Humana Press Inc, New Jersey, USA, 133–152. doi: 10.1007/978-1-59745-039-3_9
- Müller S, Stanyon R, Finelli P, Archidiacono N, Wienberg J (2000) Molecular cytogenetic dissection of human chromosomes 3 and 21 evolution. *Proceedings of the National Academy of Sciences* 97(1): 206–211. doi: 10.1073/pnas.97.1.206
- Nascimento FF, Bonvicino CR, Da Silva FCD, Schneider MPC, Seuánez H (2005) Cytochrome b polymorphisms and population structure of two species of *Alouatta* (Primates). *Cytogenetics and Genome Research* 108(1-3): 106–111. doi: 10.1159/000080808
- Neusser M, Stanyon R, Bigoni F, Wienberg J, Müller S (2001) Molecular cytogenetics of New World monkeys (Platyrrhini) - comparative analysis of five species by multi-color chromosome painting gives evidence for a classification of *Callimico goeldii* within the family of Callitrichidae. *Cytogenetics and Cell Genetics* 94(3–4): 206–215. doi: 10.1159/000048818
- Rahn MI, Mudry MD, Merani MS, Solari AJ (1996) Meiotic behavior of the $X_1X_2Y_1Y_2$ quadrivalent of the Primate *Alouatta caraya*. *Chromosome Research* 4(5): 350–356. doi: 10.1007/BF02257270
- Rens W, O'Brien PC, Grützner F, Clarke O, Graphodatskaya D, Tsend-Ayush E, Trifonov VA, Skelton H, Wallis MC, Johnston S, Veyrunes F, Graves JA, Ferguson-Smith MA (2007) The multiple sex chromosomes of platypus and echidna are not completely identical and several share homology with the avian Z. *Genome Biology* 8(11): R243. doi: 10.1186/gb-2007-8-11-r243
- Rosenberger AL, Tejedor MF, Cooke SB, Pekar S (2009) Platyrrhine Ecophylogenetics in Space and Time. In: Garber PA, Estrada A, Bicca-Marques JC, Heymann EW, Strier KB, Barrett L (Eds) *South American Primates*. Springer, New York, USA, 69–113. doi: 10.1007/978-0-387-78705-3_4
- Rylands AB (2000) An assessment of the diversity of New World Primates. *Neotropical Primates* 8(2): 61–93.
- Rylands AB, Mittermeier RA (2009) The Diversity of the NewWorld Primates (Platyrrhini): An Annotated Taxonomy. In: Garber PA, Estrada A, Bicca-Marques JC, Heymann EW, Strier KB (Eds) *South American Primates: Comparative Perspectives in the study of behavior ecology, and conservation*. Springer, New York, USA, 23–54. doi: 10.1007/978-0-387-78705-3_2
- Sankoff D (2003) Rearrangements and chromosomal evolution. *Current Opinion in Genetics and Development* 13(6): 583–587. doi: 10.1016/j.gde.2003.10.006
- Seuánez HN (1979) *The phylogeny of human chromosomes*. Springer-Verlag, New York, USA, 208 pp. doi: 10.1007/978-3-642-67260-6
- Seuánez HN, Forman L, Matayoshi T, Fanning TG (1989) The *Callimico goeldii* (Primates, Platyrrhini) genome: Karyology and middle repetitive (LINE-1) DNA sequences. *Chromosoma* 98(6): 389–395. doi: 10.1007/BF00292783

- Smith JD (1970) The systematic status of the black howler monkey, *Alouatta pigra* Lawrence. *Journal of Mammalogy* 51(2): 358–369. <http://www.jstor.org/stable/1378486>, doi: 10.2307/1378486
- Solari AJ, Rahn MI (2005) Fine structure and meiotic behaviour of the male multiple sex chromosomes in the genus *Alouatta*. *Cytogenetics and Genome Research* 108(1-3): 262–267. doi: 10.1159/000080825
- Spix JB von (1823) Simiarum et vespertilionum Brasiliensium species novae ou Histoire Naturelle des Esèces Nouvelles de Singes et de Chauves-souris observées et recueillies pendant le voyage dans l'intérieur du Brésil. Munique, Typis Francisci Seraphici Hubschmanni, VIII+72 pp.
- Stanyon R, Consigliere S, Bigoni F, Ferguson-Smith M, O'Brien PCM, Wienberg J (2001) Reciprocal chromosome painting between a New World primate, the woolly monkey, and humans. *Chromosome Research* 9(2): 97–106. doi: 10.1023/A:1009274802086
- Stanyon R, Garofalo F, Steinberg ER, Capozzi O, Di Marco S, Nieves M, Archidiacono N, Mudry MD (2011) Chromosome Painting in Two Genera of South American Monkeys: Species Identification, Conservation, and Management. *Cytogenetics and Genome Research* 134(1): 40–50. doi: 10.1159/000324415
- Stanyon R, Rocchi M, Capozzi O, Roberto R, Misceo D, Ventura M, Cardone MF, Bigoni F, Archidiacono N (2008) Primate chromosome evolution: ancestral karyotypes, marker order and neocentromeres. *Chromosome Research* 16(1): 17–39. doi: 10.1007/s10577-007-1209-z
- Stanyon R, Tofanelli S, Morescalchi MA, Agoramoorthy G, Ryder OA, Wienberg J (1995) Cytogenetic analysis shows extensive genomic rearrangements between red howler (*Alouatta seniculus*, Linnaeus) subspecies. *American Journal of Primatology* 35(3): 171–183. doi: 10.1002/ajp.1350350302
- Steinberg ER, Cortés-Ortiz L, Nieves M, Bolzán AD, García-Orduña F, Hermida-Lagunes J, Canales-Espinosa D, Mudry MD (2008) The karyotype of *Alouatta pigra* (Primates: Platyrrhini): mitotic and meiotic analyses. *Cytogenetics and Genome Research* 122(2): 103–109. doi: 10.1159/000163087
- Steinberg ER, Nieves M, Mudry MD (2007) Meiotic characterization and sex determination system of neotropical primates: Bolivian squirrel monkey *Saimiri boliviensis* (Primates: Cebidae). *American Journal of Primatology* 69(11): 1236–1241. doi: 10.1002/ajp.20430
- Swofford D (2002) PAUP*. Phylogenetic analysis using parsimony (* and other methods). Sinauer Associates, Sunderland, Massachusetts. <http://paup.csit.fsu.edu/>
- Szapkievich V, Mudry MD (2003) Polimorfismos Genéticos en el mono aullador negro de la Argentina (*Alouatta caraya*). *Historia Natural* 2(7): 37–51.
- Thomas O (1912) On small mammals from the lower Amazon. *Annals and Magazine of Natural History* 9(8): 84–90. doi: 10.1080/00222931208693108
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22(22): 4673–4680. doi: 10.1093/nar/22.22.4673

- Torres OM, Leibovici M (2001) Caracterización del cariotipo del mono aullador colorado *Alouatta seniculus* que habita en Colombia. *Caldasia* 23(2): 537–548. <http://www.revistas.unal.edu.co/index.php/cal/article/view/17722>
- Torres OM, Ramírez C (2003) Estudios Citogenético de *Alouatta palliata* (Cebidae). *Caldasia* 25(1): 193–198. [http://www.unal.edu.co/icn/publicaciones/caldasia/25\(1\)/zoologia3.pdf](http://www.unal.edu.co/icn/publicaciones/caldasia/25(1)/zoologia3.pdf)
- Villalobos F, Valerio AA, Retana AP (2004) A phylogeny of howler monkeys (Cebidae: *Alouatta*) based on mitochondrial, chromosomal and morphological data. *Revista de Biología Tropical* 52(3): 665–677. http://www.scielo.sa.cr/scielo.php?script=sci_arttext&pid=S0034-77442004000300028&lng=es&nrm=iso
- Whittaker DJ, Morales JC, Melnick DJ (2007) Resolution of the *Hylobates* phylogeny: Congruence of mitochondrial D-loop sequences with molecular, behavioral, and morphological data sets. *Molecular Phylogenetics and Evolution* 45(2): 620–628. doi: 10.1016/j.ympev.2007.08.009
- Yunis EJ, De Caballero OMT, Ramírez C, Efraín Ramírez E (1976) Chromosomal Variations in the Primate *Alouatta seniculus seniculus*. *Folia Primatologica* 25(2-3): 215–224. doi: 10.1159/000155714
- Zunino GE, González V, Kowalewski MM, Bravo SP (2001) *Alouatta caraya*. Relations among habitat, density and social organizations. *Primate Report* 61: 37–46.

Appendix 1

Data matrix. (doi: 10.3897/CompCytogen.v8i1.6716.app1) File format: Microsoft Word file (doc).

Explanation note: Data matrix contains: Chromosomal data matrix, Molecular data matrix and Combined data matrix.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Citation: Steinberg ER, Nieves M, Mudry MD (2014) Multiple sex chromosome systems in howler monkeys (Mammalia, Platyrrhini). *Comparative Cytogenetics* 8(1): 43–69. doi: 10.3897/CompCytogen.v8i1.6716 Data matrix. doi: 10.3897/CompCytogen.v8i1.6716.app1

Appendix 2

Supplementary Figure S. (doi: 10.3897/CompCytogen.v8i1.6716.app2) File format: Microsoft Word file (doc).

Explanation note: Figure S: a) and b) Most parsimonious trees obtained for the chromosomal partition c) 50% majority consensus tree obtained by “bootstrap” d) 50% majority consensus tree obtained by “bootstrap” for the molecular partition.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Citation: Steinberg ER, Nieves M, Mudry MD (2014) Multiple sex chromosome systems in howler monkeys (Mammalia, Platyrrhini). *Comparative Cytogenetics* 8(1): 43–69. doi: 10.3897/CompCytogen.v8i1.6716 Supplementary Figure S. doi: 10.3897/CompCytogen.v8i1.6716.app2
