

Similarities and differences among the chromosomes of the wild guinea pig *Cavia tschudii* and the domestic guinea pig *Cavia porcellus* (Rodentia, Caviidae)

Laura I. Walker¹, Miguel A. Soto¹, Ángel E. Spotorno¹

¹ *Laboratorio de Citogenética Evolutiva, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile. Casilla 70061, Santiago 7, Chile*

Corresponding author: *Laura I. Walker* (laurainesw@yahoo.com)

Academic editor: *E. Gornung* | Received 13 March 2014 | Accepted 10 June 2014 | Published 14 July 2014

<http://zoobank.org/23FBF6F9-9055-4772-A9F4-5E02DD447F5B>

Citation: Walker LI, Soto MA, Spotorno ÁE (2014) Similarities and differences among the chromosomes of the wild guinea pig *Cavia tschudii* and the domestic guinea pig *Cavia porcellus* (Rodentia, Caviidae). *Comparative Cytogenetics* 8(2): 153–167. doi: 10.3897/CompCytogen.v8i2.7509

Abstract

Cavia tschudii Fitzinger, 1867 is a wild guinea pig species living in South America that according to the analysis of mitochondrial genes is the closest wild form of the domestic guinea pig. To investigate the genetic divergence between the wild and domestic species of guinea pigs from a cytogenetic perspective, we characterized and compared the C, G and AgNOR banded karyotypes of molecularly identified *Cavia tschudii* and *Cavia porcellus* Linnaeus, 1758 specimens for the first time. Both species showed 64 chromosomes of similar morphology, although *C. tschudii* had four medium size submetacentric pairs that were not observed in the *C. porcellus* karyotype. Differences in the C bands size and the mean number of AgNOR bands between the karyotypes of the two species were detected. Most of the two species chromosomes showed total G band correspondence, suggesting that they probably represent large syntenic blocks conserved over time. Partial G band correspondence detected among the four submetacentric chromosomes present only in the *C. tschudii* karyotype and their subtelocentric homologues in *C. porcellus* may be explained by the occurrence of four pericentric inversions that probably emerged and were fixed in the *C. tschudii* populations under domestication. The role of the chromosomal and genomic differences in the divergence of these two *Cavia* species is discussed.

Keywords

Karyotype, G, C and AgNOR banding, active NORs, pericentric inversions, domestication

Introduction

Cavia tschudii Fitzinger, 1867 is a wild species of guinea pig (Rodentia, Caviidae) which inhabits northern Chile, southern Peru and Bolivia and northwestern Argentina (Weir 1974, Woods and Kilpatrick 2005). The domestic guinea pig *Cavia porcellus* Linnaeus, 1758 has a cosmopolitan distribution and is an experimental animal, pet, and even is consumed as food in countries of the Andean Altiplano (Tello 1972).

There is a consensus that *C. porcellus* is a domestic form derived from one of the five currently recognized wild species of guinea pigs that inhabit South America (Woods and Kilpatrick 2005). The crosses between *C. porcellus* and *C. fulgida* Wagler, 1831 yielded offspring which behaved according to the Haldane's rule (Haldane 1922), since females were fertile and males were sterile (Detlefsen 1914). By contrast, the crosses between *C. porcellus* and *C. aperea sensu* Erxleben, 1777 (Pictet and Ferrero 1951, Rood 1972) and between *C. porcellus* and *C. cutleri* Tschudi, 1844 (*sensu* Bennet, 1836) (Castle 1916) produced hybrids which were fertile in both sexes. *C. aperea* or *C. tschudii* have been repeatedly considered as the most probable ancestor of the domestic guinea pig. Later, molecular analyses of the mitochondrial cytochrome *b* and 12S RNA genes clearly showed that the closest species to *C. porcellus* is *C. tschudii* and not the genetically related *C. aperea* (Spotorno et al. 2004, Dunnun and Salazar-Bravo 2010). Based on these molecular results and on the analysis of mummified guinea pig remains found in archeological sites, Spotorno et al. (2007) suggested that the domestication of the wild guinea pig occurred in southern Peru-northern Chile.

Considering that the karyotype provides useful characters in taxonomic and systematic studies and that changes in the number and structure of chromosomes may contribute to speciation (King 1993, Searle 1993, Capanna and Redi 1994, Capanna and Castiglia 2004, Marques-Bonet and Navarro 2005, Faria and Navarro 2010), we describe and compare now for the first time the G, C and AgNOR banded karyotypes in molecularly identified specimens of the wild montane guinea pig *C. tschudii* and the domestic guinea pig *C. porcellus*. Our objective is to discover the chromosomal and genomic differences between these two species of *Cavia* in relation to the divergence associated with the domestication process.

Material and methods

Skulls, skins and liver samples for DNA analysis of all the studied animals were preserved in the collection of the Laboratorio de Citogenética de Mamíferos, Facultad de Medicina, Universidad de Chile (LCM). We examined five *Cavia tschudii* specimens, four males (LCM 3199b, 3110, 3080, 3225) and one female (LCM 3232), collected in the locality of Molinos, Valle de Lluta, 18°23'S, 69°45'W, Arica, I Región, Chile, and four *Cavia porcellus* animals, two males (LCM 2454, 3192) from the laboratory Pirbright breed, Instituto de Salud Pública, Santiago, Chile, and two females (LCM 2479, 2489) from the Andean creole breed, Arica Agromarket, Arica, Chile.

Chromosomes were obtained from marrow cells using conventional in vivo colchicine, hypotonic method, preceded by yeast injection to improve the mitotic index (Lee and Elder 1980). Metaphase cells were G-banded and C-banded by the methods described by Chiarelli et al. (1972) and Sumner (1972), respectively. The active nucleolar organizing regions (NORs) were detected by the silver staining procedure (Sánchez-Rufas et al. 1982). At least 10 good-quality metaphases for each of the staining methods per taxon were selected under a light microscope and digitally captured and stored. Chromosomes were counted, cut out and ordered by size and form using ADOBE PHOTOSHOP version 6.0. The centromeric indexes calculated by measuring the chromosomal arms in 12 metaphases of each species, allowed the classification of the chromosomes as metacentric, submetacentric, subtelocentric and telocentric (Levan et al. 1964). Chromosomes of both species were ordered in the groups defined by Fernández and Spotorno (1968) for *C. porcellus* (groups A, B and C), adding a fourth group (group D) of submetacentric chromosomes for *C. tschudii* (Fig. 1). Male and female G-banded karyotypes from each species were compared and the chromosomes were classified as having totally corresponding, partially corresponding or unique G band patterns (Spotorno 1977, Walker et al. 1979). The size and distribution of the C and AgNOR bands were evaluated in six metaphases of *C. tschudii* and nine of *C. porcellus*. To determine the total number of active AgNOR in each of the species, AgNOR⁺ sites were identified and counted in the chromosomes of 24 metaphases per species. The statistical significance of the differences was estimated using a Chi squared test.

Results

Chromosome number, size and morphology

For *Cavia tschudii* and *Cavia porcellus* we consistently found a $2n=64$, $FNa=100-102$; the variation in the FNa of both species was due to the polymorphism of chromosome 1 (Fig. 1). *Cavia tschudii* showed five pairs of submetacentric chromosomes (group D, Fig. 1a) of which four pairs (numbers 27, 28, 29 and 30) were not present in the *C. porcellus* karyotype (Fig. 1b). The X chromosome of *C. tschudii* was a large submetacentric similar to that of *C. porcellus* and the Y chromosome was a subtelocentric larger than that of *C. porcellus* (Fig. 1).

G bands

The comparison of *C. tschudii* and *C. porcellus* G-banded karyotypes revealed total correspondence for 25 of the 31 autosomal pairs and for the X chromosomes of both species (Figs. 2, 3, Table 1). The four submetacentric chromosomes present only in the *C. tschudii* karyotype showed partial G band correspondence with four *C. porcel-*




Figure 1. Conventional stained karyotypes: **a** *Cavia tschudii* male **b** *Cavia porcellus* female. Chromosomal pair N° 1 shows subtelocentric morphology in other individuals of both species (**a** and **b** insets). Bar = 5 μ m.




Figure 2. G-banded karyotypes: **a** *Cavia tschudii* male **b** *Cavia porcellus* female. Chromosomes numbered according to original karyotype descriptions (see Fig. 1). Bar = 5 μ m.

lus subtelocentric chromosomes (Fig. 6, Table 1). Only the Y chromosomes and two autosomal pairs (*C. tschudii* chromosomes 12, 13 and *C. porcellus* chromosomes 14, 18) were unique of each species karyotype (Table 1).

C bands

The chromosomal distribution of the C bands was similar in the karyotypes of the two species, being located preferentially in the centromeres and the short arms of the chromosomes (Fig. 4). However, the amount of constitutive heterochromatin was ap-




Figure 3. Chromosomes of *Cavia tschudii* (Cts) and *Cavia porcellus* (Cpo) with total G band correspondence. Cts chromosomes are at the left and Cpo at the right of each chromosomal group. Note that the long arms of subtelocentric (St) and telocentric (T) forms of pair 1 show total G band correspondence (inset). Chromosomes numbered according to original karyotype descriptions (see Fig. 1). Bar = 5 µm.

Table I. Correspondence of *Cavia tschudii* and *Cavia porcellus* chromosomes according to their G band patterns¹.

Chromosomes with total G band correspondence	
Cts	1 2 3 4 5 6 7 8 9 10 11 14 15 16 17 18 19 20 21 22 23 24 25 26 31 X
Cpo	1 2 3 4 5 6 7 8 9 11 13 21 26 19 20 22 23 24 25 27 28 29 30 31 17 X
Chromosomes with partial G band correspondence	
Cts	27 28 29 30
Cpo	10 12 15 16
Unique species chromosomes	
Cts	12 13 Y
Cpo	14 18 Y

¹Chromosome numbers are the one of each species karyotype (see Fig. 1); in the same column chromosomes with total or partial G band correspondence. Cts = *Cavia tschudii*, Cpo = *Cavia porcellus*.

precipably greater in *C. tschudii* than in *C. porcellus*, spreading over most of the short arms in several subtelocentric chromosomes (Fig. 4a). The X chromosomes of both karyotypes, equal in size, morphology and G bands (Figs 1–3), showed a C⁺ band in the paracentromeric region of the short arm (Fig. 4). Both Y chromosomes were completely heterochromatic, being larger the Y chromosome of *C. tschudii* than the *C. porcellus* one (Fig. 4).




Figure 4. C-banded karyotypes: **a** *Cavia tschudii* male **b** *Cavia porcellus* male, showing heteromorphism for chromosome 1. Most of the chromosomes of both species were tentatively identified according size and morphology. Bar = 5 μ m.




Figure 5. AgNOR-banded karyotypes: **a** *Cavia tschudii* male with four nucleolar chromosomal pairs (4, 11, 14 and 29) **b** *Cavia porcellus* male with five nucleolar chromosomal pairs (1, 3, 5, 7 and 11). The nucleolar chromosomes of both species were tentatively identified according to their size and morphology. Bar = 5 μ m.

AgNOR bands

Multiple AgNOR bands were detected in the karyotypes of both species, consistently located in the telomeres of several chromosomal pairs (Fig. 5). The analysis of some AgNOR banded metaphases per species indicated that the number of AgNOR bands was different between the two species and also among the individuals. Thus, the results showed that in *C. porcellus* the mean and maximum numbers of chromosomes with active NORs (5.76 and 9.0, respectively) were higher than those of *C. tschudii* (4.13 and 7.0, respectively). Moreover, when we examined all the 3.072 chromosomes from 48 metaphases of both species, each of them having 64 chromosomes, we found a total




Figure 6. Rough simulation of the changes associated with the occurrence of pericentric inversions in *C. tschudii* chromosomes. Submetacentric *C. tschudii* chromosomes (Cts, first column at the left) that originate the subtelo-centric *C. porcellus* chromosomes (Cpo, last column at the right): 1°) chromosomal break, 2°) rotation of the cleaved segment, 3°) rejoining and sealing with the original segment. Chromosomes numbered according to original karyotype descriptions (see Fig. 1). Bar = 5 μ m.

of 237 AgNOR⁺ sites, 138 of them located in *C. porcellus* chromosomes and 99 in *C. tschudii* chromosomes. Accordingly, the number of chromosomes bearing active NOR was significantly higher in the *C. porcellus* karyotype than in the *C. tschudii* one ($\chi^2 = 6.956$; $p < 0.05$; $df = 1$).

Discussion

Cavia tschudii and *Cavia porcellus* diploid numbers ($2n=64$), previously described with basic cytogenetic techniques (Ohno et al. 1961, Fernández and Spotorno 1968, Dunnum and Salazar-Bravo 2006), were confirmed; nevertheless the fundamental number of autosomal arms (FNa = 100-102) were different to those reported before. For *C. porcellus*, Fernández and Spotorno (1968) described an FNa = 96, while for *C. tschudii*, Dunnum and Salazar-Bravo (2006) found an individual in the Bolivian Altiplano with an FNa which ranged from 104 to 108. The FNa variability of guinea pig species may be due to polymorphisms for the presence of short arms in the chromosomes described as subtelo-centric or telocentric in these species. The polymorphism for chromosome 1 short arms detected previously for *C. cobaya* Pallas, 1766, a synonym of *C. porcellus*,

(Ohno et al. 1961, Schmid 1965, Zenzes et al. 1977) and for both species in this study, as well as the report of entirely heterochromatic short arms in the subtelocentric autosomes of *C. porcellus* (Bianchi and Ayres 1971), give support to such hypothesis.

The number and morphology of *C. tschudii* and *C. porcellus* chromosomes were similar to those reported for other subspecies and species of the genus *Cavia*. So, with the exception of *C. intermedia* Cherem, Olimpio, Ximenez, 1999, and a population of *C. magna* Ximenez, 1980, having $2n=62$ (Gava et al. 1998, Cherem et al. 1999, Gava et al. 2012), for all of the other taxa of the genus the same $2n=64$ diploid number has been described, although with different numbers of autosomal arms. An $FNa=124$ was recorded for *C. aperea pamparum* (George et al. 1972); $FNa=116$ for *C. aperea aperea* specimens from Pernambuco, Brasil (Maia 1984) and 114 for some individuals from the Bolivian lowlands (Dunnum and Salazar-Bravo 2006); $FNa=124$ for *C. magna* and *C. fulgida* (Pantaleão 1978) and 114 for *C. nana* Thomas, 1917 (*Cavia tschudii sodalis*, 1926) (Dunnum and Salazar-Bravo 2006).

The analysis of the C bands showed that although they had a similar distribution in the chromosomes of the two species, they were smaller in size in the autosomes and in the Y chromosome of *C. porcellus* than in the *C. tschudii* ones, suggesting that a loss of heterochromatin occurred during the domestication process. In accordance with this result, measurements of the genome sizes of 31 hystricognath rodent species (Gallardo et al. 2003) indicated that the genome of *C. tschudii* (9.1 pg) is larger than that of *C. porcellus* (8.2 ± 0.4 pg), having the first species the largest genome size among the 30 diploid species analyzed.

Five chromosomal pairs bearing NOR at the short arm telomeres were found by Zenzes et al. (1977) in the karyotype of the domestic *Cavia*, so being in agreement with our results. Using a double-staining procedure they could identify those chromosomes as numbers 1, 3, 9, 12 and 14 of the quinacrine banded stained karyotype. An accurate identification of the *C. porcellus* and *C. tschudii* nucleolar chromosomes described here would require the use of a similar double-staining procedure to allow the comparisons with other descriptions.

The differences in the number of AgNOR bands found between the two *Cavia* species analyzed here and among the individuals in each of them, confirmed the tendency to variability in NOR expression usually described for mammals. It has been proposed that this variability would depend mainly on the specific metabolic demands of cells and individuals (Mikelsaar et al. 1977, Mayr et al. 1987, Sánchez et al. 1989, Suzuki et al. 1990, Berríos et al. 1992, Zurita et al. 1997, Walker et al. 1999, Walker and Flores 2007). The greater number of AgNOR bands found in the *C. porcellus* karyotype than in the *C. tschudii* one would reveal a greater transcriptional activity of the ribosomal genes in the genome of the domestic form. One possible functional explanation of this result is that since *C. porcellus* has been selected for productive purposes, it would require higher rates of protein synthesis than the wild form. It should be mentioned that in a recent comparison of brain gene expression levels between four pairs of domestic/wild mammals, the largest differences were found between the domestic and wild guinea pigs, although *C. aperea*, genetically related to *C. tschudii*, was used as the wild guinea pig species (Albert et al. 2012).

Comparison of the G-banded karyotypes of the two *Cavia* species included in this study revealed that most of the autosomal pairs and the X chromosomes showed total G band correspondence, suggesting that these chromosomes constitute large syntenic blocks present in the common ancestor of both species and conserved over time. The differences in morphology and the partial G band correspondences detected between four chromosomal pairs of these two species, suggest that the four submetacentric chromosomes present only in the *C. tschudii* karyotype would have suffered pericentric inversions originating the four subtelo-centric chromosomes of *C. porcellus* (Fig. 6, Table 1).

Cavia porcellus would be the domestic successor of *C. tschudii* from which it would have originated more than 4000 and possibly 7000 years ago (Wing 1986) by a process of domestication and artificial selection in the *C. tschudii* populations which inhabit southern Peru and northern Chile (Spotorno et al. 2007, Dunnum and Salazar-Bravo 2010). While being domesticated, those populations must have been small in size and with only a few individuals participating as parents in the reproductive process, which over many generations would have produced high levels of endogamy. These characteristics would have facilitated the fixation of the pericentric inversions which must have emerged spontaneously and frequently in the populations. Specifically, the heterozygotes for the pericentric inversions would have decreased their fertility since their gametes would be unbalanced as a result of crossing-over in the inverted segment (Coyne et al. 1993, King 1993). As a consequence, gene flow between the original homozygotes and the homozygotes for the inversion would have been reduced, originating genetic divergence between the two chromosomal forms.

Nevertheless, it has been demonstrated recently that the fertility of the inversion carriers is not always reduced (Muss and Schwantz 2007). In some cases and depending on the size, genetic content, and chromosomal location of the inversion, the chromosomal inverted region pairs non-homologously with its normal partner forming a straight bivalent which does not present any loop, so causing crossing-over suppression (Torgasheva and Borodin 2010). If that is the case, the absence or reduced recombination between the inverted and non inverted genomic regions in the *Cavia* pericentric inversions, would be the cause of genetic divergence accumulation and reduction of gene flow between the two chromosomal forms, as it was proposed as a general model of speciation by several authors (Noor et al. 2001, Navarro and Barton 2003, Hoffman and Rieseberg 2008).

A critical assessment of reproductive isolation in crosses between *Cavia* species as previously reported, confronts the appropriate identification of specimens, the reliability of the taxonomy at the time, and the nature of the differences eventually found. For instance, crosses between *C. porcellus* and individuals from Arequipa, Perú identified as *C. cutleri* Bennett, 1836, which correspond to the original description of *C. cutleri* based on a single specimen from Ica, Peru (see Weir 1974), produced fertile offspring according to Castle (1916). By contrast, other wild specimens from Ica, Peru that also received the name *C. cutleri* by Tschudi in 1849, were finally renamed as *C. tschudii* by Fitzinger in 1867 (see Weir 1974); the latter is now the usually accepted name for the wild montane guinea pig (Woods and Kilpatrick 2005). Therefore, *C. cutleri* Bennett, 1838 is now considered a synonym of *C. porcellus* (Woods and Kilpatrick 2005). If

Castle in fact crossed *C. porcellus* laboratory animals with *C. cutleri* Bennet (= *C. porcellus*) specimens, in reality he might be doing intraspecific crosses, and the fertility of the descendants would be an expected result. In any case, the assignation of the individuals from Arequipa to *Cavia cutleri* Bennet was not well documented in that study, since it was based only on the smaller body size of those individuals with respect to domestic *C. porcellus* (Castle 1916). Moreover, he did not indicated the mating times taken by the crosses, neither the number of pairs in which crosses were attempted, reporting only that a large number of descendants were obtained (n=107), as many as those obtained in crosses within each form (n=108). In sum, if the chromosomal and nucleolar differences we are reporting here in molecularly identified specimens of *C. tschudii* and *C. porcellus* were also found in other populations, we predict that their eventual hybrids will show some degree of genomic incompatibility.

Reproduction of wild mammal species in captivity is a difficult and not always successful task. It is even more difficult to obtain descendants from crosses between different chromosomal races or species in the laboratory (Walker et al. 1984, 1999, Hauffe and Searle 1998, Castiglia and Capanna 2000, Franchini et al. 2008, Nunes et al. 2011). In crosses between phyllotine rodent species, we reported previously a decrease in the proportion of pairs with births and in the litter's size together with an increase in the time between mating and birth, compared to those registered for the intraspecific crosses (Walker et al. 1984, 1999). Although we repeatedly tried to cross our specimens of *C. tschudii* with *C. porcellus* in our laboratory, we have had no success yet.

If the analysis of the crosses realized between chromosomal races of *Mus* and *Sorex* rodents (Hauffe and Searle 1998, Castiglia and Capanna 2000, Franchini et al. 2008, Nunes et al. 2011) would have only considered the number of descendants obtained, it would not have been possible to reach conclusions about the fertility level of those hybrids. To estimate fertility, specific reproductive aspects must be studied, such as the success obtained in crosses between the parental forms and some hybrid characters, i.e.: the normality of their meiotic process, the histology of their gonads, the cell composition of their germinal line and the chromosomal constitution of the gametes that eventually they produce. In consequence, to evaluate the fertility level of eventual *C. tschudii* × *C. porcellus* hybrids, the reproductive characters just mentioned above must be analyzed in the descendants of crosses between individuals of the parental species taxonomically well identified. Specifically, the fertility of the heterozygotes for the pericentric inversions described here should be further investigated to evaluate the contribution of those chromosomal changes to the divergence of the two *Cavia* species.

Acknowledgements

This study was partially financed by Project FONDECYT 1011052. We thank the Servicio Agrícola y Ganadero for the permission to capture the wild animals, Dr. Mónica Acevedo for her technical assistance and Juan Oyarce for his help in collection and maintenance of the animals.

References

- Albert FW, Somel M, Carneiro M et al. (2012) A comparison of brain gene expression levels in domesticated and wild animals. *PLoS Genetics* 8(9): e1002962. doi:10.1371/journal.pgen.1002962
- Bennett ET (1836) On a new species of *Ctenomys* and other rodents collected near the Straits of Magellan by Capt. P.P. King, R.N. *Proceedings of the Zoological Society of London* 1835 (part 3, no. 36): 189–191. [Part III, no. XXXVI of the 1835 volume of the *Proceedings* was published April 8, 1836]
- Berrios S, Koifman J, Fernández-Donoso R (1992) Tissue and sex differences in the expression of nucleoli in mouse somatic cells. *European Journal of Morphology* 30: 297–303.
- Bianchi NO, Ayres J (1971) Polymorphic patterns of heterochromatin distribution in guinea pig chromosomes. *Chromosoma (Berlin)* 34: 254–260. doi:10.1007/BF00286151
- Capanna E, Redi CA (1994) Chromosomes and microevolutionary processes. *Italian Journal of Zoology* 61: 285–294.
- Capanna E, Castiglia R (2004) Chromosomes and speciation in *Mus musculus domesticus*. *Cytogenetic and Genome Research* 105: 375–384. doi: 10.1159/000078210
- Castiglia R, Capanna E (2000) Contact zone between chromosomal races of *Mus musculus domesticus*. Fertility and segregation in laboratory-reared and wild mice heterozygous for multiple Robertsonian rearrangements. *Heredity* 85: 174–156. doi: 10.1046/j.1365-2540.2000.00743.x
- Castle WE (1916) Size inheritance in guinea pig crosses. *Proceedings of the National Academy of Sciences USA* 2: 252–264. doi: 10.1073/pnas.2.4.252
- Cherem JJ, Olimpio J, Ximenez A (1999) Descrição de uma nova espécie do gênero *Cavia* Pallas 1766 (Mammalia - Caviidae) das Ilhas dos Moleques do Sul, Santa Catarina, Sul do Brasil. *Biotemas (Brasil)* 12: 95–117.
- Chiarelli B, Sarti-Chiarelli M, Shafer D (1972) Chromosome banding with trypsin. *Mammalian Chromosome Newsletter* 13: 44–45.
- Coyne JA, Meyers W, Crittenden AP, Sniegowski P (1993) The fertility effects of pericentric inversions in *Drosophila melanogaster*. *Genetics* 134: 487–496.
- Detlefsen JA (1914) Genetic studies on a cavy species cross. *Carnegie Institution of Washington Publications* 205: 1–134.
- Dunnum JL, Salazar-Bravo J (2006) Karyotypes of some members of the genus *Cavia* (Rodentia: Caviidae) from Bolivia. *Mammalian Biology* 71: 336–370. doi: 10.1016/j.mambio.2006.04.006
- Dunnum JL, Salazar-Bravo J (2010) Molecular systematics, taxonomy and biogeography of the genus *Cavia* (Rodentia: Caviidae). *Journal of Zoological Systematics and Evolutionary Research* 48: 376–388. doi: 10.1111/j.1439-0469.2009.00561.x
- Erxlebe JCP (1777) *Systema regni animalis per classes, ordines, genera, species, varietates, cum synonymia et historia animalium. Classis I. Mammalia. Weygandianis, Lipsiae*, 636 pp. doi: 10.5962/bhl.title.15933
- Faria R, Navarro A (2010) Chromosomal speciation revisited: rearranging theory with pieces of evidence. *Trends in Ecology and Evolution* 25: 660–669. doi: 10.1016/j.tree.2010.07.008

- Fernández R, Spotorno A (1968) Heteromorfismo del par cromosómico N° 1 de *Cavia porcellus* L. *Archivos de Biología y Medicina Experimentales* 5: 81–85.
- Fitzinger LJ (1867) *Versuche in der natürlichen Anordnung der Nagethiere (Rodentia)*. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-naturwissenschaftliche Classe 55: 453–515; 56: 57–168.
- Franchini P, Castiglia R, Capanna E (2008) Reproductive isolation between chromosomal races of the house mouse *Mus musculus domesticus* in a parapatric contact area revealed by an analysis of multiple unlinked loci. *Journal of Evolutionary Biology* 21: 502–513. doi: 10.1111/j.1420-9101.2007.01492.x
- Gallardo MH, Bickham JW, Kausel G, Köhler N, Honeycutt RL (2003) Gradual and quantum genome size shifts in the hystricognath rodents. *Journal of Evolutionary Biology* 16: 163–169. doi: 10.1046/j.1420-9101.2003.00507.x
- Gava A, Freitas TRO, Olimpio J (1998) A new karyotype for the genus *Cavia* from a southern island of Brazil (Rodentia-Caviidae). *Genetics and Molecular Biology* 21: 77–80. doi: 10.1590/S1415-47571998000100013
- Gava A, Dos Santos M, Quintela FM (2012) A new karyotype for *Cavia magna* (Rodentia: Caviidae) from an estuarine island and *C. apera* from adjacent mainland. *Acta Theriologica* 57: 9–14. doi: 10.1007/s13364-011-0042-0
- George W, Weir BJ, Beadford J (1972) Chromosome studies in some members of the family Caviidae (Mammalia: Rodentia). *Journal of Zoology, London* 168: 81–89. doi: 10.1111/j.1469-7998.1972.tb01339.x
- Haldane JBS (1922) Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics* 12: 101–109. doi: 10.1007/BF02983075
- Hauffe HC, Searle JB (1998) Chromosomal heterozygosity and fertility in house mice (*Mus musculus domesticus*) from Northern Italy. *Genetics* 150: 1143–1154.
- Hoffman AA, Rieseberg LH (2008) Revisiting the impact of inversions in evolution: from genetic markers to drivers of adaptive shifts and speciation. *Annual Review of Ecology, Evolution and Systematics* 39: 21–42. doi:10.1146/annurev.ecolsys.39.110707.173532
- King M (1993) *Species evolution. The role of chromosome change*. Cambridge University Press, New York, 336 pp.
- Lee M, Elder F (1980) Yeast stimulation of bone marrow mitoses for cytogenetic investigations. *Cytogenetic and Cell Genetics* 26: 36–40. doi: 10.1159/000131419
- Levan A, Fredga D, Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201–220. doi: 10.1111/j.1601-5223.1964.tb01953.x
- Linnaeus C (1758) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. 10th Edition, Laurentii Salvi, Stockholm.
- Maia V (1984) Karyotypes of three species of Caviinae (Rodentia: Caviidae). *Experientia* 40: 564–566. doi: 10.1007/BF01982332
- Marques-Bonet T, Navarro A (2005) Chromosomal rearrangements are associated with higher rates of molecular evolution in mammals. *Gene* 353: 147–154. doi: 10.1016/j.gene.2005.05.007

- Mayr B, Schleger W, Auer H (1987) Frequency of Ag-stained nucleolus organizer regions in the chromosomes of cattle. *Journal of Heredity* 78: 206–207.
- Mikelsaar AV, Schmid M, Krone H, Schwarzacher G, Schenedl W (1977) Frequency of Ag-stained nucleolus organizer regions in the acrocentric chromosomes of man. *Human Genetics* 37: 73–77. doi: 10.1007/BF00293774
- Muss B, Schwanitz G (2007) Characterization of inversions as a type of structural chromosome aberration. *International Journal of Human Genetics* 7: 141–161.
- Navarro A, Barton NH (2003) Chromosomal speciation and molecular divergence -Accelerated evolution in rearranged chromosomes. *Science* 300: 321–324. doi: 10.1126/science.1080600
- Noor MAF, Grams KL, Bertucci LA, Reiland J (2001) Chromosomal inversions and the reproductive isolation of species. *Proceedings of National Academy of Sciences* 98: 12084–12088. doi: 10.1073/pnas.221274498
- Nunes AC, Catalan J, Lopez J, Ramalhinho MG, Mathias ML, Britton-Davidian J (2011) Fertility assessment in hybrids between monobrachially homologous Rb races of the house mouse from the island of Madeira: implications for modes of chromosomal evolution. *Heredity* 106: 348–356. doi: 10.1038/hdy.2010.74
- Ohno S, Weiler C, Stenius C (1961) A dormant nucleolus organizer in the guinea pig, *Cavia cobaya*. *Experimental Cell Research* 25: 498–503.
- Pallas PS (1766) *Miscellanea zoologica quibus novae imprimis atque obscurae animalium species descriuntur et observationibus iconibusque illustrantur*. Hague Comitum: P. van Cleef, xii + 224 pp. [14 plates]
- Pantaleão E (1978) Caracterização de espécies do gênero *Cavia* por análise de seus cariótipos. MSc Dissertation, Universidade Federal do Rio Grande do Sul, Porto Alegre Brazil.
- Pictet A, Ferrero A (1951) La descendance d'un croisement interespecificque de cobayes (*Cavia aperea* D'Az x *Cavia cobaya* Marc) analysé durant 25 années. *Genetica* 25: 357–515. doi: 10.1007/BF01784834
- Rood JP (1972) Ecological and behavioural comparisons of three genera of Argentine cavies. *Animal Behaviour Monographs* 5: 1–83. doi: 10.1016/S0066-1856(72)80002-5
- Sánchez A, Burgos M, Jiménez R, Díaz de la Guardia R (1989) Quantitative analysis of silver staining of the nucleolar organizer regions in *Eliomys quercinus*. *Genome* 32: 978–982. doi: 10.1139/g89-541
- Sánchez-Rufas J, Iturra P, De Souza W, Esponda P (1982) Simple silver staining procedures for the location of nucleolus and nucleolar organizer under light and electron microscopy. *Archives of Biology* 93: 267–274
- Schmid W (1965) Heterochromatin in mammals. *Archiv der Julius Klaus-Stiftung für Vererbungsforshung, Sozialanthropologie und Rassenhygiene*. 40: 35–49.
- Searle JB (1993) Chromosomal hybrid zones in eutherian mammals. In: Harrison RG (Ed) *Hybrid zones and the evolutionary process*. Oxford University Press, Oxford, 309–353.
- Spotorno AE (1977) Phylogenetic partitioning of banded karyotypes in mammals: a model of cladistic analysis. In: Drets ME, Brum-Zorrilla N, Folle GA (Eds) *Third Latin-American Congress of Genetics*, Montevideo, February 2-12, 1977, UNESCO: 179–187.

- Spotorno AE, Valladares J, Marín J, Zeballos H (2004) Molecular diversity among domestic guinea-pigs (*Cavia porcellus*) and their close phylogenetic relationship with the Andean wild species *Cavia tschudii*. *Revista Chilena de Historia Natural* 77: 243–250. <http://www.scielo.cl/pdf/rchnat/v77n2/art04.pdf>
- Spotorno AE, Manríquez G, Fernández LA, Marín JC, González F, Wheeler J (2007) Domestication of guinea pigs from a southern Peru-northern Chile wild species and their middle pre-Columbian mummies. In: Kelt DA, Lessa EP, Salazar-Bravo J, Patton JL (Eds) *The quintessential naturalist: honoring the life and legacy of Oliver P Pearson*. University of California Publications in Zoology 134: 367–388. <http://repositories.cdlib.org/ucpress/>
- Sumner A (1972) A simple technique for demonstrating centromeric heterochromatin. *Experimental Cell Research* 75: 304–306. doi:10.1016/0014-4827(72)90558-7
- Suzuki H, Kurihara Y, Kanehisa T, Moriwaki K (1990) Variation in the distribution of silver-stained nucleolar organizing regions on the chromosomes of the wild mouse *Mus musculus*. *Molecular Biology and Evolution* 7: 271–282.
- Tello AV (1972) Efecto de cuatro raciones concentradas en el crecimiento de cuyes (*Cavia porcellus*). Tesis Universidad Nacional Agraria La Molina, Lima, Perú, 53 pp.
- Thomas O (1917) Notes on the species of the genus *Cavia*. *Annals and Magazine of Natural History*, ser. 8, 19: 152–160.
- Torgasheva AA, Borodin PM (2010) Synapsis and recombination in inversion heterozygotes. *Biochemical Society Transactions* 38: 1676–1680. doi: 10.1042/BST0381676
- Wagler J (1831) Einige Mitteilungen über Thiere Mexicos. *Isis von Oken* 24(6): 510–534.
- Walker LI, Spotorno A, Fernández-Donoso R (1979) Conservation of whole arms during chromosomal divergence of phyllotine rodents. *Cytogenetic and Cell Genetics* 24: 209–216. doi: 10.1159/000131382
- Walker LI, Spotorno A, Arrau J (1984) Cytogenetic and reproductive studies of two nominal subspecies of *Phyllotis darwini* and their experimental hybrids. *Journal of Mammalogy* 65: 220–230. doi: 10.2307/1381161
- Walker LI, Rojas M, Flores S, Spotorno A, Manríquez G (1999) Genomic compatibility between two phyllotine rodent species evaluated through their hybrids. *Hereditas* 131: 227–238. doi: 10.1111/j.1601-5223.1999.00227.x
- Walker LI, Flores SV (2007) Nucleolar activity and distribution of ribosomal genes in *Phyllotis* rodent species and their laboratory hybrids. In: Kelt D, Lessa EP, Salazar-Bravo J, Patton JL (Eds) *The quintessential naturalist: honoring the life and legacy of Oliver P Pearson*. University of California Publications in Zoology 134: 901–916. <http://repositories.cdlib.org/ucpress/>
- Weir BJ (1974) Notes on the origin of the domestic guinea pig. *Symposia of the Zoological Society of London* 34: 437–446.
- Wing E (1986) Domestication of Andean mammals. In: Vuilleumier F, Monasterio M (Eds) *High altitude tropical biogeography*. Oxford University Press, New York, 246–264.
- Woods CA, Kilpatrick CW (2005) Infraorder Hystricognathi. In: Wilson DE, Reeder DM (Eds) *Mammal species of the World: a taxonomic and geographic reference*. Johns Hopkins University Press, Baltimore, Maryland, 1538–1600.

- Ximenez A (1980) Notas sobre el género *Cavia* Pallas con la descripción de *Cavia magna* sp. n. (Mammalia-Caviidae). *Revista Nordestina de Biología* 3: 145–179.
- Zenzes MT, Schmid M, Engel W (1977) Silver-stained nucleolus organizers in the guinea pig, *Cavia cobaya*. *Cytogenetics and Cell Genetics* 19: 368–72. doi: 10.1159/000130830
- Zurita F, Sánchez A, Burgos M, Jiménez R, Díaz de la Guardia R (1997) Interchromosomal, intercellular and interindividual variability of NORs studied with silver staining and in situ hybridization. *Heredity* 78: 229–234. doi: 10.1038/hdy.1997.36