

Genomic homeology between *Pennisetum purpureum* and *Pennisetum glaucum* (Poaceae)

Gabriela Barreto dos Reis¹, Amanda Teixeira Mesquita¹, Giovana Augusta Torres¹, Larissa Fonseca Andrade-Vieira¹, Antônio Vander Pereira², Lisete Chamma Davide¹

1 Department of Biology, Federal University of Lavras, Zip Code 37200-000, Lavras, MG, Brazil **2** Laboratory of Plant Genetics, Brazilian Agricultural Research Corporation, Embrapa Dairy Cattle, Zip Code 36038-330, Juiz de Fora, MG, Brazil

Corresponding author: Lisete Chamma Davide (lisete.ufla@gmail.com)

Academic editor: Wei Wenhui | Received 15 April 2014 | Accepted 14 July 2014 | Published 8 August 2014

<http://zoobank.org/2AFB030F-49E6-4888-86DB-B28A3B387E3D>

Citation: dos Reis GB, Mesquita AT, Torres GA, Andrade-Vieira LF, Pereira AV, Davide LC (2014) Genomic homeology between *Pennisetum purpureum* and *Pennisetum glaucum* (Poaceae). Comparative Cytogenetics 8(3): 199–209. doi: 10.3897/CompCytogen.v8i3.7732

Abstract

The genus *Pennisetum* (Richard, 1805) includes two economically important tropical forage plants: *Pennisetum purpureum* (Schumacher, 1827) (elephant grass), with $2n = 4x = 28$ chromosomes and genomes A'A'BB, and *Pennisetum glaucum* (Linnaeus, 1753) (pearl millet), with $2n = 2x = 14$ chromosomes and genomes AA. The genetic proximity between them allows hybrids to be obtained ($2n = 3x = 21$) that yield forage of higher quality in relation to the parents. The study of genomic relationships provides subsidies for the knowledge about phylogenetic relations and evolution, and is useful in breeding programs seeking gene introgression. Concerning elephant grass and pearl millet, the homeology between the genomes A and A', and between these and the genome B, has been reported by conventional cytogenetic techniques. The objective of the present study was to demonstrate the degree of homeology between these genomes by means of genomic *in situ* hybridization (GISH). The results confirmed the homeology between the genomes A of pearl millet and A'B of elephant grass, and showed that there are differences in the distribution and proportion of homologous regions after hybridization. Discussion regarding the evolutionary origin of *P. purpureum* and *P. glaucum* was also included.

Keywords

Homeology, *Pennisetum purpureum*, *Pennisetum glaucum*, Genomic *in situ* hybridization

Introduction

The genus *Pennisetum* (Richard, 1805) is one of the most important in family Poaceae family. It comprises about 140 species, distributed in five sections (*Penicillaria*, *Brevivalvula*, *Gymnothrix*, *Heterostachya* and *Eu-Pennisetum*) based on morphological characteristics (Stapf and Hubbard 1934). The section *Penicillaria* includes the economically most important species: elephant grass [*Pennisetum purpureum* (Schumacher, 1827)], used as forage, and pearl millet [*Pennisetum glaucum* (Linnaeus, 1753)], used as cereal and forage (Martel et al. 2004).

Molecular analyses based on mitochondrial DNA (Chowdhury and Smith 1988), chloroplast DNA (Renno et al. 2001) and repetitive DNA sequences (Ingham et al. 1993) have revealed significant relationship between the genomes of *P. glaucum*, *P. purpureum* and *Pennisetum squamulatum* Fresen., suggesting that these three species may have a common origin (Martel et al. 2004). Among these, the cultivated species *P. glaucum* and *P. purpureum* are phylogenetically related, possessing a close resemblance between their genomes, constituting a monophyletic group with recent divergence (Martel et al. 2004). In this sense, *P. glaucum* is an annual, alogamous, diploid species ($2n = 2x = 14$, genomes AA), with genome DNA content of 4.72 pg, and constitutes the primary genic pool of this genus. In turn, *P. purpureum* belongs to the secondary genic group and is a perennial, alogamous, tetraploid species ($2n = 4x = 28$, genomes A'A'BB), with genome DNA content of 4.60 pg (Martel et al. 1997). Both species have approximately the same DNA content (pg), but differ with regards to the monoploid size genome (basic number, x). *P. purpureum* is a tetraploid and have about half the DNA content (1.15 pg) of the *P. glaucum* monoploid genome (2.36 pg). Further, *P. purpureum* have smaller chromosomes than *P. glaucum* (Martel et al. 2004). This shows that important chromosome changes may be linked to the evolution and divergence among these species (Andrade-Vieira 2010; Barbosa et al. 2003; Martel et al. 2004; Robert et al. 2011).

In spite of their integrating distinct genic groups and differing as to ploidy level, the genetic proximity between these two species becomes evident when the occurrence of natural hybridization is observed. This sexual compatibility is partial, and results in sterile triploid hybrids ($2n = 3x = 21$, genome AA'B) (Hanna 1987; Martel et al. 2004; Robert et al. 2011; Techio et al. 2006). Cytologically, the genomic proximity has been demonstrated by meiotic analyses of triploids hybrids obtained in breeding programs. During diakinesis and metaphase I in this hybrid, the formation of seven bivalents is frequently observed, resulting from the pairing among chromosomes of genomes A and A' of *P. glaucum* and *P. purpureum*, respectively, as well as seven univalents of genome B of *P. purpureum* (Jauhar 1968, 1981; Pantulu 1967; Sree Rangasamy 1972; Techio et al. 2005, 2006). The occurrence, even at low frequency, of trivalents and numbers of bivalents above seven suggests both allo- as well as autosyndetic pairing among the genomes A, A' and B (Sethi et al. 1970; Techio et al. 2005, 2006). These observations on the configurations of bivalents and univalents during meiosis, as well as the morphology of metaphase chromosomes, constitute the

pioneering works demonstrating homeology among the genomes A, A' and B. The obtained results suggest that the genome A of *P. glaucum* has larger homeology with genome A' of *P. purpureum*, whereas the origin of genome B has not been defined (Jauhar 1981; Techio et al. 2005, 2006).

In this sense, despite evidence for a common evolutionary origin between *P. glaucum* and *P. purpureum* and the economic importance of these species, there are no studies providing more conclusive data with respect to the homeology among genomes A, A' and B. Thus, the objective of this work is to describe the proportion and distribution of the homologous regions present in genomes A of *P. glaucum* and A'B of *P. purpureum*, by cytomolecular analyses using genomic *in situ* hybridization (GISH).

Material and methods

Plant material and genomic DNAs

The evaluations were carried out in mitotic metaphases of the parental *P. purpureum* (access BAG 65) and *P. glaucum* (access BN2), and of the triploid hybrid originating from this crossing (BAG 65 × BN2). The plant material and genomic DNAs were provided by the Active Germplasm Bank of Elephant Grass (BAGCE) from EMBRAPA Dairy Cattle (Brazilian Research Institute) and elephant grass breeding program, experimental field José Henrique Bruschi, municipality of Coronel Pacheco, Minas Gerais State, Brazil.

Chromosome preparation

Roots from seeds or cuttings of BAG 65, BN2 and triploid hybrid accession were collected and pretreated with a 12.5 mg.L⁻¹ cycloheximide : 150 mg.L⁻¹ 8-hydroxyquinoline solution for 2 h 45 min, at 4 °C, and fixed in ethanol : acetic acid solution (3:1), as proposed by Techio et al. (2002). Fixed root tips were digested with pectinase : cellulase (100U:200U) solution in citrate–phosphate buffer (pH 4.8) for 40 min (*P. glaucum*) and 3 h 30 min (*P. purpureum* and interspecific hybrid), at 37 °C, in moist chamber. Slides were prepared as proposed by Dong et al. (2000). A root tip was transferred to a slide and macerated with a drop of ethanol : acetic acid solution (2:1) using a fine-pointed forceps. The slide then was warmed over an alcohol flame. It could called flame-drying method.

Genomic *in situ* hybridization

Genomic DNAs of *P. glaucum* and *P. purpureum* were labeling with biotin-16-dUTP through nick-translation reaction method, thus yielding the genomic probes.

The hybridization technique was carried out according to Jiang et al. (1995). The hybridization mixture [55% formamide (v/v), 10% dextran sulfate (w/v), 2X SSC, pH=7.0, and 2 μ L of probe marked with biotin] was denatured at 95 °C, for 8 min. Chromosome preparation was denatured with 70% formamide in 2X SSC (saline sodium citrate) at 85 °C, for 1 min 20 sec (Andrade-Vieira et al. 2013) and hybridized in the mixture at 37 °C for, at least, 16 h in a moist chamber. Detection of the probe marked with biotin was performed with streptavidin conjugated with Alexa Fluor® 488. Chromosomes were counterstained with 1 μ g.mL⁻¹ 4',6-diamidino-2-phenylindole (DAPI) Vectashield® antifade solution (Vector Laboratories). The slides were evaluated under an epi-fluorescence Nikon Eclipse E600 microscope. Images of interest were digitized by means of a refrigerated monochromatic Nikon DSQi1MC camera, and processed using the software NIS-Element BR 4.00.03 (Nikon) and Adobe Photoshop CS3.

In order to evaluate the level of homeology between genomes A, A' and B, the chromosomes of five metaphases from each genome were measured, as well as the proportion occupied by the genomic probe, using the Image Tool 3.0 program. The obtained data were used to create karyograms for comparison of the evaluated genomes.

Results and discussion

Previous analyses of meiotic pairing in the triploid hybrid have showed that the genomes A and A' are more related. On the other side, between both and the genome B there are affinity/homeology reduced (Jauhar and Hanna 1998, Techio et al. 2005). In this study, hybridization of genomic probes of *P. glaucum* and *P. purpureum* were used for the first time to demonstrate and to confirm homeology among the three genomes. It was evaluated the distribution and proportion of these homeologous regions in the family constituted by the parental *P. purpureum* (BAG 65) and *P. glaucum* (BN2), and by the triploid hybrid (BAG 65 \times BN2) originating from this crossing.

The higher level of homeology between genomes A and A' was confirmed because the 14 chromosomes belonging to genome A' of *P. purpureum* were strongly marked and distinguished from the 14 chromosomes from genome B using the genomic DNA of *P. glaucum* (genome A) as probe in metaphases of *P. purpureum* (Fig. 1a). The chromosomes of genome A' presented marks in along almost role chromosome length, whereas genome B presented small marks dispersed over the length of its chromosomes (Fig. 2a). Moreover, approximately 29% of *P. purpureum* genome (A'B) was hybridized by the genome A of *P. glaucum* (Table 1). This percentual represents only the A' genome since the markers on genome B chromosomes were not record because it were dispersed on chromosomes. The observed homeology was only quantified in the genome A' of *P. purpureum*, due to the difficulty in measuring the small and dispersed marks found in genome B (Fig. 2a).

The homeology between genomes A and A' was confirmed by the extensive marking of *P. glaucum* chromosomes by the probe A'B of *P. purpureum*. All 14 chromosomes from genome A of *P. glaucum* were almost completely marked, with large blocks of probe sig-




Figure 1. Metaphases of *Pennisetum purpureum* (**A**), *Pennisetum glaucum* (**B**), and triploid hybrid (**C** and **D**). Chromosomes stained with DAPI (**A**, **B**, **C**, **D**) and probe markings in chromosomes indicated by green fluorescence (**A1**, **B1**, **C1**, **D1**). (**A1**) chromosomes of *P. purpureum* hybridized with genomic probe of *P. glaucum* (genome **A**), (**B1**) chromosome of *P. glaucum* hybridized with genomic probe of *P. purpureum* (genomes **A'B**), (**C1**) chromosomes of the triploid hybrid hybridized with genomic probe of *P. glaucum* (genome **A**), (**D1**) chromosomes of the triploid hybrid hybridized with genomic probe of *P. purpureum* (genomes **A'B**). Bar = 10 μ m (**A**); Bar = 20 μ m (**B**, **C** and **D**).




Figure 2. Karyograms of *Pennisetum purpureum* (**A**), *Pennisetum glaucum* (**B**) and triploid hybrid (**C**) identifying the chromosomes of genomes A, A' and B in each genotype. Note that in (**A**) using genome A probe (*P. glaucum*), the chromosomes of genome A' were differed from chromosomes of genome B by the staining pattern. Genome A' chromosomes showed more apparent probe markings in green than genome B chromosomes. In (**B**), using the genome A'B probe (*P. purpureum*), all chromosomes were strongly labelled (markings in green). In (**C**), using the genome A probe (*P. glaucum*), the chromosomes of the A genome were fully labeled by the probe (markings in green), the genome A' were strongly marked in the centromeric region and the genome B, poorly marked. It also could be note the difference in the labeling pattern between the genome A probe on the chromosomes of genome A' in interspecific hybrid and parental *P. purpureum*. Bar = 10 μ m.

nals observed on the chromosomes (Fig. 1b). The markings by the probe of genome A'B observed in the centromeric and pericentric regions represented 63% of the genome of *P. glaucum* (Table 1 and Fig. 2b). These marked portions result from hybridization, both between the genomes A and A' and, in smaller proportion, genomes A and B, observed both in karyograms of *P. purpureum* and triploid hybrids (Fig. 2a, c).

Table 1. Proportion of markings of genomic probes (A and A'B) on chromosomes of *Pennisetum purpureum*, *Pennisetum glaucum* and triploid hybrid.

Genotype	Total length of the chromosomes	Total length of the probe <i>P. glaucum</i> (A)	Total length of the probe <i>P. purpureum</i> (A'B)
<i>P. purpureum</i>	64,41	18,42 (28,60%)*	-
<i>P. glaucum</i>	59,01	-	37,29 (63,19%)
Triploid hybrid	73,95	40,06 (54,19%)	36,32 (49,13%)

* Proportion occupied by the probe for each genotype

In the triploid hybrid (AA'B) the hybridized portion of genome A (*P. glaucum*) corresponded to 54%, and the signal of probe A'B (*P. purpureum*) to 49% of its total genome (Table 1). Despite the similarity in proportion, the distribution pattern for the probes from the parental individuals was different in the hybrid (Fig 1c). The seven chromosome of the hybrid were entirely marked with the probe of genomic DNA from *P. glaucum*. The remaining chromosomes from *P. purpureum* parental (genome A'B) presented marks only in the centromeric and pericentromeric regions (Fig. 2c). However, when the probe with DNA of *P. purpureum* was used in chromosomes of the triploid hybrid the marks were observed mainly in centromeric and pericentromeric regions, but some chromosomes appearing almost totally marked (Fig. 1d).

The differences in marking pattern observed in the triploid hybrid, mainly between genomes A and A', could be explained by the presence of two genomes (A' and B) in the same probe. The observations evidence the changes arising from interspecific hybridization in *P. purpureum* genomes. Once combined in a polyploid hybrid nucleus, extensive reorganization may rapidly occur in the parental diploid genomes, both intra and intergenomically (Soltis and Soltis 1999; Chen et al. 2006). The rapid intergenomic rearrangements in polyploids in relation to the diploid progenitors have been demonstrated in allohexaploid F1 hybrids of *Avena sativa* (Leitch and Bennett 1997), in hexaploids of wheat (Nelson et al. 1995) and soybean (Shoemaker 1996), as well as in triploid hybrids embryos of *Pennisetum* (Campos 2007).

Besides the existing homeology among genomes A, A' and B, the utilization of GISH in the genomes of *P. glaucum*, *P. purpureum* and interspecific hybrid enable to verify the differences in chromosomes size, and also chromosomes number of these species. Analyzing cells of the interspecific hybrid, the difference in size between the parental chromosomes becomes evident, with those of *P. glaucum* being larger (Fig. 2c). It can also be observed that the total length of *P. purpureum* chromosomes did not increase proportionally in relation to those of *P. glaucum* (Table 1 and Fig 2a and b), and that the chromosomes of genome B do not differ significantly in size in relation to genome A' (Table 2 and Fig. 2a). These differences in size and chromosome number between the two species reflect their evolutionary history.

The evolutionary tendency among true grasses, which have a common and recent origin, is that the most derived species have emerged after reduction of the number and increase of the size of chromosomes in relation to the ancestors (Avdulov 1931;

Table 2. Total length (μm) of monoploid complement in genomes A, A' and B for each genotype.

Genotype	Total length of the genome A	Total length of the genome A'	Total length of the genome B
<i>P. purpureum</i>	-	23,8	23,95
<i>P. glaucum</i>	29,51	-	-
Triploid hybrid	17,63	10,54	12,44

Bennetzen 2007; Crepet and Feldman 1991; Martel et al. 2004; Paterson et al. 2004; Stebbins 1956, 1971). This tendency applies to the genus *Pennisetum*, whose common ancestral pattern presented the basic number of chromosomes $x = 9$. Furthermore, the evolutionary pattern inside the genus follows the same tendency, this way, it is observed that the ancestor of the species from section *Pennicillaria*, to which *P. purpureum* and *P. glaucum* belong, presented basic number of chromosomes $x = 7$. In this sense, analyzing the phylogeny of the genus *Pennisetum* presented by Martel et al. (2004) and the information on chromosome size and homeology of genomes A, A' and B observed in the present work, it can be inferred that the species *P. purpureum* and *P. glaucum* have concomitantly diverged from the common ancestor. The origin of *P. purpureum* occurred at the interspecific hybridization event, combining the genome A of the ancestor with genome B of a second, still unknown. Therefore, genome A' could be considered a subgenome of the ancestor A due to genomic and structural changes that occurred during evolution. On the other hand, the species from the primary genic pool of *P. glaucum* have diverged from the common ancestor through increase of chromosome size, probably by increment in the genic sequences. The increase in chromosome size in this species could be explained, as described by Poncet et al. (2002), by duplication of some genes as consequence of domestication syndrome. The presence of non-homologous recombination observed among the chromosomes of *P. glaucum* reinforces the hypothesis of genic duplication as part of the differentiation in this species (Jauhar 1970; Martel et al. 2004).

This hypothesis, presented for the evolution and divergence of *P. purpureum* and *P. glaucum* from the common ancestor, may be further reinforced by the differences observed in relation to the size of chromosomes from genomes A, A' and B, as shown in Table 2. Analyzing the size of the monoploid complement of genome A, it can be verified that it is 24% larger in relation to the length of the chromosomes of genome A' of *P. purpureum*. Considering that the genomes A and A' have evolved from an ancestor genome A, the difference in chromosome size could be related to genic duplication in *P. glaucum* and to genomic rearrangements observed in the allotetraploid hybrid *P. purpureum*. Rearrangements and loss of genomic sequences are common events after hybridization (Kellis et al. 2004), as observed in this study by comparison between different genomes combined in the triploid hybrid (Table 2). In this case, a reduction of 60% of genome A can be observed in the hybrid in relation to the parental *P. glaucum*, along with 44% and 52% reduction of genome A' and B, respectively, in relation to the parental *P. purpureum*.

In this work, GISH confirmed the homeology among genomes A, A' and B and enabled the identification and distribution of the homeologous regions in the chromosomes. Moreover, the GISH markings were able to separate the different genomes, leading the comparison on the size of the chromosomes in each of these three genomes. This distinction of the different genomes confirmed the occurrence of rearrangements after interspecific hybridization, especially when the synthetic triploid hybrid was analyzed, and prove the allotetraploid origin of *P. purpureum*. It also show that genomes A' and B have chromosomes similar in size. In evolutionary terms, the results reinforce that the genomes A and A' have diverged from an ancestral genome A by increase of chromosome size in *P. glaucum* and rearrangements and/or deletions in *P. purpureum*. The reorganizations occurring in the ancestral genome A during evolution have generated the subgenome A' of *P. purpureum*.

Acknowledgements

The authors would like to thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq, the Fundação de Amparo à Pesquisa do Estado de Minas Gerais – FAPEMIG, and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES, for financial support to the research and granting of scholarship.

References

- Andrade-Vieira LF (2010) Genomic behavior in hybrids of Napier grass and pearl millet (*Pennisetum* sp. Schum., Poaceae). PhD Thesis, Doctor in Genetics and Plant Breeding, Federal University of Lavras, Lavras, MG, 119 pp. [In Portuguese]
- Andrade-Vieira LF, Reis GB, Torres GA, Oliveira AR, Brasileiro-Vidal AC, Pereira AV, Davide LC (2013) Biparental Chromosome Elimination in Artificial Interspecific Hybrids *Pennisetum purpureum* Schum and *P. glaucum* (L.) R. Br. *Crop Science* 53: 1–8. doi: 10.2135/cropsci2013.03.0155
- Avdulov NP (1931) Karyo-systematische untersuchungen der familie gramineen. *Bulletin of Applied Botanic, of Genetic and Plant Breeding, Leningrad* 44: 119–123.
- Barbosa S, Davide LC, Pereira AV (2003) Cytogenetics of *Pennisetum purpureum* (Schumack) × *Pennisetum glaucum* L. hybrids and their parents. *Ciência e Agrotecnologia* 27: 26–35. doi: 10.1590/S1413-70542003000100003
- Bennetzen JL (2007) Patterns in grass genome evolution. *Plant Biology, Stuttgart* 10: 176–181. doi: 10.1016/j.pbi.2007.01.010
- Campos JMS (2007) Obtaining hexaploids hybrids and genomic analysis of *Pennisetum* sp. by flow cytometry. PhD Thesis, Doctor in Genetics and Plant Breeding, Federal University of Lavras, Lavras, MG, 115 pp. [In Portuguese]
- Crepet WL, Feldman GD (1991) The earliest remains of grasses in the fossil record. *American Journal Botany, Columbus* 78: 1010–1014. <http://www.jstor.org/stable/2445181>

- Chen ZJ, Ni Z (2006) Mechanisms of genomics rearrangements and gene expression changes in plant polyploids. *BioEssays*, Cambridge 28: 240–252. doi: 10.1002/bies.20374
- Chowdhury MKV, Smith RL (1988) Mitochondrial DNA variation in pearl millet and related species. *Theoretical and Applied Genetics* 76(1): 25–32. doi: 10.1007/BF00288827
- Dong F, Song J, Naess SK, Helgeson JP, Gebhardt C, Jiang J (2000) Development and applications of a set of chromosome-specific cytogenetic DNA markers in potato. *Theoretical and Applied Genetics* 101(7): 1001–1007. doi: 10.1007/s001220051573
- Hanna WW (1987) Utilization of wild relatives of pearl millet. Proceedings of the International Pearl Millet Workshop, April 7–11, 1986, 1. ICRISAT Center, Patancheru, 33–42. http://oar.icrisat.org/873/1/RA_00110.pdf
- Ingham LD, Hanna WW, Baier JW, Hannah LC (1993) Origin of the main class of repetitive DNA within selected *Pennisetum* species. *Molecular Genetics* 238(3): 350–358. doi: 10.1007/BF00291993
- Jauhar PP, Hanna WW (1998) Cytogenetics and genetics of pearl millet. *Advances in Agronomy*, San Diego 64: 1–26. doi: 10.1016/S0065-2113(08)60501-5
- Jauhar PP (1981) Cytogenetics of pearl millet. *Advances in Agronomy*, San Diego 34: 407–479. <http://eprints.icrisat.ac.in/id/eprint/6572>
- Jauhar PP (1970) Chromosomes behavior and fertility of the raw and ‘evolved’ synthetic tetraploids of pearl millet, *Pennisetum typhoides* Stapf and Hubb. *Genetica* 41(1): 407–424. doi: 10.1007/BF00958922
- Jauhar PP (1968) Inter- and intra-genomal chromosome pairing in an inter-specific hybrid and its bearing on the basic chromosome number in *Pennisetum*. *Genetica* 39(1): 360–370. doi: 10.1007/BF02324476
- Jiang J, Gill BS, Wang GL, Ronald PC, Ward DC (1995) Metaphase and interphase fluorescence in situ hybridization mapping of the rice genome with bacterial artificial chromosomes. *Genetics* 92(10): 4497–4491. <http://www.pnas.org/content/92/10/4487>
- Kellis M, Birren BW, Lander ES (2004) Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature* 428(6983): 617–624. doi: 10.1038/nature02424
- Leitch IJ, Bennett MD (1997) Polyploidy in angiosperms. *Trends Plant Science* 12: 470–476. doi: 10.1016/S1360-1385(97)01154-0
- Martel E, Poncet V, Lamy F, Siljak-Yakovlev S, Lejeune B, Sarr A (2004) Chromosome evolution of *Pennisetum* species (Poaceae): implications of ITS phylogeny. *Plant Systematics and Evolution* 249: 139–149. doi: 10.1007/s00606-004-0191-6
- Martel E, De Nay D, Siljak-Yakovlev S, Brown S, Sarr A (1997) Genome size variation and basic chromosome number in pearl millet and fourteen related *Pennisetum* species. *Journal of Heredity* 88: 139–143. doi: 10.1093/oxfordjournals.jhered.a023072
- Nelson JC, Deynze AE, Sorrells ME, Autrique E, Lu YH, Merlino M, Atkinson M, Leroy P (1995) Molecular mapping of wheat. Homeologous group 2, *Genome* 38: 516–524. doi: 10.1139/g95-067
- Pantulu JV (1967) Pachytene pairing and meiosis in the F1 hybrid of *Pennisetum typhoides* and *P. purpureum*. *Cytologia* 32: 532–541. doi: 10.1508/cytologia.32.532

- Paterson AH, Bowers JE, Chapman BA (2004) Ancient polyploidization predating divergence of the cereals, and its consequences for comparative genomics. *Proceedings of the National Academy of Sciences, Washington*, 101(26): 9903–9908. doi: 10.1073/pnas.0307901101
- Poncet V, Martel E, Allouis S, Devos KM, Lamy F, Sarr A, Robert T (2002) Comparative analysis of QTLs affecting domestication traits between two domesticated × wild pearl millet (*Pennisetum glaucum* L., Poaceae) crosses. *Theoretical Applied Genetics* 104(6–7): 965–975. doi: 10.1007/s00122-002-0889-1
- Renno JF, Mariac C, Poteaux C, Bezancon G, Lumaret R (2001) Haplotype variation of cp-DNA in the agamic grass complex *Pennisetum* section *Brevivalvula* (Poaceae). *Heredity* 86(5): 537–544. doi: 10.1046/j.1365-2540.2001.00854.x
- Robert T, Khalfallah N, Martel E, Lamy F, Poncet V, Allinne C, Remigereau M-S, Rekima S, Leveugle M, Lakis G, Siljak-Yakovlev S, Sarret A (2011) *Pennisetum*. In: Kole C (Ed.) Springer-Verlag. *Wild Crop Relatives: Genomic and Breeding Resources, Millets and Grasses*, Berlin Heidelberg, 217–255. doi: 10.1007/978-3-642-14255-0_13
- Sethi GS, Kalia HR, Ghai BS (1970) Cytogenetical studies of three interspecific hybrids between *Pennisetum typhoides* Stapf and Hubb. and *P. purpureum* Schumach. *Cytologia* 35(1): 96–101. doi: 10.1508/cytologia.35.96
- Shoemaker R, Olson T, Kanazin V (1996) Soybean genome organization: evolution of a legume genome. In: Gustafson JP, Flavell RB (Eds) Plenum Press. *Genomes of Plants and Animals: 21st Stadler Genetics Symposium*, New York, 139–150.
- Soltis DE, Soltis PS (1999) Polyploidy: recurrent formation and genome evolution. *Tree* 14(9): 348–352. doi: 10.1016/S0169-5347(99)01638-9
- Sree Rangasamy SR (1972) Cytological studies on diploid and polyploid taxa of the genus *Pennisetum* Rich. *Genetica* 43(2): 257–273. doi: 10.1007/BF00123633
- Stapf O, Hubbard CE (1934) *Pennisetum*. In: Prain D (Ed.) *Flora of Tropical Africa*, Part 6. Reeve & Co, Ltd, Ashford, Kent, 954–1070.
- Stebbins GL (1971) *Chromosomal evolution in higher plants*. E. Arnold, London, 216 pp.
- Stebbins GL (1956) Cytogenetics and evolution in the grass family. *American Journal of Botany* 43(10): 890–905. doi: 10.2307/2439006
- Techio VH, Davide LC, Pereira AV (2006) Meiosis in elephant grass (*Pennisetum purpureum*), pearl millet (*Pennisetum glaucum*) (Poaceae, Poales) and their interspecific hybrids. *Genetics and Molecular Biology* 29(2): 353–362. doi: 10.1590/S1415-47572006000200025
- Techio VH, Davide LC, Pereira AV (2005) Genomic analysis in *Pennisetum purpureum* and *P. glaucum* hybrids. *Caryologia* 8(1): 28–33. doi: 10.1080/00087114.2005.10589428
- Techio VH, Davide LC, Pereira AV, Bearzoti E (2002) Cytotaxonomy of some species and of interspecific hybrids of *Pennisetum* (Poaceae, Poales). *Genetics and Molecular Biology* 25(2): 203–209. doi: 10.1590/S1415-47572002000200014